AHC Media

Tuesday September 29, 2015

Ambulatory Surgery Centers: Infection Control Standards & Safe Injection Practices from CMS

The information provided in AHC Media Webinars does not, and is not intended to constitute medical or legal advice. Opinions, references and links provided by our speakers are provided for your convenience and do not represent our endorsement of such opinions, products or services.

Speaker

- Sue Dill Calloway RN, Esq. CPHRM, CCMSCP, CCMSCP
- AD, BA, BSN, MSN, JD
- President of Patient Safety and Education Consulting
- 5447 Fawnbrook Lane
- Dublin, Ohio 43017
- 614 791-1468
- sdill1@columbus.rr.com

Objectives

- Explain the necessity for the ASC in-house policy on safe injection practices.
- Describe how to best communicate these mandatory infection control standards and safe injection practices to new-employees during their on-boarding/orientation training.
- Explain new and revised standards, regulations, and laws put forth by CMS, TJC and the federal government.
- Evaluate compliance requirements and penalties.

You Don’t Want One of These

Introduction to the CMS CfCs on Infection Control

Conditions for Coverage (CFC)

- CMS published **Conditions For Coverage** (CfC) for Ambulatory Surgical Centers or ASC
- Similar to Conditions of Participation or CoPs published for hospitals
- ASC facilities that accept Medicare payment
  - Must follow the CfC regulations for all patients
  - Not just for Medicare or Medicaid patients
  - Don’t forget the Infection Control Worksheet

ASC Interpretive Guidelines

- CMS posted many revisions since May 15, 2009
- Current manual is dated April 1, 2015
  - First publishes in Federal Register
  - Next publishes the interpretive guidelines
  - Some standards have a survey procedures which are like the test questions to the test
- Renumbered the tag numbers and 151 pages (Q tag numbers 001-267)
  - Infection control worksheet (Exhibit 351) not attached to interpretive guidelines but form is available off the CMS website and periodically updated
- Available on CMS website[1]

[1]www.cms.gov/Regulations-and-Guidance/Guidance/Manuals/downloads/som107_exhibit_351.pdf

2 Policy memos at www.cms.hhs.gov/SurveyCertificationGenInfo/PMSR/list.asp#TopOfPage

Subscribe to the Federal Register Free



http://listserv.access.gpo.gov/cgi-bin/wa.exe?SUBED1=FEDREGTOC-L&A=1

CMS ASC CFC Manual

State Operations Manual
Appendix L - Guidance for Surveyors: Ambulatory Surgical Centers

Table of Contents
(Rev. 137, 04-01-15)



Transmittals for Appendix L

Part I - Ambulatory Surgical Center Survey Protocol

Introduction

Regulatory and Policy References

Tasks in the Survey Protocol

Task 1 – Off-Site Survey Preparation
Task 2 – Entrance Activities
Task 3 – Information Gathering/Investigation
Task 4 – Preliminary Decision Making and Analysis of Findings
Task 5 – Exit Conference
Task 6 – Post-Survey Activities

www.cms.gov/Regulations-and-Guidance/Guidance/Manuals/downloads/som107_Appendixtoc.pdf

CMS Rewrites ASC Manual in 2009

DEPARTMENT OF HEALTH & HUMAN SERVICES
Centers for Medicare & Medicaid Services
7500 Security Boulevard, Mail Stop S2-12-25
Baltimore, Maryland 21244-1850

www.cms.hhs.gov/SurveyCertificationGenInfo/downloads/SCLetter09_37.pdf

Center for Medicaid and State Operations/Survey and Certification Group

Ref: S&C-09-37

DATE: **May 15, 2009**

TO: State Survey Agency Directors

FROM: Director
Survey and Certification Group

SUBJECT: State Operations Manual (SOM) Appendix L, Ambulatory Surgical Centers (ASC) Comprehensive Revision

Memorandum Summary

- New ASC Conditions for Coverage (CfCs) take effect on May 18, 2009.
- SOM Appendix L has been revised:
  - Part 1, Survey Protocol has been added.
  - The Interpretive Guidelines have been extensively updated to reflect the new and revised CfCs, as well as to provide more detailed guidance for existing CfCs. Feedback on this guidance is welcome; as needed, further refinements will be made in the future.
- ASPEN Tags for ASC surveys have been revised and renumbered. ASPEN surveyor kits with the new tags and regulatory text will be available shortly after May 18th.

On November 18, 2008 the Centers for Medicare & Medicaid Services (CMS) adopted the Hospital Outpatient Prospective Payment System final rule (73 FR 68502). The final rule included revisions to the ASC CfCs in 42 CFR 416.2 – 416.52 that take effect on May 18, 2009. Among the regulatory changes are the following:

Original ASC Infection Control Worksheet

Attachment 2

Exhibit 351

INFECTION CONTROL SURVEYOR WORKSHEET

Instructions: The following is a list of items that must be assessed during the on-site survey, in order to determine compliance with the infection control Condition for Coverage. Items are to be assessed primarily by surveyor observation, with interviews used to provide additional confirming evidence of observations. In some cases information gained from interviews may provide sufficient evidence to support a deficiency citation.

The interviews and observations should be performed with the most appropriate staff person(s) for the items of interest (*e.g.*, the staff person responsible for sterilization should answer the sterilization questions).

A minimum of one surgical procedure must be observed during the site visit, unless the ASC is a low volume ASC with no procedures scheduled during the site visit. The surveyor(s) must identify at least one patient and follow that case from registration to discharge to observe pertinent practices. For facilities that perform brief procedures, e.g., colonoscopies, it is preferable to follow at least two cases.

When performing interviews and observations, any single instance of a breach in infection control would constitute a breach for that practice.

Citation instructions are provided throughout this instrument, indicating the applicable regulatory provision to be cited on the Form CMS 2567 when deficient practices are observed.

PART 1 - ASC CHARACTERISTICS

1) ASC name: _____

2) Address: _____

_____

State: _____

3) 10-digit CMS Certification Number: _____

4) What year did the ASC open for operation? _____

5) Please list date(s) of site visit: _____ (mm/dd/yyyy) to _____ (mm/dd/yyyy)

6) What was the date of the most recent previous federal (CMS) survey: _____ (mm/dd/yyyy)

7) Does the ASC participate in Medicare via accredited "deemed" status? 1 ☐ YES 2 ☐ NO

7a) If YES, by which CMS-recognized accreditation organization? (Check only ONE):

1 ☐ Accreditation Association for Ambulatory Health Care (AAAHC)
2 ☐ American Association for Accreditation of Ambulatory Surgery Facilities (AAAASF)

Conditions for Coverage (CFC)

- All CMS manuals found at website[1]
- Appendix **L** in the State Operations Manual Section 1832 of SSA ASC must meet quality and safety standards
- They are **Q** standards and Infection Control section starts at Q-0240
- Every ASC should have a copy of this
- Consider placing it on your intranet or making it available on all units

[1] http://www.cms.hhs.gov/manuals/downloads/som107_Appendicestoc.pdf

Location of CMS Hospital CoP Manuals

Medicare State Operations Manual

Appendix

- Each Appendix is a separate file that can be accessed directly from the SOM Appendices Table of Contents, as applicable.
- The appendices are in PDF format, which is the format generally used in the IOM to display files. Click on the red button in the "Download" column to see any available file in PDF.
- To return to this page after opening a PDF file on your desktop, use the browser "back" button. This is because closing the file usually will also close most browsers

CMS Hospital CoP Manuals new address

www.cms.hhs.gov/manuals/downloads/som107_Appendixtoc.pdf

App. No.	Description	PDF File
A	Hospitals	2,185 KB
AA	Psychiatric Hospitals	606 KB

I	Life Safety Code	140 KB
J	Intermediate Care Facilities for Persons With Mental Retardation	1,810 KB
K	Comprehensive Outpatient Rehabilitation Facilities	202 KB
L	Ambulatory Surgical Services Interpretive Guidelines and Survey Procedures	263 KB
M	Hospice	720 KB
N	Pharmaceutical Service Requirements in Long-Term Care Facilities	Deleted
P	Survey Protocol for Long-Term Care Facilities	929 KB
PP	Interpretive Guidelines for Long-Term Care Facilities	1,440 KB
Q	Determining Immediate Jeopardy	326 KB
R	Resident Assessment Instrument for Long-Term Care Facilities	38 KB
S	Mammography Suppliers	Deleted

Sample Page from ASC CfC Manual

Q-0240
(Rev. 56, Issued: 12-30-09, Effective/Implementation: 12-30-09)

§416.51 Condition for Coverage – Infection control

The ASC must maintain an infection control program that seeks to minimize infections and communicable diseases.

Interpretive Guidelines: §416.51

This regulation requires the ASC to maintain an active program for the minimization of infections and communicable diseases. The National Institute of Allergy and Infectious Diseases (NIAID) defines an infectious disease as a change from a state of health to a state in which part or all of a host's body cannot function normally because of the presence of an infectious agent or its product. An infectious agent is defined by the NIAID as a living or quasi-living organism or particle that causes an infectious disease, and includes bacteria, viruses, fungi, protozoa, helminthes, and prions. NIAID defines a communicable disease as a disease associated with an agent that can be transmitted from one host to another. (See NIAID website glossary)

The ASC's infection control program must:

CMS ASC Infection Control Worksheet



Updated Infection Control Worksheet

- CMS issued survey memo on June 26, 2015
- Announced revisions to the infection control worksheet
- Revised to bring the worksheet into compliance with accepted standards of practices
- Also to reflect currently released guidance
- Also to clarify questions that were being asked of the past worksheet

CMS Memo Revising IC Worksheet

DEPARTMENT OF HEALTH & HUMAN SERVICES
Centers for Medicare & Medicaid Services
7500 Security Boulevard, Mail Stop C2-21-16
Baltimore, Maryland 21244-1850

CENTERS FOR MEDICARE & MEDICAID SERVICES

Center for Clinical Standards and Quality/Survey & Certification Group

Ref: S&C: 15-43-ASC

DATE: **June 26, 2015**

TO: State Survey Agency Directors

FROM: Director
Survey and Certification Group

SUBJECT: Advanced Copy - Update to Ambulatory Surgical Center (ASC) Infection Control Surveyor Worksheet (ICSW)

Memorandum Summary

- *ASC Infection Control Surveyor Worksheet Revisions*: The Centers for Medicare & Medicaid Services (CMS) has made minor revisions to the Infection Control Surveyor Worksheet, Exhibit 351 of the State Operations Manual (SOM) for assessing compliance with the Medicare ASC Infection Control Condition for Coverage (CfC).
- *Change:* Revisions were made to bring the worksheet into alignment with current accepted standards of practice; reflect recently released guidance; and improve the clarity of certain questions. The worksheet is used by State and Federal surveyors on all survey activity in ASCs when assessing compliance with the infection control CfC.

Revised IC Worksheet 17 Pages

Exhibit 351

***Ambulatory Surgical Center (*ASC*)* INFECTION CONTROL SURVEYOR WORKSHEET**

(Rev: 142, Issued: 07-17-15, Effective: 07-17-15, Implementation: 07-17-15)

Name of State Agency or AO (please specify) _____

Instructions: The following is a list of items that must be assessed during the on-site survey, in order to determine compliance with the infection control Condition for Coverage. Items are to be assessed primarily by surveyor observation, with interviews used to provide additional confirming evidence of observations. In some cases information gained from interviews may provide sufficient evidence to support a deficiency citation.

The interviews and observations should be performed with the most appropriate staff person(s) for the items of interest (*e.g.*, the staff person responsible for sterilization should answer the sterilization questions). A minimum of one surgical procedure must be observed during the site visit. The surveyor(s) must identify at least one patient and follow that case from registration to discharge to observe pertinent practices. For facilities that perform brief procedures, e.g., colonoscopies, it is preferable to follow at least two cases. When performing interviews and observations, any single instance of a breach in infection control would constitute a breach for that practice.

Citation instructions are provided throughout this instrument, indicating the applicable regulatory provision to be cited on the Form CMS-2567 when deficient practices are observed.

PART 1 – ASC CHARACTERISTICS

1. ASC Name _____

2. Address, State and Zip Code _____
Address

CMS ASC Infection Control Worksheet

The following are sections discussed in the worksheet:

- Infection control program requirements
  - Must follow nationally recognized standards
- Qualified infection preventionist
  - Enough hours to direct program
- System to identify infections related to procedures
- Reporting system to state health agency
- Training of staff in orientation and annually

CMS ASC Infection Control Worksheet

- Hand hygiene requirements
  - Direct care givers do not wear artificial fingernails
- Safe injection practices
- Single use devices, sterilization and high level disinfection
  - Chemical indicator in every load and weekly biological indicator for each sterilizer and every load of implantable items
- Immediate use steam sterilization (IUSS)
- Environmental infection control

The CMS Infection Control Pilot

- When CMS first piloted the IC worksheet in 2008 in 67 ASFs and current version 7-17-15
- 68% had lapses in at least one IC category
- 18% has lapses in three or more of the five categories
  - Handling of blood glucose monitoring equipment, safe injection practices, equipment reprocessing and handling, hand hygiene, and environmental cleaning

CMS Survey Memos on Infection Control



CMS Policy and Memo Website

- Check the CMS website for updates
  - Consider doing this every month especially the survey and certification memos
  - Assign responsibility to one person
  - Check below website once a month
- Website includes information on CMS Survey and Certification memoranda, guidance, clarifications and instructions to State Survey Agencies and CMS Regional Offices

Source: http://www.cms.hhs.gov/SurveyCertificationGenInfo/PMSR/list.asp#TopOfPage

CMS Policy & Memo to States

Policy & Memos to States and Regions

CMS Survey and Certification memoranda, guidance, clarifications and instructions to State Survey Agencies and CMS Regional Offices. www.cms.gov/Medicare/Provider-Enrollment-and-Certification/SurveyCertificationGenInfo/Policy-and-Memos-to-States-and-Regions.html

Show entries: 10

Filter On:

Title	Memo #	Posting Date	Fiscal Year
Final Rule: SNF Medicare FY 2016 Payments, Quality Reporting, Value-Based Purchasing and Staffing Data Collection Requirements – Informational Only	15-49-NH	2015-08-07	2015
Publication of Medicare and Medicaid Programs; Reform of Requirements for Long-Term Care Facilities; Proposed Rule (CMS-3260-P) – Informational Only	15-46-NHs	2015-07-17	2015
Medication-Related Adverse Events in Nursing Homes	15-47-NH	2015-07-17	2015
Intermediate Care Facilities for Individuals with Intellectual Disabilities (ICF/IID) Appendix J, Part II - Clarifications to the Interpretive Guidance at Tag W187 for §483.430(d)(3)	15-48-ICF/IID	2015-07-17	2015
Advanced Copy - Update to Ambulatory Surgical Center (ASC) Infection Control Surveyor Worksheet (ICSW)	15-43-ASC	2015-06-26	2015
Use of Portable Reverse Osmosis (RO) Units and Block Carbon	15-44-ESRD	2015-06-26	2015

Example of a Survey Memo Informed Consent

Ref: S&C-11-21-ASC

DATE: **April 8, 2011**

TO: State Survey Agency Directors

FROM: Director
Survey and Certification Group

SUBJECT: Clarifications to the Ambulatory Surgical Center (ASC) Interpretive Guidelines, State Operations Manual Appendix L – Obtaining consent before observing surgical procedures

Memorandum Summary

ASCs may not limit surveyor ability to observe patient care and surveyors are not required to:

- Sign documents or provide proof of vaccinations in order to observe a surgical procedure.
- Obtain the consent of the operating physician prior to observing a surgical procedure.

Surveyors will:

- Obtain patient's consent.
- Conduct themselves in a manner that protects patient safety and that does not interfere with the operating physician or procedure.

It has come to our attention that some ASCs are attempting to require surveyors satisfy certain

CRE and ERCP Survey Memo

DEPARTMENT OF HEALTH & HUMAN SERVICES
Centers for Medicare & Medicaid Services
7500 Security Boulevard, Mail Stop C2-21-16
Baltimore, Maryland 21244-1850

CMS
CENTERS FOR MEDICARE & MEDICAID SERVICES

Center for Clinical Standards and Quality/Survey & Certification Group

Ref: S&C-15-32 Hospitals/CAHs/ASCs

DATE: **April 3, 2015**

TO: State Survey Agency Directors

FROM: Director
Survey and Certification Group

SUBJECT: Alert Related to Outbreaks of Carbapenem-Resistant *Enterobacteriaceae* (CRE) during gastrointestinal endoscopy, particularly Endoscopic Retrograde Cholangiopancreatography (ERCP)

Memorandum Summary

- ***Situation:*** Recent newspaper articles, medical publications, and adverse event reports associate multidrug-resistant bacterial infections caused by CRE with patients who have undergone ERCP. Duodenoscopes used to perform ERCP are difficult to clean and disinfect, even when manufacturer reprocessing instructions are followed correctly, and have been implicated in these outbreaks. The U.S. Food and Drug Administration (FDA) has issued a Safety Communication warning, with related updates, that the design of duodenoscopes may impede effective cleaning.

SAFE Injections in ASCs



Unsafe Injection Practices and Disease Transmission

Reuse of syringes combined with the use of single-dose vials for multiple patients undergoing anesthesia can transmit infectious diseases. The syringe does not have to be used on multiple patients for this to occur.



1. A clean syringe and needle are used to draw the sedative from a new vial.

2. It is then administered to a patient who has been previously infected with hepatitis C virus (HCV). Backflow into the syringe contaminates the syringe with HCV.

3. The needle is replaced, but the syringe is reused to draw additional sedative from the same vial for the same patient, contaminating the vial with HCV.

4. A clean needle and syringe are used for a second patient, but the contaminated vial is reused. Subsequent patients are now at risk for infection.

Source: www.**southernnevadahealthdistrict**.org

CDC

CMS Memo on Safe Injection Practices

- June 15, 2012 CMS issues a 7 page memo on safe injection practices
- Discusses the safe use of single dose medication to prevent healthcare associated infections (HAI)
- Notes new exception which is important especially in medications shortages
- General rule is that single dose vial (SDV)can only be used on one patient
- Will allow SDV to be used on multiple patients if prepared by pharmacist under laminar hood following USP 797 guidelines

Safe Injection Practices Memo

DEPARTMENT OF HEALTH & HUMAN SERVICES
Centers for Medicare & Medicaid Services
7500 Security Boulevard, Mail Stop C2-21-16
Baltimore, Maryland 21244-1850

CMS
CENTERS for MEDICARE & MEDICAID SERVICES

Office of Clinical Standards and Quality/Survey & Certification Group

Ref: S&C: 12-35-ALL

DATE: **June 15, 2012**

TO: State Survey Agency Directors

FROM: Director
Survey and Certification Group

SUBJECT: Safe Use of Single Dose/Single Use Medications to Prevent Healthcare-associated Infections

Memorandum Summary

- ***Under certain conditions, it is permissible to repackage single-dose vials or single use vials (collectively referred to in this memorandum as "SDVs") into smaller doses, each intended for a single patient:*** The United States Pharmacopeia (USP) has established standards for compounding which, to the extent such practices are also subject to regulation by the Food and Drug Administration (FDA), may also be recognized and enforced under §§501 and 502 of the Federal Food, Drug and Cosmetics Act (FDCA). These USP compounding standards include USP General Chapter 797, *Pharmaceutical Compounding - Sterile Preparations* ("USP <797>"). Under USP <797>, healthcare facilities may repackage SDVs into smaller doses, each intended for use with one patient. Among other things, these standards currently require that:
  - The facility doing the repackaging must use qualified, trained personnel to do so, under International Organization for Standardization (ISO) Class 5 air quality conditions within an ISO Class 7 buffer area. All entries into a SDV for purposes of repackaging under these conditions must be completed within 6 hours of the initial needle puncture.
  - All repackaged doses prepared under these conditions must be assigned and labeled with a beyond use date (BUD), based on an appropriate determination of contamination risk level in accordance with USP <797>, by the licensed healthcare professional supervising the repackaging process.
- ***Administering drugs from one SDV to multiple patients without adhering to USP <797>***

CMS Memo on Safe Injection Practices

- All entries into a SDV for purposes of repackaging must be completed with 6 hours of the initial puncture in pharmacy following USP guidelines
- Only exception of when SDV can be used on multiple patients
- Otherwise using a single dose vial on multiple patients is a violation of CDC standards
- CMS will cite hospital under the hospital CoP infection control standards since must provide sanitary environment
  - Also includes ASCs, hospice, LTC, home health, CAH, dialysis, etc.

CMS Memo on Safe Injection Practices

- Bottom line is you can not use a single dose vial on multiple patients
- CMS requires hospitals to follow nationally recognized standards of care like the CDC guidelines
- SDV typically lack an antimicrobial preservative
- Once the vial is entered the contents can support the growth of microorganisms
- The vials must have a beyond use date (BUD) and storage conditions on the label

CMS Memo on Safe Injection Practices

- Make sure pharmacist has a copy of this memo
- If medication is repackaged under an arrangement with an off site vendor or compounding facility ask for evidence they have adhered to 797 standards
- ASHP Foundation has a tool for assessing contractors who provide sterile products
- Go to www.ashpfoundation.org/MainMenuCategories/PracticeTools/SterileProductsTool.aspx
- Click on starting using sterile products outsourcing tool now

ASHP Foundation

FOSTERING SAFE AND EFFECTIVE MEDICATION USE

Contact Us | Donate Now | ASHPorg | SEARCH

HOME | ABOUT US | PROGRAM NEWS | LEADERSHIP | RESEARCH | ADVANCING PRACTICE | AWARDS | EDUCATION | SUPPORT THE ASHP FOUNDATION

• Print this page • Email this page

Advancing Practice

- Optimizing Antithrombotic Management: An Assessment Tool
- Bar Code Guide
- My Medicine List℠
- Outsourcing Sterile Products Preparation: Contractor Assessment Tool
- Pharmacy Practice Model Initiative

Outsourcing Sterile Products Preparation: Contractor Assessment Tool

Developed with support from PharMEDium Services, LLC

Now available!

Preparation of sterile parenteral products is a critical component of health-system pharmacy practice. For departments that choose to outsource the preparation of parenteral medications, this web-based tool can be used to evaluate proposals during the selection of an external organization that would provide parenteral product preparation services.

The assessment tool helps you evaluate each of these areas:

- Regulatory compliance
- Quality and patient safety measures
- Medication administration safety features
- Service excellence

Start using the Sterile Products Outsourcing Tool now!

www.ashpfoundation.org/MainMenuCategories/PracticeTools/SterileProductsTool.aspx

Safe Injection Practices

EMERGENCY MEDICINE PATIENT SAFETY FOUNDATION

Safe Injection Practices Patient Safety Brief
Emergency Medicine Patient Safety Foundation

By: Sue Dill Calloway RN MSN JD CPHRM
Ruth Carrico PhD RN FSHEA CIC

July 2012

Not All Vials Are Created Equal

SINGLE-DOSE OR MULTI-DOSE?

NOT ALL VIALS ARE CREATED EQUAL.

Dozens of recent outbreaks have been associated with reuse of single-dose vials and misuse of multiple-dose vials. As a result of these incidents, patients have suffered significant harms, including death. CDC and the One & Only Campaign urge healthcare providers to recognize the differences between single-dose and multiple-dose vials and to understand appropriate use of each container type.

This information can literally save a life.

ONEANDONLYCAMPAIGN.ORG

DO YOU PROVIDE TREATMENT FOR PATIENTS WITH CANCER?

PROTECT YOUR PATIENTS, YOURSELF, AND YOUR BUSINESS

Since 2002, at least nine serious infectious disease outbreaks have occurred in cancer clinics. These outbreaks involved unsafe injection practices, including the reuse of syringes. As a result, hundreds of patients became infected and thousands more required notification and testing for bloodborne pathogens.

REMEMBER! WHEN PREPARING MEDICATIONS AND INJECTIONS...

NEVER reuse these items:

Needles or syringes that have been used for any purpose

Vials with "single-dose vial" printed on the label

Saline bags

Intravenous tubing

ALWAYS follow aseptic technique* when:

Preparing any medication

Disinfecting a vial's septum

Accessing a central line

Injecting any medications

*Aseptic technique is used by health care workers to prevent the contamination of clean areas, equipment, and sterile medications. This will help prevent the spread of infection. Please refer to CDC's Basic Infection Control and Prevention Plan for Outpatient Oncology Settings for more information.

LEARN MORE ABOUT WAYS YOU CAN KEEP YOUR PATIENTS

Watch Award Winning Video

Safe Injection Practices - How to Do It Right

www.youtube.com/watch?v=6D0stMoz80k&feature=youtu.b

CMS Memo May 30, 2014

- CMS publishes 4 page memo on infection control breaches and when they warrant referral to the public health authorities
- This includes a finding by the state agency (SA), like the Department of Health, or an accreditation organization
  - TJC, DNV Healthcare, CIHQ, or AOA HFAP
- CMS has a list and any breaches should be referred
- Referral is to the state authority such as the state epidemiologist or State HAI Prevention Coordinator

Infection Control Breaches

DEPARTMENT OF HEALTH & HUMAN SERVICES
Centers for Medicare & Medicaid Services
7500 Security Boulevard, Mail Stop C2-21-16
Baltimore, Maryland 21244-1850

CMS
CENTERS FOR MEDICARE & MEDICAID SERVICES

Center for Clinical Standards and Quality/Survey & Certification Group

Ref: S&C: 14-36-All

DATE: **May 30, 2014**

TO: State Survey Agency Directors

FROM: Director
Survey and Certification Group

SUBJECT: Infection Control Breaches Which Warrant Referral to Public Health Authorities

Memorandum Summary

- ***Infection Control Breaches Warranting Referral to Public Health Authorities:*** If State Survey Agencies (SAs) or Accrediting Organizations (AOs) identify any of the breaches of generally accepted infection control standards listed in this memorandum, they should refer them to appropriate State authorities for public health assessment and management.

- ***Identification of Public Health Contact:*** SAs should consult with their State's Healthcare Associated Infections (HAI) Prevention Coordinator or State Epidemiologist on the preferred referral process. Since AOs operate in multiple States, they do not have to confer with State public health officials to set up referral processes, but are expected to refer identified breaches to the appropriate State public health contact identified at: http://www.cdc.gov/HAI/state-based/index.html

CMS Memo Infection Control Breaches

- If any of the listed breaches are observed, then will take appropriate enforcement action
- And will make the public health authority aware
  - Includes LTC, ASCs, hospice, hospitals, home health agencies, CAH, rural health clinics and dialysis facilities
- CDC is working closely with SA on HAI prevention
- List of breaches to be referred include:
- Using the same needle for more than one individual;

CMS Memo Infection Control Breaches

- Using the same (pre-filled/manufactured/insulin or any other) syringe, pen or injection device for more than one individual
- Re-using a needle or syringe which has already been used to administer medication to an individual to subsequently enter a medication container (e.g., vial, bag), and then using contents from that medication container for another individual
- Using the same lancing/fingerstick device for more than one individual, even if the lancet is changed

CMS Memo on Insulin Pens

- CMS issues memo on insulin pens on May 18, 2012
- Insulin pens are intended to be used on one patient only
- CMS notes that some healthcare providers are not aware of this
- Insulin pens were used on more than one patient which is like sharing needles
- Every patient must have their own insulin pen
- Insulin pens must be marked with the patient's name

Insulin Pens

DEPARTMENT OF HEALTH & HUMAN SERVICES
Centers for Medicare & Medicaid Services
7500 Security Boulevard, Mail Stop C2-21-16
Baltimore, Maryland 21244-1850

CMS
CENTERS for MEDICARE & MEDICAID SERVICES

Office of Clinical Standards and Quality/Survey & Certification Group

Ref: S&C: 12-30-ALL

DATE: May 18, 2012

TO: State Survey Agency Directors

FROM: Director
Survey and Certification Group

SUBJECT: Use of Insulin Pens in Health Care Facilities

www.cms.gov/Medicare/Provider-Enrollment-and-Certification/SurveyCertificationGenInfo/Policy-and-Memos-to-States-and-Regions.html

Memorandum Summary

Insulin Pen devices: The Centers for Medicare & Medicaid Services (CMS) has recently received reports of use of insulin pens for more than one patient, with at least one 2011 episode resulting in the need for post-exposure patient notification. These reports indicate that some healthcare personnel do not adhere to safe practices and may be unaware of the risks these unsafe practices pose to patients. **Insulin pens are meant for use by a single patient only.** Each patient/resident must have his/her own. Sharing of insulin pens is essentially the same as sharing needles or syringes, and must be cited, consistent with the applicable provider/supplier specific survey guidance, in the same manner as re-use of needles or syringes.

Background
Insulin pens are pen-shaped injector devices that contain a reservoir for insulin or an insulin cartridge. These devices are designed to permit self-injection and are intended for single-person use. In healthcare settings, these devices are often used by healthcare personnel to administer insulin to patients. Insulin pens are designed to be used multiple times by a single patient/resident, using a new needle for each injection. Insulin pens must never be used for more

CDC Has Flier on Insulin Pens

CDC CLINICAL REMINDER

Insulin Pens
Must Never Be Used for More than One Person

Summary

The Centers for Disease Control and Prevention (CDC) has become increasingly aware of reports of improper use of insulin pens, which places individuals at risk of infection with pathogens including hepatitis viruses and human immunodeficiency virus (HIV). This notice serves as a reminder that insulin pens must **never** be used on more than one person.

Background

Insulin pens are pen-shaped injector devices that contain a reservoir for insulin or an insulin cartridge. These devices are designed to permit self-injection and are intended for single-person use. In healthcare settings, these devices are often used by healthcare personnel to administer insulin to patients. Insulin pens are designed to be used multiple times, for a single person, using a new needle for each injection. Insulin pens must **never** be used for more than one person. Regurgitation of blood into the insulin cartridge can occur after injection [1] creating a risk of bloodborne pathogen transmission if the pen is used for more than one person, even when the needle is changed.

In 2009, in response to reports of improper use of insulin pens in hospitals, the Food and Drug Administration (FDA) issued an alert for healthcare professionals reminding them that insulin pens are meant for use on a single patient only and are not to be shared between patients [2]. In spite of this alert, there have been continuing reports of patients placed at risk through inappropriate reuse and sharing of insulin pens, including an incident in 2011 that required notification of more than 2,000 potentially exposed patients [3]. These events indicate that some healthcare personnel do not adhere to safe practices and may be unaware of the risks these unsafe practices pose to patients.

Recommendations

Insulin Pen Posters and Brochures Available

ONE NEEDLE, ONE SYRINGE, ONLY ONE TIME.
Safe Injection Practices Coalition
www.ONEandONLYcampaign.org

About the Campaign | Safe Injection Practices | Healthcare Provider Information | Patient Information | Campaign Resources | News | Contact Us

Insulin Pen Safety – One Insulin Pen, One Person

www.oneandonlycampaign.org/content/insulin-pen-safety

The Safe Injection Practices Coalition created an insulin pen poster and brochure for healthcare providers as a reminder that insulin pens and other injectable medications are meant for one person and should never be shared. PDFs of these educational materials are linked below:

Specific Materials for Safe Use of Insulin Pens – for Clinicians and Patients

- Poster
- Brochure

Click here to order free copies of these materials from the Centers for Disease Control and Prevention (CDC) (*publication numbers 22-1501 and 22-1503*).

Additional Resources

- VA Patient Safety Alert: Multi-Dose Pen Injectors (Department of Veterans Affairs, January 2013)

CMS Memo Humidity in the OR

- CMS issues memo related to the relative humidity (RH) on April 19, 2013
  - AORN use to say temperature maintained between 68-73 degrees and humidity between 30-60% in OR, PACU, cath lab, endoscopy rooms and instrument processing areas
  - Tag 60 and 101 in ASC manual
- CMS says if no state law can write policy or procedure or process to implement the waiver
- Waiver allows RH between 20-60%
- In anesthetizing locations- see definition in memo

Humidity in Anesthetizing Areas

DEPARTMENT OF HEALTH & HUMAN SERVICES
Centers for Medicare & Medicaid Services
7500 Security Boulevard, Mail Stop C2-21-16
Baltimore, Maryland 21244-1850

Center for Clinical Standards and Quality/Survey & Certification Group

Ref: S&C: 13-25-LSC & ASC

DATE: April 19, 2013

TO: State Survey Agency Directors

FROM: Director
Survey and Certification Group

SUBJECT: Relative Humidity (RH): Waiver of Life Safety Code (LSC) Anesthetizing Location Requirements; Discussion of Ambulatory Surgical Center (ASC) Operating Room Requirements

Memorandum Summary

- ***RH of ≥20 Percent Permitted in Anesthetizing Locations:*** The Centers for Medicare & Medicaid Services (CMS) is issuing a categorical LSC waiver permitting new and existing ventilation systems supplying hospital and critical access hospital (CAH) anesthetizing locations to operate with a RH of ≥20 percent, instead of ≥35 percent. We are also recommending that RH not exceed 60 percent in these locations.
- ***This Waiver Does Not Apply:***
  - When more stringent RH control levels are required by State or local laws and regulations; or
  - Where reduction in RH would negatively affect ventilation system performance.
- ***Hospitals & CAHs Must Elect to Use the Categorical Waiver:***
  - Individual waiver applications are not required, but facilities are expected to have written documentation that they have elected to use the waiver.
  - At the entrance conference for any survey assessing LSC compliance, a facility that has elected to use this waiver must notify the survey team.
- ***Ongoing Requirements:***
  - Facilities must monitor RH in anesthetizing locations and take corrective actions when needed to ensure RH remains at or above 20 percent.
- ***ASCs:*** ASCs are not subject to all of the same LSC requirements as hospitals, but are required, consistent with 42 CFR 416.44(a)(1), to maintain RH in operating rooms in accordance with nationally accepted guidelines.
- ***State Operations Manual (SOM) Appendices A, E, L, & W are being updated accordingly.***

Impact of Lowering the Humidity

- Lowering humidity can impact some equipment and supplies
- Can affect shelf life and product integrity of some sterile supplies including EKG electrodes
- Some electro-medical equipment may be affected by electrostatic discharge especially older equipment
  - Can cause erratic behavior of software and premature failure of the equipment
  - It can affect calibration of the equipment
- Follow the manufacturers instructions for use that explains any RH requirements

CMS Memo on Low Relative Humidity

DEPARTMENT OF HEALTH & HUMAN SERVICES
Centers for Medicare & Medicaid Services
7500 Security Boulevard, Mail Stop C2-21-16
Baltimore, Maryland 21244-1850

Center for Clinical Standards and Quality /Survey & Certification Group

Ref: S&C: 15-27-Hospital, CAH & ASC

DATE: **February 20, 2015**

TO: State Survey Agency Directors

FROM: Director
Survey and Certification Group

SUBJECT: Potential Adverse Impact of Lower Relative Humidity (RH) in Operating Rooms (ORs)

Memorandum Summary

- ***Information on OR RH*** is provided for Ambulatory Surgical Centers (ASCs) & Supplemental Information for Hospitals & Critical Access Hospitals (CAHs) Using the Categorical Waiver of Life Safety Code (LSC) Anesthetizing Location RH Requirements
  - The Association for the Advancement of Medical Instrumentation (AAMI) coordinated the release on January 5, 2015 of a Joint Communication of multiple healthcare-related organizations on how a RH of <30% in ORs may affect the performance of some sterile supplies and electro-medical equipment.
- ***S&C 13-25-LSC & ASC*** permits hospitals and CAHs to use a LSC categorical waiver to establish an RH level <35% in anesthetizing locations. Before electing or continuing to use this categorical waiver, hospitals and CAHs are expected to ensure that the humidity levels in their ORs are compatible with the manufacturers' instructions for use (IFUs) for the supplies and equipment used in that setting.
- ***ASCs do not require a categorical waiver*** in order to use a lower RH level in their ORs but also need to ensure they comply with the IFUs for their OR supplies and equipment.

Impact of Lowering the Humidity



American Hospital Association

ASHE

AHRMM

Quality Advisory

January 21, 2015

01-21-2015 Accessed ; https://www.magnetmail.net/actions/email_web_version.cfm?recipient_id=1331564405&message_id=8663272&user_id=AHA_8&group_id=1105177&jobid=25267573

NEW GUIDANCE ON HUMIDITY LEVELS IN THE OPERATING ROOM

THE ISSUE

A change in the standards regulating a hospital's physical environment in the operating room (OR) may conflict with the instructions for use on some equipment and supplies routinely used in surgery. To ensure patient safety during surgery, the AHA in collaboration with its personal membership groups, the American Society for Healthcare Engineering (ASHE) and the Association for Healthcare Resource & Materials Management (AHRMM), urge hospitals to examine their humidity levels in the OR and consider the effects on equipment and products used during surgery. This advisory and associated attachments will assist in your assessment.

BACKGROUND

Many safety codes and standards regulating the health care physical environment now require relative humidity levels in **ORs** (not other areas of the facility) to be at least 20 percent, a change from the 30 percent minimum humidity required by some previous editions of codes. The 20 percent threshold provides hospitals with flexibility during

Lowering Humidity Can Have Other Effects

RELATIVE HUMIDITY LEVELS IN THE OPERATING ROOM
JOINT COMMUNICATION TO HEALTHCARE DELIVERY ORGANIZATIONS
January 2015

This is an important communication to the multiple stakeholders in healthcare whose work touches sterile supplies and electro-medical equipment used in delivering care to patients. The subject is about how relative humidity (RH) levels lower than 30% can impact the integrity and functionality of some of these products, with a special emphasis on RH levels in the operating room (OR). The following professional organizations have collaborated in the development of this communication: Ambulatory Surgery Center Association (ASCA), American College of Clinical Engineering (ACCE), American Hospital Association (AHA), American Society for Healthcare Engineering (ASHE), American Society of Anesthesiologists (ASA), American Society of Heating, Refrigeration and Air Conditioning Engineers (ASHRAE), Association for Healthcare Resource & Materials Management (AHRMM), Association for the Advancement of Medical Instrumentation (AAMI), Association of periOperative Registered Nurses (AORN), Association of Surgical Technologists (AST), Health Industry Distributors Association (HIDA), and the International Association of Healthcare Central Service Materials Management (IAHCSMM).[1]

HHS Action Plan

- HHS estimated that HAIs incur nearly $20 billion in excess healthcare cost each year
  - Many are preventable
- Top priority of HHS now and CMS is under HHS
- Develop HHS Action Plan to Prevent HAIs
- Every infection preventionist (IP) should have a copy of this document
- HHS get a billion dollars to enforce IC and has a video every healthcare practitioner should see
  - Partnering to heal video at http://www.hhs.gov/partneringtoheal

[1]http://hhs.gov/ophs/initiatives/hai/index.html

Video on Preventing HAI

U.S. Department of Health & Human Services

HHS.gov

www.hhs.gov/ash/initiatives/hai/training/

Email Updates | Font Size

HHS Home > ASH Home > Initiatives > Healthcare-Associated Infections > Training

- ASH Home
- Key Personnel
- Regional Health Administrators
- Initiatives
  - Viral Hepatitis
  - Tobacco Control and Prevention
  - Healthcare-Associated Infections
    - Action Plan
    - National Targets and Metrics
    - Projects
    - Steering Committee
    - Events

Partnering to Heal:

TEAMING UP AGAINST HEALTHCARE-ASSOCIATED INFECTIONS

Partnering to Heal is a computer-based, video-simulation training program on infection control practices for clinicians, health professional students, and patient advocates.

The training highlights effective communication about infection control practices and ideas for creating a "culture of safety" in healthcare institutions to keep patients from getting sicker. Users assume the identity of the following five main characters and make decisions about preventing healthcare-associated infections (HAIs):

A Physician, Nathan Green, Director of a Hospital Post -op Unit, ready to start new prevention efforts in the unit;

A Registered Nurse, Dena Gray, working to learn effective communications skills that could make the difference for her patients;

State Specific ASC Regulations

- Many states have state specific regulations on freestanding ASCs
- Every ASC should be familiar with their state specific law
  - For example, Ohio has OAC 3701-83, amended 6/8/09[1]
  - Most all states have their ASC rules accessible on the internet
  - Provide citation to all your policies to both the CFC and state law provision

[1] http://www.odh.ohio.gov/rules/final/finalRules.aspx

CMS Resources

- CMS has website on Ambulatory Surgical Center[1]
- CMS has other resources on ASC
  - Medicare Learning Network (MLN) Spotlights
  - ASC Transmittals
  - Join ASC Mailing list
  - Join the Open door forum
  - ASC Internet only manuals
  - ASC Payment website[2]

[1] http://www.cms.hhs.gov/center/asc.asp

[2] http://www.cms.hhs.gov/ASCPayment/

Conditions for Coverage (CFC)

- Sign up for ASC Open Door Forums to stay informed on changes to ASC
- This is a free service by CMS
- Consider taping it and then administrator can listen and direct relevant sections to the appropriate staff
- Generally CMS does not have an outline of what is discussed in the open door forum
- CMS has an ASC website with resources

Source: http://www.cms.gov/Outreach-and-Education/Outreach/OpenDoorForums/index.html

Join ASC Listserv Policies and Information



CMS Ambulatory Surgical Center



Important Links

Coding and Billing

- Ambulatory Surgical Center (ASC) Payment
- ASC Payment System Questions and Answers
- CMS Recognized P-C IOLS and A-C IOLs [PDF, 154KB]
- New Technology Intraocular Lenses (NTIOLs)
- Electronic Billing & EDI Transactions
- HCPCS - General Information
- ICD-10 Coordination and Maintenance Committee
- ICD-10
- Claim Adjustment Reason and Remittance Advice Remark Codes The latest HIPAA codes are available at the Washington Publishing Company (WPC) Website.

CMS Manuals & Transmittals

-
- Transmittals
- Internet-Only Manuals (IOMs)

Health Care Consumer Initiatives

- Ambulatory Surgical Centers

Educational Resources

- MLN General Information
- MLN Educational Web Guides
- MLN Matters Articles (Search Tips)
- MLN Products

Contacts

- Quality Improvement Organizations (QIO)
- CMS Regional Offices
- (COBC)

How to Stay Informed

- Subscribe now to receive the weekly *MLN Connects Provider eNews* for the latest Fee-For-Service program information, event announcements, claims and pricer information, and MLN educational product updates.
- Open Door Forums
- Press Releases

CMS Ambulatory Surgical Center



www.cms.gov/Medicare/Medicare-Fee-for-Service-Payment/ASCPayment/index.html

National ASCA Association



CDC's Ebola Website

Centers for Disease Control and Prevention
CDC 24/7: Saving Lives. Protecting People.™

SEARCH

CDC A-Z INDEX

Ebola (Ebola Virus Disease)

Recommend Tweet Share

Language: English



About Ebola

2014 West Africa Outbreak

The 2014 Ebola epidemic is the largest in history, affecting multiple countries in West Africa. One imported case from Liberia and associated locally acquired cases in healthcare workers have been reported in the United States. CDC and partners are taking precautions to prevent the further spread of Ebola within the United States.

Latest CDC Outbreak Information

Updated October 16, 2014

What's New

October 19, 2014: When Caring for Suspect or Confirmed Patients with Ebola

October 17, 2014: Frequently Asked Questions about Dallas and Ohio Flights

October 17, 2014: Interim Guidance for Healthcare

Approved Accreditation Agencies

- There are four Medicare approved accreditation agencies for ASCs for deemed status
  - The Joint Commission (TJC)
  - American Association for Accreditation of Ambulatory Surgical Facilities (AAAASF)
  - Accreditation Association for Ambulatory Health Care (AAAHC)
  - American Osteopathic Association (AOA) or Healthcare Facility Accreditation program (HFAP)

CFC or the CoPs?

- Hospital elects for ASC to operate as a department of the hospital (CoPs) or decides to be paid as a ASC (CFC)
- Hospitals that operate a freestanding ASC as a department of the hospital follows the CMS CoPs
  - 510 pages long with tag numbers from 0001 to 1164
  - Radiology section changed in 2013 and no longer the same in CoP and CFC

[1] http://edocket.access.gpo.gov/2008/pdf/E8-26212.pdf

CMS Oversight of Infection Control

- CMS pilot show poor infection control standards
- This is why CMS gets a $50 million dollar grant and $10 million for ASCs so surveyors more knowledgeable
- Part 2 of the infection control worksheet is to provide detailed guidelines for assessing if the ASC has a sanitary environment
- CMS had mandatory training for ASC surveyors and heavy focus on the infection control standards and work sheet
  - Focus on instrument reprocessing also

Best Practices HAIs in Ambulatory Surgery

Best Practices HAIs in Ambulatory Surgery

- CMS requires ASCs to follow national standards of care and practice
- This is why best practices and evidenced based practice is important to implement in the ASC
- PPSA publishes use of evidence-based practices for prevention of HAIs in ambulatory surgery in 2013
- Infection control plan should be updated annually
- A structured education and competency program is implemented to integrate knowledge at the bedside
- Process and outcome measures are evaluated

Top Five Surgical Site Infections

Table 1. Top Five Infection Sites for Surgical Site Infections (SSIs), as Reported to the Pennsylvania Patient Safety Authority, March 2004 through July 2012

SURGICAL SITE	NO. OF SSIs	PROCEDURE WITH HIGHEST NO. OF SSIs (n)
Knee or shoulder joint	86	Arthroscopy/rotator cuff repair (41)
Ankle or foot	83	Bunionectomy (28)
Eye	79	Cataract surgery (40)
Abdomen	52	Hernia repairs (22)
Hand or wrist	49	Carpal tunnel surgery (15)

Best Practices HAIs in Ambulatory Surgery

PATIENT SAFETY AUTHORITY
An independent agency of the Commonwealth of Pennsylvania

Monitoring the Use of Evidence-Based Best Practices for Prevention of Healthcare-Associated Infections (HAIs) in Ambulatory Surgery

Monitoring compliance with best practices aimed at preventing HAIs is fundamental to improvement. This sample tool is designed to assess facility best-practice strategies and compliance in five categories: surveillance techniques, standardized educational programs, safe injections and point-of-care devices, sterilization and high-level disinfection, and environmental infection control. This tool provides assessment questions consistent with current infection control guidelines and may be adapted or revised as per facility policy. This tool can also be used to identify barriers to implementation of best practices at the bedside and to identify gaps in the effectiveness of the infection control program.

This tool can be used to assist an ambulatory surgical facility with assessment and process monitoring of the level of implementation of current evidence-based best practices. Questions in each category are assessed based on their consistency with the guidelines and on the following elements:

— Infection control goals are consistent with the infection control plan and are updated at least annually.
— Policies and procedures are up to date and reviewed annually.
— A structured educational/competency program is implemented to integrate knowledge into practice at the bedside.
— Sufficient resources are provided to enable staff compliance with policies.
— Process and outcome measures are evaluated and reviewed by the quality assurance and process improvement committee.
— Accountability is assigned for administrative support, resources, and implementation of best-practice strategies.

Application Considerations

— Facility self-assessment: This tool can be used by the facility to evaluate the current level of best-practice implementation.
— Clinical observation: Multiple items can be evaluated by clinical observation or simulation.
— Staff interview: Questions can be answered verbally, prior to clinical observation of actual practice.
— Record review: Ensure documentation of each category in the medical record, policy manual, and education records.

Sources

— A complete list of current guidelines for each category is located on the last page of the assessment tool.

Directions for Use

— In the implementation category cells next to each of the individual practices listed, enter the score for level of implementation (0, 1, or 2).
— In the remediation plan column, describe the action to improve scores less than 100%.

http://patientsafetyauthority.org/ADVISORIES/AdvisoryLibrary/Documents/2013_prepubtool_assessment.pdf

PATIENT SAFETY AUTHORITY
An independent agency of the Commonwealth of Pennsylvania

Ambulatory Surgical Facility Infection Control Practice Assessment

Scoring: 2 = 100% implemented; 1 = Partially implemented; 0 = Not implemented

	BEST PRACTICES	IMPLEMENTATION CATEGORIES: Infection control plan and goals	Policies and Procedures	Education/competency program	Resources	Accountability assigned	Monitoring of processes and outcomes assigned	REMEDIATION PLAN
I.	**SURVEILLANCE[1-4]**							
1	Postoperative surgical site infections (SSIs) are actively identified and tracked.							
2	There is written documentation confirming tracking of SSIs.							
3	The facility complies with state notifiable disease reporting requirements.							
4	SSI tracking information is shared with facility clinicians.							
II.	**STANDARDIZED EDUCATION[1-4]**							
1	A designated, licensed healthcare professional directs the infection control program.							
2	The healthcare professional directing infection control has received formal training in infection control and prevention.							
3	All facility staff receive ongoing, job-specific infection control education.							
4	The infection control education includes all licensed medical-							

Safe Injections and Point of Care Devices

	BEST PRACTICES	IMPLEMENTATION CATEGORIES						REMEDIATION PLAN
		Infection control plan and goals	Policies and Procedures	Education/competency program	Resources	Accountability assigned	Monitoring of processes and outcomes assigned	
2	The rubber septum is scrubbed with alcohol and dried prior to puncture.							
3	Injections are prepared in a clean area free from contamination or contact with blood, body fluids, or contaminated equipment.							
4	Needles, syringes, prefilled syringes, and insulin pen cartridges are used for only one patient.							
5	Single-dose and multidose does vials are always entered with a new needle and a new syringe, even for the same patient.							
6	Manufactured prefilled syringes, single-dose vials, ampoules, and intravenous solutions are used for only one time, for one dose, and for one patient.							
7	Medication administration tubing and connectors are used for only one patient.							
8	Multidose vials are dated when opened or punctured and discarded within 28 days or per manufacturer's instructions for an expiration date after opening, whichever is first.							
9	Multidose vials used on multiple patients are kept in a centralized, clean medication area and do not enter the immediate patient treatment area.							
10	Single-dose vials are discarded within one hour of opening or puncturing.							
11	Glucose meters are cleaned and disinfected after each patient.							

	BEST PRACTICES	IMPLEMENTATION CATEGORIES						REMEDIATION PLAN
		Infection control plan and goals	Policies and Procedures	Education/competency program	Resources	Accountability assigned	Monitoring of processes and outcomes assigned	
IV.	**STERILIZATION AND HIGH-LEVEL DISINFECTION[1-4,6]**							
1	Items are precleaned according to manufacturer's directions or evidence-based guidelines.							
2	Semicritical equipment, such as laryngoscope blades and respiratory therapy equipment, that contacts nonintact skin or mucosa is high-level disinfected.							
3	Critical equipment that enters sterile tissue or the vascular system, such as surgical instruments or implants, is sterilized.							
4	Medical devices and instruments are visually inspected for residual soil and recleaned if necessary before packaging and sterilization.							
5	Documentation of chemical, biological, and mechanical indicators is completed.							
6	Items are contained, handled, and stored in a manner to avoid compromised sterility.							
7	Sterile packages are inspected for integrity.							
8	Documentation is in place that chemicals for high-level disinfection are prepared, tested, and replaced according to manufacturer's instructions.							
9	Instruments undergoing high-level disinfection are disinfected for the appropriate length of time and at the appropriate temperature and allowed to dry as specified by manufacturer or evidence-based guidelines.							

Best Practices HAIs in Ambulatory Surgery

- PPSA had 733 events related to HAI surgical site infections and sterilization issues
  - From March 2004 to July 2012
  - Looked at adverse events classified as infections due to procedural complications, breaks in sterile technique, and sterilization problems due to equipment, supplies, or devices
  - 84% of the 733 infection control related events
  - Positive cultures in 43% of the 613 SSI event reports with Staph accounting for 59%
  - 16% due to sterilization problems with equipment, devices, surgical supplies or breaks in sterile technique

Ambulatory Surgical Facilities: Strategies to Integrate Basic Infection Control Concepts into Clinical Practice

Sharon Bradley, RN, CIC
Senior Infection Prevention Analyst
Pennsylvania Patient Safety Authority

ABSTRACT

In the last decade, the Centers for Medicare and Medicaid Services and the Centers for Disease Control and Prevention conducted investigations of national outbreaks of viral and bacterial infections and assessments of infection control practices in ambulatory surgical facilities (ASFs) across the country. Lax infection control practices were found to be widespread in ASFs, mostly due to clinicians misunderstanding or disregarding the standards of care. From March 2004 through July 2012, Pennsylvania ASFs reported 733 events related to healthcare-associated surgical site infections (SSI) and sterilization errors through the Pennsylvania Patient Safety Reporting System. Development

BACKGROUND

Since 2001, more than 150,000 patients in the United States have been exposed to viral and bacterial pathogens as a result of unsafe infection control and injection practices in ambulatory surgical facilities (ASFs), resulting in infectious outbreaks of meningitis, hepatitis, sacral abscesses, and joint infections.[1] These numbers may represent only a fraction of actual cases. Many outbreaks and sporadic transmissions associated with ASFs go unrecognized because patients may be treated for the infection in a physician office or a hospital, and some diseases (e.g., hepatitis) have long incubation periods.[2,3] Outbreaks have been identified in hospital and nursing home settings; however, outbreaks with the most widespread impact have occurred in outpatient settings.[2] Results of outbreak investigations by the Centers for Disease Control and Prevention (CDC)[3,4] and, more recently, national surveys by departments of health[5,6] point to lax infection control practices in ASFs, mostly due to clinicians' misunderstanding or disregarding the standards of care.[7]

In response to these outbreaks, in 2008, the Centers for Medicare and Medicaid Services (CMS) developed and piloted the Infection Control Surveyor Worksheet in ASFs in three states, assessing compliance with five categories of infection control.[8] Of the 68 ASFs inspected, 68% (n = 46) had lapses in at least one infection control category. Surveyors found that 18% (n = 12) had lapses in three or more of the following five categories: handling of blood glucose monitoring equipment, safe injection practices, equipment reprocessing and handling, hand hygiene, and environmental cleaning.[5]

In Pennsylvania, ASFs are required to report to the Pennsylvania Patient Safety Authority through its Pennsylvania Patient Safety Reporting System (PA-PSRS) adverse events classified as infections due to procedural complications, breaks in sterile technique, and sterilization problems due to equipment, supplies, or devices.[9,10] From March 2004

What They Found

- The flush step in the sterilization process of cleaning colonoscopes was omitted. All staff were reeducated to the endoscopy cleaning process.
- The nurse noticed 20 minutes into a case that the indicator in the arthroscopy instruments had not changed color to indicate that sterilization had occurred.
- It was noted that the antireflux valve was missing from the 24-hour tubing.
- At the end of the day the tech collecting autoclave sheets discovered that the set of instruction had been put in the autoclave and not run so unsterilized instruments were used on the patient.

The CMS CfCs for Infection Control





Infection Control Q-0240

- There are many sections on infection control in the CFC manual
- Infection control is a very important section and is being hit hard in the surveys
  - CMS got a $50 million dollar grant to enforce infection control standards and HHS 1 billion so surveyors are better educated
- Issue receiving interest after ASCs in Nevada were found to have multiple problems
  - 40,000 patients had to be tested who were seen over 4 yr period
  - The reuse of syringes having colonoscopies caused patients to get HIV, Hepatitis B and C

MMWR
Weekly
May 16, 2008 / 57(19);513-517

Acute Hepatitis C Virus Infections Attributed to Unsafe Injection Practices at an Endoscopy Clinic --- Nevada, 2007

On January 2, 2008, the Nevada State Health Division (NSHD) contacted CDC concerning surveillance reports received by the Southern Nevada Health District (SNHD) regarding two persons recently diagnosed with acute hepatitis C. A third person with acute hepatitis C was reported the following day. This raised concerns about an outbreak because SNHD typically confirms four or fewer cases of acute hepatitis C per year. Initial inquiries found that all three persons with acute hepatitis C underwent procedures at the same endoscopy clinic (clinic A) within 35--90 days of illness onset. A joint investigation by SNHD, NSHD, and CDC was initiated on January 9, 2008. The epidemiologic and laboratory investigation revealed that hepatitis C virus (HCV) transmission likely resulted from reuse of syringes on individual patients and use of single-use medication vials on multiple patients at the clinic. Health officials advised clinic A to stop unsafe injection practices immediately, and approximately 40,000 patients of the clinic were notified about their potential risk for exposure to HCV and other bloodborne pathogens. This report focuses on the six cases of acute hepatitis C identified during the initial investigation, which is ongoing; additional cases of acute hepatitis C associated with exposures at clinic A might be identified. Comprehensive measures involving viral hepatitis surveillance, health-care provider education, public awareness, professional oversight, licensing, and improvements in medical devices can help detect and prevent transmission of HCV and other bloodborne pathogens in health-care settings.

MRSA Safe Injection Practices July 13, 2012

www.cdc.gov/mmwr/preview/mmwrhtml/mm6127a1.htm?s_cid=mm6127a1_w

Centers for Disease Control and Prevention
MMWR
Morbidity and Mortality Weekly Report
Weekly / Vol. 61 / No. 27
July 13, 2012

Invasive *Staphylococcus aureus* Infections Associated with Pain Injections and Reuse of Single-Dose Vials — Arizona and Delaware, 2012

Transmission of life-threatening bacterial infections can occur when health-care personnel do not adhere to Standard Precautions and instead use medication in containers labeled as single-dose or single-use for more than one patient (*1*). This report summarizes the investigation of two outbreaks of invasive *Staphylococcus aureus* infection confirmed in 10 patients being treated for pain in outpatient clinics. In each outbreak, the use of single-dose or single-use vials (SDVs) for more than one patient was associated with infection transmission. In both investigations, clinicians reported difficulty obtaining the medication type or vial size that best fit their procedural needs. These outbreaks are a reminder of the serious consequences that can result when SDVs are used for more than one patient. Clinician adherence to safe injection practices, particularly when appropriately sized SDVs are unavailable, is important to prevent infection transmission. If SDVs must be used for more than one patient, full adherence to *U.S. Pharmacopeia* standards is critical to minimize the risks of multipatient use.

Pain Management Clinic — Arizona

On April 8, 2012, the Arizona Department of Health Services was notified of a patient with acute mediastinitis with blood and pleural fluid cultures positive for methicillin-resistant *Staphylococcus aureus* (MRSA). The report indicated this patient and two other patients with culture-confirmed invasive MRSA infections had undergone procedures recently

patients; two new syringes were used to withdraw 5 mL each from a 10 mL SDV of contrast medium (300 mgI/mL) and a 10 mL SDV of saline solution. The contents from each syringe then were transferred to the alternate vial, resulting in two 10 mL vials of diluted contrast solution, one for use in the morning and one reserved for the afternoon. Among patients receiving contrast on the day of the outbreak, six received injections from the morning vial and four from the afternoon vial. All of the patients with MRSA infections received diluted contrast from the afternoon vial.

The three patients with MRSA infections went to a local hospital 4–8 days after their outpatient pain remediation procedures. They required inpatient care for severe infections, including acute mediastinitis, bacterial meningitis, epidural abscess, and sepsis. Hospitalization ranged from 9 to 41 days, with additional long-term acute care required for one patient. The fourth recipient of diluted contrast from the afternoon vial was found deceased at home, 6 days after treatment at the clinic. The cause of death was reported as multiple-drug overdose; however, invasive MRSA infection could not be ruled out.

Samples from six unopened vials of contrast medium from the lot in use by the facility at the time of the outbreak were sent to CDC for analysis. No intrinsic contamination was identified using standard bacterial culture methods. Unopened vials of saline were not cultured because the lot in use at the time of the outbreak was not available and saline was routinely used

Unsafe Injection Practices Outbreaks

Acute Hepatitis C Virus Infections Attributed to Unsafe Injection Practices at an Endoscopy Clinic — Nevada, 2007

Outbreak of *Burkholderia cepacia* Bloodstream Infection at an Outpatient Hematology and Oncology Practice

Transmission of Hepatitis B and C Viruses in Outpatient Settings — New York, Oklahoma, and Nebraska, 2000–2002

Outbreak of *Serratia marcescens* Bloodstream and Central Nervous System Infections After Interventional Pain Management Procedures

75% of ASCs Cited in Infection Control

- July 2010 article states 75% of ASCs in one state received infection control citations
  - Reusing single use devices
  - Improper surgical attire
  - No infection control training (including training physicians)
  - OR nurse wearing ring
  - Doctor whose mask was not tied
  - Nurse picks up something off the OR floor and doesn't reglove
  - Instrument in OR on mayo stand after room cleaned

General Categories

Benchmarking
General Surgery
Accrediting/Quality
Anesthesia
Code/Bill/Reimburse
Building/Renovating

show all categories

Advertisement

OR EXCELLENCE

Harbor Beach Marriott Resort and Spa
Ft. Lauderdale, Fla.
Oct. 27 - 29, 2010

More Than 75% of Ill. Surgery Centers Cited in Infection Control Inspections

Dropped instruments and contaminated hands are among the lapses cited in state reports.

An OR nurse picks something off the floor mid-procedure, discards it, and then returns to work without washing her hands. A dropped surgical instrument is picked up, placed on a clean table and left there — even after the case is over and the room is cleaned. A surgeon enters the OR with his mask untied and dangling from his face.

These are just some of the infection control lapses discovered at more than three-quarter of Iliinois surgery centers inspected over the past year, according to state reports obtained by the Associated Press.

In 22 out of 29 centers surveyed, inspectors found clinics cleaning and reusing single-

Safe Injection Practices



- The surveyors will hit safe injection practices hard during the survey
- CMS has an infection control worksheet that everyone should be familiar with
- CDC has guidelines on single dose and multidose vials
  - Available at http://www.cdc.gov/ncidod/dhqp/injectionSafetyPractices.html
- Every ASC should have a safe injection practice P&P

CDC Injection Safety Website

CDC 10 Standards on Safe Injection Practices

Injection Safety
- CDC's Role
- CDC Statement
- Information for Providers
- Information for Patients
- Preventing Unsafe Injection Practices
- ▶**Safe Injection Practices**
- CDC Clinical Reminder: Spinal Injection Procedures
- Infection Prevention during Blood Glucose Monitoring and Insulin Administration
- Recent Publications
- Recent Meetings
- The One & Only Campaign

Related Links
- One & Only Campaign
- HICPAC
- 2007 Guideline for Isolation Precautions
- HHS Action Plan to Prevent HAIs

Injection Safety > Preventing Unsafe Injection Practices

Email page
Print page

Recommend 8 Tweet Share

Safe Injection Practices to Prevent Transmission of Infections to Patients

Download the complete 2007 Guideline for Isolation Precautions: Preventing Transmission of Infectious Agents in Healthcare Settings [PDF - 3.80 MB]

III.A.1.b. Safe Injection Practices The investigation of four large outbreaks of HBV and HCV among patients in ambulatory care facilities in the United States identified a need to define and reinforce safe injection practices 453. The four outbreaks occurred in a private medical practice, a pain clinic, an endoscopy clinic, and a hematology/oncology clinic. The primary breaches in infection control practice that contributed to these outbreaks were 1) reinsertion of used needles into a multiple-dose vial or solution container (e.g., saline bag) and 2) use of a single needle/syringe to administer intravenous medication to multiple patients. In one of these outbreaks, preparation of medications in the same workspace where used needle/syringes were dismantled also may have been a contributing factor. These and other outbreaks of viral hepatitis could have been prevented by adherence to basic principles of aseptic technique for the preparation and administration of parenteral medications 453, 454. These include the use of a sterile, single-use, disposable needle and syringe for each injection given and prevention of contamination of injection equipment and medication.

Contact Us:

Centers for Disease Control and Prevention
1600 Clifton Rd
Atlanta, GA 30333

800-CDC-INFO
(800-232-4636)
TTY: (888) 232-6348

New Hours of Operation
8am-8pm ET/Monday-Friday
Closed Holidays

cdcinfo@cdc.gov

Face Masks and Spinal Injections



Revised IC Worksheet 17 Pages

Exhibit 351

***Ambulatory Surgical Center (*ASC*)* INFECTION CONTROL SURVEYOR WORKSHEET**

(Rev: 142, Issued: 07-17-15, Effective: 07-17-15, Implementation: 07-17-15)

Name of State Agency or AO (please specify) _____

Instructions: The following is a list of items that must be assessed during the on-site survey, in order to determine compliance with the infection control Condition for Coverage. Items are to be assessed primarily by surveyor observation, with interviews used to provide additional confirming evidence of observations. In some cases information gained from interviews may provide sufficient evidence to support a deficiency citation.

The interviews and observations should be performed with the most appropriate staff person(s) for the items of interest (*e.g.*, the staff person responsible for sterilization should answer the sterilization questions). A minimum of one surgical procedure must be observed during the site visit. The surveyor(s) must identify at least one patient and follow that case from registration to discharge to observe pertinent practices. For facilities that perform brief procedures, e.g., colonoscopies, it is preferable to follow at least two cases. When performing interviews and observations, any single instance of a breach in infection control would constitute a breach for that practice.

Citation instructions are provided throughout this instrument, indicating the applicable regulatory provision to be cited on the Form CMS-2567 when deficient practices are observed.

PART 1 – ASC CHARACTERISTICS

1. ASC Name _____

2. Address, State and Zip Code _____
Address

Safe Injection Practices

II. Injection Practices (injectable medications, saline, other infusates)
Observations are to be made of staff *preparing* and *administering* medications and *performing* injections (e.g., anesthesiologists, certified registered nurse anesthetists, nurses).
Unless otherwise indicated, a "No" response to any question below must be cited as a deficient practice in relation to 42 CFR 416.51(a).

Practices to be Assessed	Was Practice Performed?	Manner of Confirmation
A. Needles are used for only one patient	○ Yes ○ No ○ N/A	○ Observation ○ Interview ○ Both

Practices to be Assessed	Was Practice Performed?	Manner of Confirmation
B. Syringes are used for only one patient	○ Yes ○ No ○ N/A	○ Observation ○ Interview ○ Both
C. The rubber septum on a medication vial is disinfected with alcohol prior to piercing.	○ *Yes* ○ *No* ○ *N/A*	○ *Observation* ○ *Interview* ○ *Both*

Safe Injection Practices IC Worksheet

II. Injection Practices (injectable medications, saline, other infusates)

Observations are to be made of staff preparing and administering medications and performing injections (e.g., anesthesiologists, certified registered nurse anesthetists, nurses).

Unless otherwise indicated, a "No" response to any question below **must** be cited as a deficient practice in relation to 42 CFR 416.51(a).

If unable to observe is selected, please clarify in the surveyor notes box why it was not observed and attempt to assess by means of interview or documentation review.

NOTE: Some types of infection control breaches, including some specific to medication administration practices, pose a risk of bloodborne pathogen transmission that warrant engagement of public health authorities. When management review confirms that a survey has identified evidence of one or more of the breaches described in S&C: 14-36-All, in addition to taking appropriate enforcement action to ensure the deficient Medicare practices are corrected, the SA should also make the responsible State public health authority aware of the identified breach.

Practices to be Assessed	Was Practice Performed?	*Surveyor Notes*
A. Needles are used for only one patient.	○ Yes ○ No ○ *Unable to observe*	

Safe Injection Practices IC Worksheet

- One needle and one syringe for only one patient
- Disinfect rubber septum with alcohol even if new
- Enter medication vial only with new needle
- Pre-drawn medication are labeled with date and time drawn, initial of person, name, strength, time, and BUD
- Bag of IV, tubing, and connectors used for only one patient
- Voluntary P&P that multi-dose vials are used for one patient only

Safe Injection Practices IC Worksheet

- Date multi-dose vials to expire in 28 days unless sooner by the manufacturer which is different from BUD
- Do not take multi-dose vials into patient care areas including the OR and anesthesia cart unless used on that patient then discarded
- Sharps are disposed of in puncture resistant containers
- Replace sharps containers when fill line is reached

CDC Resources Safe Injection Practices



CDC 10 Safe Injection Practices

- Information for Patients
- Preventing Unsafe Injection Practices
- **Safe Injection Practices**
  - CDC Clinical Reminder: Spinal Injection Procedures
- Drug Diversion
- Infection Prevention during Blood Glucose Monitoring and Insulin Administration
- Recent Publications
- Recent Meetings
- The One & Only Campaign
- Patient Notification Toolkit

Related Links

- CDC's HAI site
- 2007 Guideline for

Safe Injection Practices to Prevent Transmission of Infections to Patients

Download the complete 2007 Guideline for Isolation Precautions: Preventing Transmission of Infectious Agents in Healthcare Settings [PDF - 3.80 MB]

III.A.1.b. Safe Injection Practices The investigation of four large outbreaks of HBV and HCV among patients in ambulatory care facilities in the United States identified a need to define and reinforce safe injection practices 453. The four outbreaks occurred in a private medical practice, a pain clinic, an endoscopy clinic, and a hematology/oncology clinic. The primary breaches in infection control practice that contributed to these outbreaks were 1) reinsertion of used needles into a multiple-dose vial or solution container (e.g., saline bag) and 2) use of a single needle/syringe to administer intravenous medication to multiple patients. In one of these outbreaks, preparation of medications in the same workspace where used needle/syringes were dismantled also may have been a contributing factor. These and other outbreaks of viral hepatitis could have been prevented by adherence to basic principles of aseptic technique for the preparation and administration of parenteral medications 453, 454. These include the use of a sterile, single-use, disposable needle and syringe for each injection given and prevention of contamination of injection equipment and medication.

Whenever possible, use of single-dose vials is preferred over multiple-dose vials, especially when medications will be administered to multiple patients. Outbreaks related to unsafe

Control and Prevention
1600 Clifton Rd
Atlanta, GA 30333

800-CDC-INFO
(800-232-4636)
TTY: (888) 232-6348

Contact CDC-INFO

www.cdc.gov/injectionsafety/IP07_standardPrecaution.html

Advancing ASC Quality

- ASC Quality Collaboration has ASC tool kit for infection prevention
- Includes one on hand hygiene and safe injection practices
  - Excellent resources on safe injection practices
- Includes a basic and expanded version of the toolkit
  - These are available at http://www.ascquality.org/advancing_asc_quality.cfm

ASC Quality Collaboration Toolkits

ASC Quality Collaboration

Partners in ASC Quality

http://ascquality.org/advancing_asc_quality.cfm

ASC Quality Collaboration | Quality Measures and Guide | Quality Report

Advancing ASC Quality: ASC TIPs

- Hand Hygiene Toolkit
- Safe Injection Practices Toolkit
- Point of Care Devices Toolkit
- Environmental Infection Prevention Toolkit
- Single-Use Device Reprocessing Toolkit
- Endoscope Reprocessing Toolkit

Advancing ASC Quality

To support the ASC industry's focus on high quality care, the ASC Quality Collaboration is assembling **ASC Tools for Infection Prevention**, or **ASC TIPs**. Our goal is to make infection prevention resources readily accessible to ASCs by bringing them together in one location.

The following **ASC TIPs** are now available:

- Hand Hygiene Toolkit
- Safe Injection Practices Toolkit
- Point of Care Devices Toolkit
- Environmental Infection Prevention Toolkit
- Single-Use Device Reprocessing Toolkit
- Endoscope Reprocessing Toolkit
- Sterilization and High-Level Disinfection Toolkit



Safe Injection Practices: What CMS Surveyors Are Looking For

CMS surveyors use a worksheet to assess infection control practices during ASC surveys. The section of the worksheet used to assess safe injection practices is reproduced below. **Because this is the SAME TOOL a CMS surveyor will use to assess safe injection practices, it is also a useful SELF-ASSESSMENT tool for an ASC.**

Unless otherwise indicated, a "No" response to any question below will be cited as a deficient practice.

II. Injection Practices (injectable medications, saline, other infusates)
Observations are to be made of staff who prepare and administer medications and perform injections (e.g., anesthesiologists, certified registered nurse anesthetists, nurses).

Practices to be Assessed	Was Practice Performed?	Manner of Confirmation
A. Needles are used for only one patient	○ Yes ○ No ○ N/A	○ Observation ○ Interview ○ Both
B. Syringes are used for only one patient	○ Yes ○ No ○ N/A	○ Observation ○ Interview ○ Both
	○ Yes	○ Observation

CDC One & Only Campaign



www.oneandonlycampaign.org

Injection Safety Checklist

INJECTION SAFETY CHECKLIST

The following Injection Safety checklist items are a subset of items that can be found in the *CDC Infection Prevention Checklist for Outpatient Settings: Minimum Expectations for Safe Care*.

The checklist, which is appropriate for both inpatient and outpatient settings, should be used to systematically assess adherence of healthcare providers to safe injection practices. Assessment of adherence should be conducted by direct observation of healthcare personnel during the performance of their duties.

Injection Safety	Practice Performed?	If answer is No, document plan for remediation
Proper hand hygiene, using alcohol-based hand rub or soap and water, is performed prior to preparing and administering medications.	Yes No	
Injections are prepared using aseptic technique in a clean area free from contamination or contact with blood, body fluids, or contaminated equipment.	Yes No	
Needles and syringes are used for only one patient (this includes manufactured prefilled syringes and cartridge devices such as insulin pens).	Yes No	
The rubber septum on a medication vial is disinfected with alcohol prior to piercing.	Yes No	
Medication vials are entered with a new needle and a new syringe, even when obtaining additional doses for	Yes No	

Infection Control Q-0240

- The infection control section starts at Tag number 240
- **Standard** states that the ASC must maintain an infection control program that seeks to minimize infections and communicable diseases
- The ASC must have an active program to minimize infections and communicable diseases
- CMS has a definition of what is an infectious disease and infectious agent
  - Definition from The National Institute of Allergy and Infectious Diseases (NIAID)

Infection Control Q 240

- An **infectious disease** is defined as a change from a state of health to a state in which part or all of a host's body cannot function normally because of the presence of an infectious agent or its product
- An **infectious agent** is defined by the NIAID as a living or quasi-living organism or particle that causes an infectious disease, and includes bacteria, viruses, fungi, protozoa, helminthes, and prions. NIAID defines a communicable disease as a disease associated with an agent that can be transmitted from one host to another

Infection Control Q 240

- An infection control program must;
  - Provide a functional and sanitary environment for surgical services, to avoid sources and transmission of infections and communicable diseases
  - What constitutes a functional and sanitary environment is discussed in detail under tag 241
  - Be based on nationally recognized infection control guidelines
  - These organizations have many great resources

Requirements of an Infection Control Program

- Nationally recognized organizations include:
  - CDC or the Center for Disease Control
  - SHEA or The Society for Healthcare Epidemiology of American
  - APIC or Association for Professionals in Infection Control and Epidemiology
  - American Association of PeriOperative Registered Nurses or AORN
  - Institute for Healthcare Improvement or IHI
  - IDSA or the Infectious Diseases Society of America
  - Pa Patient Safety Authority or PaPSA

APIC Website

www.apic.or

APIC Spreading knowledge. Preventing infection.
Association for Professionals in Infection Control and Epidemiology

Media Membership Volunteering Sign In Search keywords

Consumers Professional Practice Education & Certification Resources Public Policy About

Spreading knowledge. Preventing infection.® APIC

Learn more

NEWS More Stories SPOTLIGHT

Measles: What you should know

Read More >

Low influenza vaccination rates among nursing home employees put residents at risk, study finds

Read More >

AORN

About AORN
Foundation
Career Center
Customer Service
Nurse Leaders
Educators
Industry Partners
Login
Join/Renew
AORN
Association of periOperative Registered Nurses
Topics of Interest
Search
Membership
Clinical Practice
Education
Events
Books & Publications
Get Involved
Advocacy
Have Questions About Surgical Attire?
Learn more about the new Guideline on Surgical Attire at AORN Central Booth 2121 at the Surgical Conference & Expo!
www.aorn.org
See implementation tools »
New Surgical Attire Module
AORN
Event Calendar
Product Catalog
Surgical Conference & Expo 2015 | March 7 - 11, 2015
Denver, CO
OR Executive Summit™ | March 8 - 10, 2015
Denver, CO
Leadership Development Summit™ | March 8 - 10, 2015
Denver, CO
View All Events
Annual Conference
Executive Summit
Research to Practice
OPEN
March 8-10
SOLUTIONS
GUIDELINES
25% New and Revised Content
One big show. One big opportunity. Get answers and solutions to your challenges, all in one place.
Learn, share ideas and solutions, and network with your peers at this exclusive conference.
Reduce Risk and Ensure Safest Patient Care with the 2015 Guidelines for Perioperative Practice.
ORNurseLink™
Connecting the periOperative Community
My Home
User Guide
NIFA
RNFA
Ask about our Early Bird Disc

AORN Guidelines for Perioperative Practice

Guidelines for Perioperative Practice

Introducing AORN's *Guidelines for Perioperative Practice* (previously titled *Perioperative Standards and Recommended Practices*). With a new name that more accurately reflects the content, this edition offers extensively revised and new, evidence-based guidelines for perioperative RNs and other members of the team to help standardize practice and promote patient and worker safety.

Available in several formats to suit your needs:

Print Edition

Easy hands-on access to AORN's evidence-rated guidelines in a traditional print format.
Order now

eBook Mobile App

Keep the most current information at your fingertips. Easily search, take notes and highlight your favorite content.
See all the benefits

eSubscription

The online edition is perfect for facilities to share among multiple users.
See all the benefits

CD

This CD's intuitive interface is ideal for teaching and presentations.
Order now

Packages

Save up to 15% on bundled formats and combine with other key publications.
See all packages

The 2015 edition features extensively revised and new guidelines for:

Institute for Healthcare Improvement IHI



SHEA Website

CDC Website


En Español
CDC
Centers for Disease Control and Prevention
CDC 24/7: Saving Lives, Protecting People™
SEARCH
CDC A-Z INDEX
Diseases & Conditions
Healthy Living
Travelers' Health
Emergency Preparedness
More CDC Topics
2015 Measles Cases and Outbreaks
Current information for ongoing Measles outbreak
www.cdc.gov
Recommend
Tweet
Share
What's New
Vital Signs
Secondhand Smoke
Learn about the unequal danger that
Outbreak
Ebola Update
CDC response to Ebola in United States and
Flu
Take 3: Actions To Fight The Flu
Learn CDC's recommended actions to
Feature
Prepare for Winter
Plan ahead and be prepared as cold

Requirements of an Infection Control Program

- Be directed by a designated health care professional with training in infection control
  - Infection preventionist should be qualified through education, training, experience, or certification
  - Certification is not required
- Be integrated into the ASC's QAPI program
- Be ongoing
- Include actions to prevent, identify and manage infections and communicable diseases; and

Infection Control Q 240

- Include a mechanism to immediately implement corrective actions and preventive measures that improve the control of infection within the ASC
  - Administrators need to listen to IP if they identify a problem
- The ambulatory care setting presents unique challenges for infection control
- Patients remain in common areas for a prolonged period of time such as the lobby
  - Some newer ASCs have lobbies with hepa filtration devices and lights that kill bacteria and viruses in the air

Infection Control Q 240

- Surgery and recovery rooms are turned around quickly
- May be hard to identify patients with infectious or communicable diseases
- Patients are in the ASC a short time which makes it hard sometimes to identify infections associated with the ASC
- Information has to be obtained after discharge and not directly linked if the patient was in the hospital

Infection Control Q 240

- Want to have physicians report back when postoperative infections are seen in the office
- The ASC can send an email to the primary care physician after discharge (See tag 245)
- Be sure to add this to the NPP (HIPAA Notice of Privacy Practice) and to the form the patient is signing so they are aware of this
- Considering asking the patient for their email address so you can send a follow up email
  - Send a follow up email asking if they have any problems including any postoperative infections

Infection Control 240

- ASCs must have a comprehensive infection control program
- In the past, poor infection control programs related to injections of medications, saline or other infusates in some ASCs have resulted in the transmission of communicable diseases
- Thousand of patients had to be tested for communicable diseases
  - This include hepatitis B and C and HIV

Infection Control 240

- The survey procedure discusses that one of the surveyors must complete the infection control worksheet
  - Remember every nurse, department director, and pharmacists should be familiar with the sections covered in the revised infection control worksheet
  - Infection preventionist should do training to physicians and staff
- However, each member of the survey team is to assess compliance with the ASC CfC requirements
  - Surveyors of ASC training session had heavy focus on infection control

Infection Control Sanitary Environment Q 241

- **Standard:** The ASC must provide a functional and sanitary environment for the provision of surgical services by adhering to professionally acceptable standards of practice (SOP).
- Necessary to avoid transmission of infections and communicable diseases
- All areas must be clean and sanitary
  - Include surgical prep area, PACU, and OR
- ASC must monitor environmental services (housekeeping) and maintenance activities

Policies for Sanitary Environment

- Ventilation and water quality control issues, including measures taken to maintain a safe environment during internal or external construction/renovation
- Maintaining safe air handling systems in areas of special ventilation, such as the OR
- Techniques for food sanitation if employee food storage and eating areas are provided

Policies for Sanitary Environment

- Techniques for cleaning and disinfecting environmental surfaces, carpeting, and furniture
- Techniques for disposal of regulated and non-regulated waste and
- Techniques for pest control
- These policies should be consistent with SOC such as CDC, AORN, APIC, SHEA etc.
  - CMS does not prescribe guidelines but ASC should document choices

CDC Guidelines for Environmental IC

Guidelines for Environmental Infection Control in Health-Care Facilities

Recommendations of CDC and the Healthcare Infection Control Practices Advisory Committee (HICPAC)

Prepared by
Lynne Sehulster, Ph.D.[1]
Raymond Y.W. Chinn, M.D.[2]
[1]*Division of Healthcare Quality Promotion*
National Center for Infectious Diseases
[2]*HICPAC member*
Sharp Memorial Hospital
San Diego, California

www.cdc.gov/mmwr/preview/mmwrhtml/rr5210a1.htm

The material in this report originated in the National Center for Infectious Diseases, James M. Hughes, M.D., Director; and the Division of Healthcare Quality Promotion, Steven L. Solomon, M.D., Acting Director.

Summary

The health-care facility environment is rarely implicated in disease transmission, except among patients who are immunocompromised. Nonetheless, inadvertent exposures to environmental pathogens (e.g., Aspergillus *spp. and* Legionella *spp.) or airborne pathogens (e.g.,* Mycobacterium tuberculosis *and varicella-zoster virus) can result in adverse patient outcomes and cause illness among health-care workers. Environmental infection-control strategies and engineering controls can effectively prevent these infections. The incidence of health-care--associated infections and pseudo-outbreaks can be minimized by 1) appropriate use of cleaners and disinfectants; 2) appropriate maintenance of medical equipment (e.g., automated endoscope reprocessors or hydrotherapy equipment); 3) adherence to water-quality standards for hemodialysis, and to ventilation standards for specialized care environments (e.g., airborne infection isolation rooms, protective environments, or operating rooms); and 4) prompt management of water intrusion into the facility. Routine environmental sampling is not usually advised, except for water quality determinations in hemodialysis settings and other situations*

AORN Toolkit Environmental Cleaning

www.aorn.org/toolkits/environmentalcleaning/

Survey Procedure Q 241

- Surveyor is instructed to use the specific questions on the infection control worksheet related to environmental infection control
- Instructed to observe how clean the ASC is including PACU, floors, horizontal surfaces, patient equipment, air inlets, supply, storage areas etc.
- Interview staff to determine whether cleaning and disinfection takes place and using appropriate materials

Survey Procedure Q 241

- Determine if has procedure for decontamination after gross spills of blood or other body fluids
- Determine whether sharps are disposed of properly
  - Needlestick Safety and Prevention Act
  - Part of the OSHA Bloodborne Pathogen Standard (29 CFR 1910.1030)
  - Have sharps containers at the bedside in all rooms
- Determine if the ASC re-uses devices that are marketed for single use

CMS IC Worksheet Part IV

- Tool for assessing compliance with Standard that the ASC provides a functional and sanitary environment by adhering to professionally acceptable standards of practice
- Observations are made of staff performing environmental cleaning including surgical technicians
- CMS also added the citation instructions
  - Unless otherwise indicated in the body of the worksheet , a “No” response to any question must be cited as a deficient practice

Environmental Infection Control

IV. Environmental Infection Control

Observations are to be made of staff performing **environmental cleaning (e.g., surgical technicians, cleaning staff, etc.)**

If unable to observe is selected, please clarify in the surveyor notes box why it was not observed and attempt to assess by means of interview or documentation review.

*Unless otherwise indicated, a "No" response to any question below **must** be cited as a deficient practice in relation to 42 CFR 416.51(a).*

Practices to be Assessed	Was Practice Performed?	*Surveyor Notes*
A. Operating rooms are cleaned and disinfected after each surgical or invasive procedure with an EPA-registered disinfectant	○ Yes ○ No ○ *Unable to observe*	
B. Operating rooms are terminally cleaned daily	○ Yes ○ No ○ *Unable to observe*	
C. *Environmental surfaces in patient care areas are cleaned and disinfected, using an EPA-registered disinfectant on a regular basis (e.g., daily), when spills occur and when*	○ Yes ○ No ○ *Unable to observe*	

Environmental Infection Control

- OR disinfected after each surgical or invasive procedure with an EPA registered disinfectant
- OR terminally cleaned daily
- Surfaces in patient care area cleaned and disinfected on a regular basis
  - And when spills occur or surfaces are visibly contaminated
- Have P&P to decontaminate gross spills of blood
- Will observe nurse doing finger stick for blood sugar
  - Hand hygiene done before and after with gloves

Environmental Infection Control

- Finger stick device must be totally disposable including lancet holding device
- Point-of-care device, like glucose meter, is disinfected after every use and according to manufacturer's instructions
- If manufacturer does not have cleaning instructions then must be limited to one patient use

Infection Control Program Q 242

- **Standard**: The ASC must maintain an ongoing program designed to prevent, control, and investigate infections and communicable diseases.
- In addition, the infection control and prevention program must include documentation that the ASC has considered, selected, and implemented nationally recognized infection control guidelines.

Infection Control Program Q 242

- Program must be ongoing
- Must prevent, control, and investigate infections and communicable diseases
- Must have an active surveillance program
  - Must cover patients and staff
  - Must include detection through collecting and analyzing the data
  - CDC has good surveillance resources
- Program must be based on national standards
  - Specify CDC, AORN, SHEA, APIC, in P&P, IC plan etc.

CDC Surveillance

CDC Monitoring HAI



I. Essential Elements of Surveillance[1, 2, 3]

A. Assess the population and identify those at greatest risk for the outcome or process of interest

1. Healthcare-associated infections (HAI) (outcomes)

2. Patient care practices aimed at preventing HAI (processes)

B. Select the outcome or process for surveillance

1. Examples of outcomes: HAI, infection or colonization with a specific organism, pyrogenic reaction or vascular access infection in hemodialysis patients, sharps injuries, etc.

2. Examples of processes: Central line insertion practices (CLIPs), surgical care processes (e.g., preoperative antimicrobial prophylaxis), medication errors, influenza vaccination rates, hepatitis B immunity rates, personnel compliance with protocols, etc.

3. Examples of other events: Occurrence of reportable diseases and conditions, communicable diseases in personnel, organisms or syndromes indicative of bioterrorist events, etc.

C. Determine observation time period

D. Choose the surveillance methodology (See Section II)

E. Monitor for the outcome or process using standardized definitions for all data

Elements of Effective State-Based Surveillance for Multidrug-Resistant Organisms Related to Healthcare-Associated Infections: Summary of Discussions from a Council of State and Territorial Epidemiologists and Centers for Disease Control and Prevention Experts Meeting



Jonathan Duffy, MD, MPH[a]; Dawn Sievert, PhD, MS[a]; Catherine Rebmann, MPH[a]; Marion Kainer, MD, MPH[b,c]; Ruth Lynfield, MD[b,d]; Perry Smith, MD[b,e]; Scott Fridkin, MD[a]

[a]Centers for Disease Control and Prevention
[b]Council of State and Territorial Epidemiologists
[c]Tennessee Department of Health
[d]Minnesota Department of Health
[e]New York State Department of Health

So What's In Your Infection Control Plan?

GETTING TO ZERO
INFECTION CONTROL AND PREVENTION (ICP) ACTION PLAN

Action Area One: Collaborative Approach: New Ways of Thinking

Key Issues: Tackling healthcare associated infections requires commitment from all levels of the organization and an enhanced local and system infrastructure committed to "getting to zero".

PRIORITY AREA	ACTION REQUIRED	MEASUREMENT OF SUCCESS	LEAD	PRIORITY	COMPLETION DATE
Infrastructure	1. Form a Hospital X-wide multidisciplinary Infection Control and Prevention (ICP) workgroup from inpatient and outpatient services, including Physician, Nursing, Pharmacy, Laboratory, Housekeeping, Facilities, Risk, Quality and Safety departments, etc. to: • Oversee the development and implementation of Hospital X's Infection Control and Prevention strategies • Monitor performance against the action plan • Review scorecard metrics (including all deaths associated with infections) • Serve as champions to facilitate intervention strategies	Formation of ICP workgroup with quarterly progress reports	VP Clinical Safety, Local Infectious Disease physician to co-chair ICP workgroup	HIGH	January 09
	2. Assess and recommend appropriate local structure to ensure accountability in meeting "getting to zero" goals	Distribution of draft recommendations for local accountability structure	VP Clinical Safety, VP Medical Services, Physician Leadership Council, Nursing Leadership Council	HIGH	February 09

PRIORITY AREA	ACTION REQUIRED	MEASUREMENT OF SUCCESS	LEAD	PRIORITY	COMPLETION DATE
Management and Communication of Critical Data and Information	1. Establish a process for thorough review and appropriate action steps related to the following: • Hospital X deaths or serious harm associated with infections, including mechanism to share action plans and findings Hospital X-wide to prevent reoccurrence. • Notification & inclusion of ICP leaders in renovation or new construction design • Incidences of infection control breakdown	Established review process and action plan drafted	Patient Safety Center with assistance of multidisciplinary ICP workgroup	HIGH	March 09
	2. Evaluate available ICP alert and reporting software to maximize ICP efficiency, documentation, screening and surveillance	Completed evaluation with recommendation of ICP alert and reporting software	Patient Safety Center with assistance of multidisciplinary ICP workgroup	HIGH	September 09
	3. Define metrics for: • Appropriate use of targeted antibiotics • MRSA • SSI • VAP • CLBSI • CA-UTI	Monthly report of metrics	Hospital X Pharmacy & Therapeutics Committee, Quality Management Department and Patient Safety Center	HIGH	July 09


IP Tools
upload. share. prevent.
LOOKING FOR RESOURCES AND A PLACE TO COLLABORATE WITH OTHER IPS?
IP TOOLS IS A RESOURCE FOR INFORMATION SHARING AMONG INFECTION PREVENTIONISTS.
Home
Tools & Resources
Help
Invite Others
GO
search...
Home
www.infectionpreventiontools.com/
Top 10 Downloads
Scope Reprocessing Competency Package (3712)
Risk Assessment (3105)
Sterile Processing Assessment Recommendation Collection (2952)
Sterile Processing Monitoring Checklist (2664)
UTI Catheter Related Infections Audit Tool (2575)
Infection Control Report - Data Summary for Reporting (2563)
Risk Assessment (2422)
IC Program Template and Checklists (1970)
UTI Bundle (1864)
Bed Bug POLICY (1812)
Forgot your Username or Password?
JOIN THE THOUSANDS OF USERS FROM 53 DIFFERENT COUNTRIES WHO ARE VISITING AND ACCESSING THESE TOOLS AND RESOURCES.
What is IP Tools?
Our Vision
Questions?
What is IP Tools?
IP Tools is devoted to the sharing of information among Infection Preventionists.
The goal is to enable this group of professionals to post documents and files they feel would be useful to others as well as download documents and files that have been posted by others.
Every download is free.
So, if you are ready to share some of your own tools and resources or you are ready to see what others have shared, click Tools and Resources.
Additional resources available at: www.publichealthtools.com
Why register?
Registration is important, it allows users access to all of

Infection Control Program Q 242

- ASC needs to select one or more sets of guidelines for the following key functions:
  - Maintain sanitary environment
  - Develop and implement IC activities for MS, employees, contract workers including nurses who are employed by the physician, housekeeping etc
  - Mitigate the risks associated with HAI
    - Correct name is healthcare-association infections

5 Key Functions of Infection Control

- Identifying infections
- Monitoring compliance with all policies, procedures, protocols and other infection control program requirements
  - Need to make sure all P&P are followed
  - Make sure staff are aware of all IC P&P
- Program evaluation and revision of the program, when indicated
  - ASC should consider doing this once a year and documenting when it is done

Infection Control Activities Q 242

- Evaluate staff immunization status for designated infectious diseases
  - Follow the recommendation of the CDC ACIP or Advisory committee on Immunization Practices
  - Recommendations at www.cdc.gov/vaccines/recs/acip/default.htm
  - See also www.flu.gov
- Need a policy regarding when the ASC is going to screen staff for infections likely to cause significant infectious diseases
  - Must include when staff are restricted from providing direct care to patients

Advisory Committee on Immunization Practices



CDC Flu and Vaccination Information

www.cdc.gov/flu/index.htm

Infection Control Activities Q 242

- P&P must also include when staff must remain away from the ASC all together
- Need P&P for reportable disease under state law
  - Every state has a list of communicable diseases that must be reported to the state department of health like MRSA or VRE
- New employees need orientation and regular update training in preventing and controlling HAI
  - Do not forget to provide infection control training to the physicians and LIPs also

Infection Control Activities 242

- Infection preventionist should also provide training on methods to prevent exposure and transmission of infections and communicable diseases
  - Hand hygiene is one of the most important ways to prevent infection
  - Studies show that compliance rates could be improved
- ASC needs methods to evaluate staff exposure to patients with infectious and communicable diseases
  - Nurse has needle stick injury and patient is HIV positive
  - Nurse exposed to patient who is diagnosed with TB

Mitigation of Risks Contributing to HAI

- The CDC defines an HAI as a localized or systemic condition resulting from an adverse reaction to the presence of an infectious agent or its toxin for purposes of surveillance
- There must be no evidence that the infection was present or incubating at the time of admission to the ASC
- HAI can be from an endogenous
  - Such as from skin, nose, mouth, GI trach, or vagina that are normally inhibited by microorganisms
  - Note the Jan 7, 2010 edition of NEJM

Mitigation of Risks Contributing to HAI 242

- This article found that one out of every seven patients have staph aureus in their nose
  - If culture positive can irrigate and add nasal drops and reduce surgical site infections
  - Recent study found one in four patient have positive nasal culture for staph
  - Jan 2010 NEJM also found that using chlorahexidine instead of betadine could reduce surgical site infections dramatically
- HAI can be from an exogenous source which means outside the body

Surgery Risk Mitigation Measures

- Examples of exogenous sources would include visitors, patient care staff, patient equipment, medical devices, or the health care environment
- Surgery related infection risk mitigation
  - Implement prophylaxis (antibiotics) to prevent surgical site infection
  - Must be given at the appropriate time (usually within one hour of the incision)
  - Must be discontinued appropriately after surgery
  - Do not forget if long case to look to see if patient needs rebolusing of the antibiotic

Surgery Risk Mitigation Measures 242

- Need to address aseptic techniques used in surgery including sterilization or high level disinfection of instruments
  - Sterilization of scopes and other equipment is very important and recent problems have made this an area that is hit hard during the survey
  - CMS issues memo on sterilization of equipment including flash sterilization (immediate use steam is new name)
  - AORN Perioperative Standards and Recommended Practices has standards on this now called Guidelines for PeriOperative Practices
  - Need to clean instruments with lap sponge, soak in enzymatic solution and then sterilize so no bioburden

IUSS Question in IC Worksheet

Practices to be Assessed	Was Practice Performed?
H. Is immediate-use steam sterilization (IUSS) performed on-site? ***If NO, skip to "High Level Disinfection Section"*** ***If YES, you must also assess the practices at questions "I - K":*** *(A "No" answer does not result in a citation)*	○ *Yes* ○ *No*
I. If IUSS is performed, all of the following criteria are met: • *Work practices ensure proper cleaning and decontamination, inspection, and arrangement of the instruments into the recommended sterilizing trays or other containment devices before sterilization.* • *Once clean, the item is placed within a container intended for immediate use. The sterilizer cycle and parameters used are selected according to the manufacturers' instructions for use for the device. container. and sterilizer.*	○ *Yes* ○ *No* ○ *Unable to observe* ○ *N/A*

IUSS Question in IC Worksheet

- If IUSS used, must ensure proper cleaning, decontamination, inspection, and arrangement of instruments in sterilizing trays
- One clean, item is placed in container for immediate use and placed on field in aseptic manner
- Must follow manufacturer instructions
- Sterilizer function is monitored such as mechanical, biologic, and chemical
  - Not a routine or frequent practice
- Not performed on implants, CJ instruments, etc.

Immediate Use Sterilization

- CMS issues a memo on immediate use sterilization or IUSS
  - Multiple society went together and named immediate use sterilization; AORN, AAMI, APIC, AAAHC, etc.
- CMS instructs hospitals to follow manufactures recommendation
- Not intended to be used to process items used at a later date
- Intended for immediate use so used during a procedure for which it was sterilized and in manner that minimizes exposure to air and other contaminates

Immediate Use Steam Sterilization IUSS

DEPARTMENT OF HEALTH & HUMAN SERVICES
Centers for Medicare & Medicaid Services
7500 Security Boulevard, Mail Stop C2-21-16
Baltimore, Maryland 21244-1850

Center for Clinical Standards and Quality /Survey & Certification Group

Ref: S&C: 14-44-Hospital/CAH/ASC

DATE: **August 29, 2014**

TO: State Survey Agency Directors

FROM: Director
Survey and Certification Group

SUBJECT: Change in Terminology and Update of Survey and Certification (S&C) Memorandum 09-55 Regarding Immediate Use Steam Sterilization (IUSS) in Surgical Settings

Memorandum Summary

- ***Change in Terminology: "Flash" Sterilization vs. IUSS:*** Nationally recognized organizations with expertise in infection prevention and control and instrument sterilization processes, and other professional organizations recommend abandoning the use of the term "flash" sterilization, which is now considered outmoded, and replacing it with the term "IUSS."

- ***Update of S&C Memorandum 09-55 Regarding Standards for Immediate Use Sterilization in Surgical Settings:*** This memo reiterates and updates information regarding nationally recognized infection prevention and control guidelines and professionally acceptable standards of practice with respect to immediate use sterilization and supersedes S&C Memorandum 09-55.

Introduction

Immediate-Use Steam Sterilization

www.aami.org/publication/standards/ST79_Immediate_Use_Statement.pdf

Immediate-Use Steam Sterilization

"Flash sterilization" has traditionally been used to describe steam sterilization cycles where unwrapped medical instruments are subjected to an abbreviated steam exposure time and then used promptly after cycle completion without being stored. This is in contrast to traditional "terminal sterilization" cycles, where instruments are sterilized within containers, wrappers, or primary packaging designed to maintain the instruments' sterility and allow the devices to be stored for later use. The term "flash" arose out of the abbreviated time of exposure of the unwrapped device.

Today, however, "flash sterilization" is an antiquated term that does not fully describe the various steam sterilization cycles now used to process items not intended to be stored for later use. Current guidelines may require longer exposure times and/or the use of single wrappers or containers designed to allow for aseptic transfer of an item to the point of use. The term "immediate-use steam sterilization" more accurately reflects the current use of these processes. The same critical reprocessing steps (such as cleaning, decontaminating, and transporting sterilized items) must be followed regardless of the specific sterilization cycle employed; a safe process does not include short-cuts or work-arounds.

"Immediate use" is broadly defined as the shortest possible time between a sterilized item's removal from the sterilizer and its aseptic transfer to the sterile field. Immediacy implies that a sterilized item is used during the procedure for which it was sterilized and in a manner that minimizes its exposure to air and other environmental contaminants. A sterilized item

Single Use, High Level Disinfection, Sterilization

III. Single Use Devices, Sterilization, and High Level Disinfection

Pre-cleaning must always be performed prior to sterilization and high-level disinfection

Sterilization must be performed for critical equipment (i.e., instruments and equipment that enter normally sterile tissue or the vascular system, such as surgical instruments)

High-level disinfection must be performed for semi-critical equipment (i.e., items that come into contact with non-intact skin or mucous membranes such as reusable flexible endoscopes, laryngoscope blades)

Observations are to be made of staff *performing* equipment reprocessing (e.g., surgical techs), unless these activities are performed under contract or arrangement off-site from the ASC.

Unless otherwise indicated, a "No" response to any question below **must** be cited as a deficient practice in relation to 42 CFR 416.51(a).

SINGLE-USE DEVICES

(Choose N/A if single-use devices are never reprocessed and used again) (Surveyor to confirm there is a contract or other documentation of an arrangement with a reprocessing facility by viewing it)

Practices to be Assessed		Was Practice Performed?	*Surveyor Notes*
A.	a. If single-use devices are reprocessed, they are devices that are approved by the FDA for reprocessing	○ Yes ○ No ○ N/A	

Surgery Risk Mitigation Measures 242

- Need to promote hand hygiene among staff
  - Follow CDC guidelines or the WHO hand hygiene guidelines
  - TJC has great document on how to measure hand hygiene adherence
- Need measures specific to prevention of organizations that are antibiotic resistant
  - Concerns because of MRSA and VRE and other super bugs
  - We need new antibiotics to fight there emerging infections

CDC Hand Hygiene Guidelines



www.cdc.gov/mmwr/PDF/rr/rr5116.pdf

Guideline for Hand Hygiene in Health-Care Settings

Recommendations of the Healthcare Infection Control Practices Advisory Committee and the HICPAC/SHEA/APIC/IDSA Hand Hygiene Task Force

CDC Posters

www.cdc.gov/handhygiene/Resources.html

World Health Organization | Patient Safety
A World Alliance for Safer Health Care



http://whqlibdoc.who.int/publications/2009/9789241597906_eng.pdf

WHO Guidelines on Hand Hygiene in Health Care

First Global Patient Safety Challenge
Clean Care is Safer Care

CDC Hand Hygiene Recommendations

- CDC published guidelines Oct 25, 2002 at www.cdc.gov/handhygiene/
- In CDC MMWR Recommendations and Reports,
- Report available at www.cdc.gov/mmwr/preview/mmwrhtml/rr5116a1.htm or go to www.cdc.gov,
- TJC published document in 2009 on Measuring Hand Hygiene Adherence: Overcoming the Challenges and this is an important document,
- Monitored during infection control tracer,

HAND HYGIENE ADHERENCE: OVERCOMING THE CHALLENGES

http://www.jointcommission.org/assets/1/18/hh_monograph.pdf

This monograph was authored by The Joint Commission in collaboration with the following organizations:

- The Association for Professionals in Infection Control and Epidemiology, Inc.
- The Centers for Disease Control and Prevention
- The Institute for Healthcare Improvement
- The National Foundation for Infectious Diseases
- The Society for Healthcare Epidemiology of America
- The World Health Organization World Alliance for Patient Safety

This monograph was supported in part by an unrestricted educational grant provided by GOJO Industries, Inc., Akron, Ohio

Surgery Risk Mitigation Measures 242

- ASC must measure specific safe practices for injecting medications and saline and other infusates
  - Recommend policy on safe injection practices with 10 CDC recommendation
  - Include single and multi-dose vials recommendations
- Germicides and disinfectants must be used in accordance with manufacturers instructions
- Appropriate use of facility and medical equipment
  - Includes hepa air filtration, UV lights, and other equipment to control the spread of infectious agents

AJIC special article

www.apic.org/AM/Template.cfm?Section=Position_Statements&Template=/CM/ContentDisplay.cfm&ContentID=15088

APIC position paper: Safe injection, infusion, and medication vial practices in health care

Susan A. Dolan, RN, MS, CIC,[a] Gwenda Felizardo, RN, BSN, CIC,[b] Sue Barnes, RN, BSN, CIC,[c] Tracy R. Cox, RN, CIC,[d] Marcia Patrick, RN, MSN, CIC,[e] Katherine S. Ward, RN, BSN, MPH, CIC,[f] and Kathleen Meehan Arias, MS, CIC[g]
Washington, DC

Outbreaks involving the transmission of bloodborne pathogens or other microbial pathogens to patients in various types of health care settings due to unsafe injection, infusion, and medication vial practices are unacceptable. Each of the outbreaks could have been prevented by the use of proper aseptic technique in conjunction with basic infection prevention practices for handling parenteral medications, administration of injections, and procurement and sampling of blood. This document provides practice guidance for health care facilities on essential safe injection, infusion, and vial practices that should be consistently implemented in such settings.

Key Words: Bloodborne pathogens; injection; infusion; medication vial practices; aseptic technique; parenteral medications; administration of injections; procurement of blood.

Copyright © 2010 by the Association for Professionals in Infection Control and Epidemiology, Inc. Published by Elsevier Inc. All rights reserved. (Am J Infect Control 2010;38:167-72.)

The transmission of bloodborne viruses and other microbial pathogens to patients during routine health care procedures continues to occur because of the use of unsafe and improper injection, infusion, and medication vial practices by health care professionals in the exposure of >100,000 individuals to viral hepatitis and the transmission of either hepatitis B virus (HBV) or hepatitis C virus (HCV) to more than 500 patients.[1-3] The unsafe practices used by health care personnel in these outbreaks can be categorized as (1)

Surgery Risk Mitigation Measures 242

- Educate patients, visitors and staff about infections and communicable diseases and methods to reduce transmission in the ASC and in the community
  - APIC has a number of free educational brochures
  - CDC has MRSA education both for staff and community MRSA
  - TJC has the speak up brochure series including five things you can do to reduce infections
  - CDC has a video on hand hygiene and standard precautions at www.cdc.gov/handhygiene/training/interactiveEducation/

Educational Brochures

http://www.apic.org/AM/Template.cfm?Section=Education_Resources&Template=/TaggedPage/TaggedPageDisplay.cfm&TPLID=91&ContentID=8738

APIC works to provide information to both the general public and healthcare professionals. The brochures on this page are regularly reviewed and updated as needed to insure that the information provided is current. These materials are available for you to download, copy and distribute free of charge. These pamphlets are intended to provide a general reference to each topic. No brochure can adequately diagnose a medical condition. If in doubt regarding your symptoms, please contact a healthcare professional.

- 10 tips for preventing the spread of infection
- Antibiotic Safety
- Chlamydia
- Companion Animals and Your Health
- Food Safety in Your Home
- Hand Hygiene for Consumers
- Hand Hygiene for Healthcare Workers
- Healthy Skin - An Ounce of Prevention
- Hepatitis A (HAV)
- Hepatitis B (HBV)
- Hepatitis C
- Hepatitis C (HCV)
- How to Dine Out Safely
- Los Hechos Sobre Chlamydia
- Meningococcal Meningitis
- Mold in Your Home
- Patient Safety - Protecting Yourself from Medical Errors
- Pneumonoccal Disease - Are You at Risk?
- Prevent Needlesticks
- Preventing Infection During Your Chemotherapy Treatment
- Preventing Ventilator Associated Pneumonia
- Respiratory Etiquette - An Important Cultural Change
- Respiratory Syncytial Virus
- Scarlet Fever
- Speak Up: Five Things You Can Do To Prevent Infection
- Traveling Overseas - Tips on How to Avoid Infection

www.cdc.gov/mrsa/mrsa_initiative/skin_infection/index.html

CDC Home

Centers for Disease Control and Prevention
Your Online Source for Credible Health Information

SEARCH

A-Z Index A B C D E F G H I J K L M N O P Q R S T U V W X Y Z #

National MRSA Education Initiative: Preventing MRSA Skin Infections

National MRSA Education Initiative

- About MRSA Skin Infections
- Consumer Educational Materials
- MRSA Photos
- Information for Healthcare Professionals
- Materials for Healthcare Professionals
- Treatment Algorithm
- Public Service Announcements
- Press Room

Text size: S M L XL

Email page
Print page
Bookmark and share

The goal of the National MRSA Education Initiative is to help Americans better recognize and prevent MRSA skin infections.

Why is this important? Recent data show that Americans visit the doctor approximately 12 million times each year to get checked for suspected Staph or MRSA skin infection.

The good news is that a few simple steps can prevent and reduce the spread of MRSA.

Contact Us:

Centers for Disease Control and Prevention
National MRSA Education Initiative
1600 Clifton Rd, MS A35
Atlanta, GA 30333

800-CDC-INFO
(800-232-4636)
TTY: (888) 232-6348
24 Hours/Every Day

cdcinfo@cdc.gov

Five Things You Can Do To Prevent Infection is supported by

American Hospital Association
www.hospitalconnect.com

Association for Professionals in Infection Control and Epidemiology, Inc.
www.apic.org

Centers for Disease Control and Prevention
www.cdc.gov

Infectious Diseases Society of America
www.idsociety.org

The Joint Commission
www.jointcommission.org

Society for Healthcare Epidemiology of America
www.shea-online.org

The Joint Commission is the largest health care accrediting body in the United States that promotes quality and safety.

Helping health care organizations help patients

The Joint Commission

SpeakUP™

5 Five Things You Can Do To Prevent Infection

www.jointcommission.org/GeneralPublic/Speak+Up/about_speakup.

Avoiding contagious diseases like the common cold, strep throat, and the flu is important to everyone. Here are five easy things you can do to fight the spread of infection.

1.

Clean your hands.

- Use soap and warm water. Rub your hands really well for at least 15 seconds. Rub your palms, fingernails, in between your fingers, and the backs of your hands.
- Or, if your hands do not look dirty, clean them with alcohol-based hand sanitizers. Rub the sanitizer all over your hands, especially under your nails and between your fingers, until your hands are dry.
- Clean your hands before touching or eating food. Clean them after you use the bathroom, take out the trash, change a diaper, visit someone who is ill, or play with a pet.

2.

Make sure health care providers clean their hands or wear gloves.

- Doctors, nurses, dentists and other health care providers come into contact with lots of bacteria and viruses. So before they treat you, ask them if they've cleaned their hands.
- Health care providers should wear clean gloves when they perform tasks such as taking throat cultures, pulling teeth, taking blood, touching wounds or body fluids, and examining your mouth or private parts. Don't be afraid to ask them if they should wear gloves.

3.

Cover your mouth and nose.

Many diseases are spread through sneezes and coughs. When you sneeze or cough, the germs can travel 3 feet or more! Cover your mouth and nose to prevent the spread of infection to others.

- Use a tissue! Keep tissues handy at home, at work and in your pocket. Be sure to throw away used tissues and clean your hands after coughing or sneezing.
- If you don't have a tissue, cover your mouth and nose with the bend of your elbow or hands. If you use your hands, clean them right away.

4.

If you are sick, avoid close contact with others.

- If you are sick, stay away from other people or stay home. Don't shake hands or touch others.
- When you go for medical treatment, call ahead and ask if there's anything you can do to avoid infecting people in the waiting room.

5.

Get shots to avoid disease and fight the spread of infection.

Make sure that your vaccinations are current—even for adults. Check with your doctor about shots you may need. Vaccinations are available to prevent these diseases:

- Chicken pox
- Measles
- Tetanus
- Shingles
- Mumps
- Diphtheria
- Hepatitis
- Meningitis
- Flu (also known as influenza)
- Whooping cough (also known as Pertussis)
- German measles (also known as Rubella)
- Pneumonia (*Streptococcus pneumoniae*)
- Human papillomavirus (HPV)

The goal of the Speak Up™ program is to help patients become more informed and involved in their health care.

CDC Hand Hygiene Resources and Video

www.cdc.gov/handhygiene/Patient_materials.html

CDC Home

Centers for Disease Control and Prevention

Your Online Source for Credible Health Information

SEARCH

A-Z Index A B C D E F G H I J K L M N O P Q R S T U V W X Y Z #

Hand Hygiene in Healthcare Settings

Hand Hygiene in Healthcare Settings

- Hand Hygiene Features
- Hand Hygiene Basics
- Guidelines
- Training
- **Patient Materials**
- Promotional Campaigns
- Measurement

Hand Hygiene in Healthcare Settings

Text size: S M L XL

Email page

Print page

Bookmark and share

Patient Materials

Hand Hygiene Saves Lives: Patient Admission Video

This video, available in English and Spanish, teaches two key points to hospital patients and visitors to help prevent infections: the importance of practicing hand hygiene while in the hospital, and that it is appropriate to ask or remind their healthcare providers to practice hand hygiene as well.

In the video, a nurse character named Gayle demonstrates hand hygiene using an alcohol-based hand sanitizer while emphasizing

Modeled after the video that airline passengers are required to view prior to take-off on a flight, this new video is intended to be shown to patients upon admission to the hospital. The goal is that the video

Get email updates

To receive email updates about Hand Hygiene, enter your email address:

What's this?

Submit

Identifying Infections 242

- ASC must monitor activities through out the entire facility to identify infection risks and communicable disease problems
  - Need to document the monitoring and tracking activities
- Surveillance should be done in accordance with recognized infection control surveillance activities such as the CDC National Healthcare Safety Network
- Need to monitor follow up of patients after discharge to gather information to see if they got an infection

CDC Home

Centers for Disease Control and Prevention

CDC 24/7: Saving Lives. Protecting People.™

NHSN
All CDC Topics
Choose a topic above
SEARCH

A-Z Index A B C D E F G H I J K L M N O P Q R S T U V W X Y Z #

National Healthcare Safety Network (NHSN)

www.cdc.gov/nhsn

CDC's National Healthcare Safety Network is the nation's most widely used healthcare-associated infection tracking system. NHSN provides facilities, states, regions, and the nation with data needed to identify problem areas, measure progress of prevention efforts, and ultimately eliminate healthcare-associated infections.

In addition, NHSN allows healthcare facilities to track blood safety errors and important healthcare process measures such as healthcare personnel influenza vaccine status and infection control adherence rates.

New Research

REDUCE MRSA Trial: Simple steps slash infection rates

Learn More»

1 2 3

Email page link
Print page

NHSN Login
Tips for navigating the new NHSN website [PDF - 1.6 MB]

Webinar Registration

The NHSN Biovigilance Component Team will be conducting a training webinar for the Hemovigilance Module.
Date: August 29, 2013
Time: 2:00 – 3:00 pm
Registration deadline August 26, 2013

Go to registration »

Continuing Education Opportunities

About NHSN

CDC's NHSN is the largest HAI reporting system in the U.S.

Data & Reports

See national and state reports using NHSN data

Guidelines and Recommendations

Review CDC HAI prevention guidelines

New to NHSN? Enroll Facility Here.

For first time facility enrollment.

Reporting & Surveillance Resources for Enrolled Facilities

Training, protocols, forms, support materials, analysis resources, and FAQs

Group Users

View resources for group users here.

Get email updates

To receive email updates about this page, enter your email address:

National Healthcare Safety Network NHSN

- CDC has great resources
- Includes an internet surveillance system that is secure
- Can monitor adverse reactions and incidents related to blood or blood products
- There is no cost
- Has a resource library
- Has many training resources with webcast training lectures on device associated module, catheter associated UTI, MRDO surveillance etc.

Identifying Infections 242

- ASC must develop and implement interventions to address issues identified through the detection activities
- ASC must then monitor the effectiveness of the intervention to make sure it worked
  - Can collect data and analysis to show it worked
- Must not only have policies but must make sure they are being followed
  - Do staff use hand hygiene before providing care to patients
  - Is the equipment sterilized and disinfected properly

CDC Guidelines for Environmental IC

- Chapters on;
- Air and water including dialysis water control
- Laundry and bedding
- Cleaning spills of blood
- Carpeting and pest control
- Regulated medical waste etc.
- See also Guidelines for the Disinfection and Sterilization in Healthcare Facilities
  - Cleaning equipment and scopes

www.cdc.gov/hicpac/Disinfection_Sterilization/toc.html

Guideline for Disinfection and Sterilization in Healthcare Facilities, 2008

William A. Rutala, Ph.D., M.P.H.[1,2], David J. Weber, M.D., M.P.H.[1,2], and the Healthcare Infection Control Practices Advisory Committee (HICPAC)[3]

[1]Hospital Epidemiology
University of North Carolina Health Care System
Chapel Hill, NC 27514

[2]Division of Infectious Diseases
University of North Carolina School of Medicine
Chapel Hill, NC 27599-7030

APIC Guidelines

Scientific guidelines

A A A 2

Centers for Disease Control and Prevention

The following guidelines have been released by the Centers for Disease Control and Prevention (CDC). They are listed in reverse chronological order.

Guidelines for the prevention of intravascular catheter-related infections were developed for healthcare personnel who insert intravascular catheters and for persons responsible for surveillance and control of infections in hospital, outpatient, and home healthcare settings.

Guideline for the prevention and control of norovirus gastroenteritis outbreaks in healthcare settings addresses prevention and control of norovirus gastroenteritis outbreaks in healthcare settings, and includes specific recommendations for implementation, performance measurement, and surveillance.

Guideline for prevention of catheter-associated urinary tract infections reviews the available evidence on CAUTI prevention for patients requiring chronic indwelling catheters and individuals who can be managed with alternative methods of urinary drainage.

Guidance for control of infections with carbapenem-resistant or carbapenemase-producing Enterobacteriaceae in acute care facilities provides new guidance for CRE infection prevention and control in an effort to limit the further emergence of these organisms.

Guidance on public reporting of healthcare-associated infections: recommendations is based on established principles for public health and healthcare-associated infection (HAI) reporting systems. It is intended to assist those tasked with designing and implementing public reporting systems for HAIs.

Guideline for disinfection and sterilization in healthcare facilities presents evidence-based recommendations on the preferred methods for cleaning, disinfection, and sterilization of patient-care medical devices and for cleaning and disinfecting the healthcare environment.

Guideline for isolation precautions in hospitals updates and expands the *1996 Guideline for isolation precautions in hospitals*.

2006 guide for management of multidrug resistant organisms in healthcare settings provides guidance for the implementation of strategies and practices to prevent the transmission of MRSA, VRE, and other MDROs.

Guidelines for preventing the transmission of Mycobacterium tuberculosis in healthcare facilities updates tuberculosis (TB) control recommendations reflecting shifts in the epidemiology of TB, advances in scientific understanding, and changes in healthcare practice that have occurred in the United States.

Guideline for preventing healthcare-associated pneumonia, 2003 updates, expands, and replaces the previously

http://www.cdc.gov/hicpac/pubReportGuide/publicReportingHAI.html

Program Evaluation 242

- IC program must be part of the PI program
- ASC are performing more invasive procedures increasing the risk for developing and transmitting infections
- Patients have varying degrees of wellness which makes this IC challenging in the ASC
  - May include elderly or patients with immuno-compromised state
- ASC should design their infection control program with this in mind

Program Evaluation 242

- Need to take precautions for patients who present with risk of transmission by airborne or droplet route
  - Would include prompt separation
  - Implement respiratory hygiene/ cough etiquette protocols
  - Cover your cough campaign
- Appropriate isolation precautions based on the routes of transmission of the suspected infection
  - CDC has isolation guidelines for hospitals that can be helpful for the ASC infection preventionist
  - www.cdc.gov/hicpac/2007IP/2007isolationPrecautions.html

CDC Home

Centers for Disease Control and Prevention

Your Online Source for Credible Health Information

SEARCH

A-Z Index A B C D E F G H I J K L M N O P Q R S T U V W X Y Z #

Healthcare Infection Control Practices Advisory Committee (HICPAC)

www.cdc.gov/hicpac/pdf/isolation/isolation2007.pdf

HICPAC

- About HICPAC
- Member Roster
- Charter
- Event Calendar
- Methodology Guideline
- Projects in Progress
- Publications
  - 2008 Disinfection & Sterilization Guideline
  - **Guideline for Isolation Precautions 2007**
    - Executive Summary
    - Part I: Review of

HICPAC > Publications

2007 Guideline for Isolation Precautions: Preventing Transmission of Infectious Agents in Healthcare Settings

Download the complete PDF version Guideline for Isolation Precautions: Preventing Transmission of Infectious Agents in Healthcare Settings 2007
PDF (3.80 MB / 225 pages)

Jane D. Siegel, MD; Emily Rhinehart, RN MPH CIC; Marguerite Jackson, PhD ; Linda Chiarello, RN MS; the Healthcare Infection Control Practices Advisory Committee

Acknowledgement: The authors and HICPAC gratefully acknowledge Dr. Larry Strausbaugh for his many contributions and valued guidance in the preparation of this guideline.

Text size: S M L XL

Email page

Print page

Bookmark and share

Download page:

PDF [157 KB]

Contact HICPAC:

Centers for Disease Control and Prevention
Healthcare Infection Control Practices Advisory Committee (HICPAC)
MS A07
1600 Clifton Rd
Atlanta, GA 30333

Cover Your Cough Posters



www.cdc.gov/flu/protect/covercough.htm

Survey Procedures 242

- Surveyor is told to use the IC tool to assess compliance with this standard
- Determine if there is an ongoing program to prevent, control, and investigate diseases among all staff, contract workers, and volunteers
- Make sure P&P are implemented correctly and consistent with national standards of care
- Make sure ASC mitigates risk that attribute to HAIs like hand hygiene
- Make sure ASC is tracking and monitoring all activities

Infection Control Program 243

- **Standard:** The program must be under the direction of a designated and qualified professional with training in infection control
- Called an infection preventionist (IP) now by APIC
- Must designate in writing who this is

Infection Control Program 243

- ASC must make sure this person has training in the principles and methods of infection control
- Certification in IC is desirable but not required
- Must maintain qualifications through education and training (will look at personnel file of IP)
  - Recommend national and local meetings such as APIC and SHEA
- Does not specify how many FTEs this is but need to have adequate resources to get the job done
  - APIC and SHEA publish studies and recommendations on resource allocation

Infection Control Program 244

- **Standard:** IC program is an integral part of the PI program
  - IC program must be integrated into the PI program
  - Data and program activities are a part of PI
- The program measures indicators and activities related to IC on an ongoing basis
- Surveyor to make sure IC activities result in specific actions to improve IC
- Two of the common things measured are geared at reducing infections

What do you Measure for PI?

- Patient Burns and Patient Fall in the ASC
- Prophylactic IV Antibiotic on Time
- Hospital Transfer/Admission
- Wrong Site, Side, Patient, Procedure or Implant
- Appropriate Surgical Site Hair Removal
- Normothermia Outcome
- Unplanned Anterior Vitrectomy
- Safe surgery checklist and volume data on selected surgical procedures, Flu vaccine rates among personnel, endoscopy surveillance and follow up and surveillance of patients with adenomatous polyps, cataract improvement in 90 days

ASC Quality Collaboration

Infection Control Program 245 12-12-2013

- **Standard**: The IC program must provide a plan of action for preventing, identifying, and managing infections and communicable disease
- IC program must implement the corrective and preventive measures that result in improvement
- ASC must have a comprehensive plan
  - Consider electronic surveillance
  - Could include hand hygiene, infectious waste management, bloodborne pathogens, bioterrorism readiness, patient outcomes, duties of the IP, surveillance strategies, environmental monitoring, antibiotic utilization studies, outbreak management, communicable disease reporting, education, P&P required and followed, etc.

CMS ASC Quality Reporting



www.cms.gov/Medicare/Quality-Initiatives-Patient-Assessment-Instruments/ASC-Quality-Reporting/index.html

CMS ASC Measures Flu in 2016

Measures for CY 2015 Payment Determination

- ASC-1 Patient Burn
- ASC-2 Patient Fall
- ASC-3 Wrong Site, Wrong Side, Wrong Patient, Wrong Procedure, Wrong Implant
- ASC-4 Hospital Transfer/Admission
- ASC-5 Prophylactic Intravenous (IV) Antibiotic Timing
- ASC-6 Safe Surgery Checklist Use
- ASC-7 ASC Facility Volume Data on Selected ASC Surgical Procedures

Measures for CY 2016 Payment Determination

- ASC-1 Patient Burn
- ASC-2 Patient Fall
- ASC-3 Wrong Site, Wrong Side, Wrong Patient, Wrong Procedure, Wrong Implant
- ASC-4 Hospital Transfer/Admission
- ASC-5 Prophylactic Intravenous (IV) Antibiotic Timing
- ASC-6 Safe Surgery Checklist Use
- ASC-7 ASC Facility Volume Data on Selected ASC Surgical Procedures
- ASC-8 Influenza Vaccination Coverage among Healthcare Personnel

Infection Control Plan 245

- IC plan must result in improvements in outcomes
- Plan must include the following:
  - Maintenance of a sanitary environment
  - Development and implementation of infection control measures related to ASC personnel
  - Mitigation of risks associated with patient infections present upon admission
  - Mitigation of risks contributing to healthcare-associated infections

Infection Control Plan 245

- Active surveillance
- Monitoring compliance with all P&P, protocols, and other IC program requirements
- Plan evaluation and revision of the plan when indicated
- Coordination as required by law with federal, state, and local emergency preparedness and health authorities to address communicable and infectious disease threats and outbreaks
- Compliance with reportable disease requirements of the local health authority

Infection Control Plan 245

- ASCs are required to have a process to follow up each patient after discharge
- This is done to identify and track infections associated with the ASC visit
- If ASCs learns of an infection in the post-operative period from the physician or the patient
  - ASC may consider inquiring if there a lab confirmation of an infectious disease
  - Are there indications the infection was associated with the stay
  - Can delegate part of follow-up to physicians on staff if information is reported back to ASC and documented in the medical record

Infection Control Plan 245

- Surveyor is suppose to make sure plan has all the required elements of IC
- Suppose to ask for documentation of actual examples of how issues were identified and corrected
- Make sure those actions resulted in improvement
- IP to go over IC plan with surveyor and explain how it addresses the fundamental elements of an IC program

Infection Control Plan 245

- Surveyor procedure
  - Surveyor is suppose to ask leadership how it tracks infections among both patients and staff
  - Surveyor is to ask for documentation that infections are tracked for all patients
  - Surveyor is suppose to ask the leadership what diseases are reported to the staff to verify that the ASC is aware of the state department of health reporting requirements
  - Surveyor will ask if the ASC has ever reported a reportable disease to the state
    - If yes, will review documentation of the case

ASC Infection Control Plan

OUTCOME AND PROCESS MEASURES

Process	Goal	Opportunities to Reduce Risk	Assessment	Measure of Success	Responsible Party
Hand Hygiene	Staff will perform hand hygiene at every opportunity following CDC guidelines with an observation rate > 80%.	Staff will perform hand hygiene with Alcohol Hand Rub or Soap & Water before & after patient contact. After removing gloves and after contact with blood, body fluids or contaminated surfaces. Gloves changed between patients/activities. Gloves are worn when coming in contact with a potentially infectious patients or blood/body fluids or contaminated surfaces.	Observation	All opportunities for hand hygiene were taken. Observation rate is > 80%.	All staff and visitors

ASC Infection Control Plan

SSI	The rate of SSI's will be < 1% with a target of 0%.	Hand Hygiene/Scrub Patient preparation. Surgical attire procedures are followed. Sterile Field requirements are met. All per AORN guidelines. Environmental cleaning is done per CDC requirements.	Observation Reported breaches of surgical asepsis.	The ASC rate of SSI's is < 1%. All opportunities to reduce the risk of SSI's are being practiced.	All Staff

ASC Infection Control Plan

Outcome and Process Measures

Process	Goal	Opportunities to Reduce Risk	Assessment	Measure of Success	Responsible Party
Endoscope Reprocessing	Prevent HAIs' due to improper processing of endoscopes which can result in cross contamination or outbreaks. Rate will be < 1% with a target rate of 0%.	Limiting operating room traffic. Continuing education. Endoscopes will be processed per manufacturer's directions. Staff will wear appropriate PPE during reprocessing. Continuing education & competency assessment will be done at least annually.	Reprocessing logs are monitored. Observation of cleaning.	No cross-contamination or outbreaks are reported.	Staff
Injection Practices	Reduction/elimination of exposure to blood borne pathogens. Rate will be < 0.5% with a target rate of 0%.	Follow APIC's/CDC's guidelines for safe injection practices which include: Never use medication in a syringe for more than one patient even if the needle is changed between patients. Utilize sharps safety devices whenever possible.	Observe staffs injection practices. Vials are labeled with expiration date. Safety devices are activated when available.	Employee health nurse reports a rate of 0% exposure to BBP through unsafe injection practices.	All Staff

ASC Infection Control Plan

Outcome and Process Measures

Process	Goal	Opportunities to Reduce Risk	Assessment	Measure of Success	Responsible Party
Sterilization	Medical and surgical instruments do not transmit infectious pathogens to patients or staff. The ASC is sterilizing equipment using standard methods per guidelines of APIC, AAMI, AORN & CDC.	Sterilized equipment is cleaned and visually inspected prior to sterilization. All parameters of steam sterilization are met i.e. time, temperature level, pressure and moisture. Chemical indicators are placed in each load and inspected for integrity and reprocessed if there is any doubt of improper sterilization. Biological indicators are done	Packages pass visual inspection and chemical parameters. Flash sterilization is <1% of loads.	No cases of transmission of pathogens to patient or staff.	All staff

So What Are the CDC 10 Recommendations?

- The CDC has a page on Injection Safety that contains the excerpts from the Guideline for Isolation Precautions: Preventing Transmission of Infectious Agents in Healthcare Settings
- Summarizes their 10 recommendations
- CMS expects hospitals to follow the CDC guidelines
- Available at http://www.cdc.gov/ncidod/dhqp/injectionSafetyPractices.html

CDC Safe Injection Recommendations

- Use aseptic technique to avoid contamination of sterile injection equipment.
- Do not administer medications from a syringe to multiple patients, even if the needle or cannula on the syringe is changed.
  - Needles, cannula and syringes are sterile, single-use items; they should not be reused for another patient nor to access a medication or solution that might be used for a subsequent patient.

CDC Safe Injection Recommendations

- Use fluid infusion and administration sets (i.e., intravenous bags, tubing and connectors) for one patient only and dispose appropriately after use
- Consider a syringe, needle, or cannula contaminated once it has been used to enter or connect to a patient's intravenous infusion bag or administration set

CDC Safe Injection Recommendations

- Use single-dose vials for parenteral medications whenever possible
- Do not administer medications from single-dose vials or ampules to multiple patients or combine leftover contents for later use
- If multidose vials must be used, both the needle or cannula and syringe used to access the multidose vial must be sterile

CDC Safe Injection Recommendations

- Do not keep multidose vials in the immediate patient treatment area and store in accordance with the manufacturer's recommendations;
  - Discard if sterility is compromised or questionable
- Do not use bags or bottles of intravenous solution as a common source of supply for multiple patients

CDC Safe Injection Recommendations

- Worker safety; Adhere to federal (OSHA) and state requirements for protection of healthcare personnel from exposure to blood borne pathogens
- Wear a mask when placing a catheter or injecting material into the spinal canal or subdural space
  - Example, during myelograms, lumbar puncture and spinal or epidural anesthesia.

The End! Questions???

- Sue Dill Calloway RN, Esq. CPHRM, CCMSCP, CCMSCP
- AD, BA, BSN, MSN, JD
- President of Patient Safety and Education Consulting
- 5447 Fawnbrook Lane
- Dublin, Ohio 43017
- 614 791-1468
- sdill1@columbus.rr.com
- (Call with questions, No emails)

Resources

- There are many places in the rest of the ASC manual that reference infection control
- Information is provided on those sections
- References to the national standards of care and practice are also included such as APIC, SHEA, etc.

Infection Control Other Sections

- Board section (states that surveyor is suppose to ask who performs infection control
  - APIC now calls them a infection preventionist
- Surgical Services section states that surgical procedures must be performed in a safe manner
- Safe manner is defined and states must follow acceptable standards of practice including the APIC standards
  - APIC stands for Association of Professionals in Infection Control and Epidemiology

Infection Control

- Surgical services also states that infection control includes the following
  - The conformance to aseptic and sterile technique by all individuals in the surgical area
  - That there is appropriate cleaning between surgical cases and appropriate terminal cleaning applied
    - AORN has section on how to clean the OR and between cases
  - That operating room attire is suitable for the kind of surgical case performed, that persons working in the operating suite must wear only clean surgical costumes, that surgical costumes are designed for maximum skin and hair coverage

Infection Control

- AORN has section on what is suitable attire in the OR in their Perioperative Standards and Recommended Practices
- Surgical services also states that infection control includes the following
  - That equipment is available for rapid “emergency” high-level disinfection/ sterilization of operating room materials
  - This is an important issue with both TJC (the Joint Commission) and CMS
  - CMS issues a memo on this for ASCs

Infection Control

- That sterilized materials are packaged, handled, labeled, and stored in a manner that ensures sterility e.g., in a moisture- and dust-controlled environment, and policies and procedures for expiration dates have been developed and are followed in accordance with accepted standards of practice.
- That temperature and humidity are monitored and maintained within accepted standards of practice
  - AORN in the Perioperative Standards and Recommended Practices state what the temperature and humidity should be in the OR and PACU

Infection Control

- ASC that identifies problems with IC thru the PI program should take action to reduce the potential for transmission (p
- Tag 81 discusses that the ASC must measure, analyze, and track quality indicators, adverse events and infection control performance
- ASC should have outcome indicators and focus on complication rates and HAI rates
- ASC should look at surgical site infection rate and make sure antibiotic prophylaxis meets the SOC such as the CDC guidelines

Infection Control

- The location of the OR within the ASC and the access to it must conform to accepted standards of practice, particularly for infection control, with respect to the movement of people, equipment and supplies in and out of the OR
- Discuss rapid emergency sterilization of OR equipment and says should not be routinely used since intended for emergency use
- Can allow individuals to accompany patients in the PACU as long as consistent with SOP and infection control guidelines

Infection Control

- Tag 103 on the physical environment has many sections on infection control including
  - The ASC must establish a program for identifying and preventing infections, maintaining a sanitary environment, and reporting the results to appropriate authorities
  - Must have program to follow up every patient after discharge to identify and track if they had an infection associated with the stay in the ASC
  - If find out about an infection from the surgeon then needs to investigate to see if from the ASC
- Surveyor to ask LD how it tracks infections among staff and patients

Infection Control

- Tag 122 discusses that ASC needs to track infection rate when evaluating whether to recredential the physicians
- Tag 181 on administration of drugs discusses using standard infection control practices when using injectible medications
- Radiology standards require enforcing infection control standards
- Tag 232 on patient rights says staff should follow all current infection control standards

APIC Resources for Ambulatory Care

Infection Control in Ambulatory Care

- Guidelines have been developed by the CDC's Healthcare Infection Control Practices Advisory Committee (HICPAC) www.cdc.gov/ncidod/dhqp/hicpac_pubs.html
  - Infection control plan for ambulatory care
  - Guidelines for Disinfection and Sterilization in Healthcare Facilities 2008
  - Guidelines for Isolation Precautions 2007
  - Management of Multidrug-Resistant Organisms 2006
  - Influenza Vaccination of Healthcare Personnel 2006 and CDC has flu website at www.flu.gov with information on universal flu vaccine(all strains of the flu for decades)

www.infectioncontroltoday.com/

Bioterrorism

- ASC should be knowledgeable about emergency preparedness, including bioterrorism
- Terrorists could use bioterrorism
- There is a long list of bioterrorism agents
  - Anthrax, arena viruses, botulism, brucellosis, cholera, Ebola virus, hemorrhagic fever, E. coli, Lassa fever, plague, ricin toxin, salmonella, and cryptosporidium
- For a comprehensive list go to website[1]

[1]http://www.emergency.cdc.gov/agent/agentlist.asp

http://emergency.cdc.gov/bioterrorism/

Emergency Preparedness and Response

Emergency Preparedness & Response

- Specific Hazards
- **Bioterrorism**
  - Chemical
  - Gulf Oil Spill 2010
  - Radiation
  - Mass Casualties
  - Natural Disasters & Severe Weather
  - Recent Outbreaks & Incidents
- Preparedness for All Hazards
- What CDC Is Doing
- What You Can Do
- What's New

EPR > Specific Hazards



Bioterrorism

More images

Specific Bioterrorism Agents

- A–Z
- List by Category

Info for the General Public

- Overview
- Agent-Specific Fact Sheets

Info for Professionals

- Case Definitions
- Training
- First Responders
- Lab Info
- Surveillance
- Preparation & Planning
- Communicating in the First Hours: Initial Communication With the Public During a Potential Bioterrorism Event

Text size: S M L XL

Email page
Print page
Bookmark and share
Subscribe to RSS

View page in

中文 (Chinese)
Français (French)
Deutsch (German)
Italiano (Italian)
한국어 (Korean)
Русский (Russian)
Español (Spanish)
Tiếng Việt (Vietnamese)

Get email updates

To receive email updates about this

APIC
Spreading knowledge.
Preventing infection.™
MyAPIC Store Join APIC APICList Careers Site Map Contact Us
login / logoff
+ text
SEARCH
About APIC
Member Services
Education & Certification
Research
Guidelines & Standards
Public Policy
Resources/ Information Services
Emergency Preparedness
EMERGENCY PREPAREDNESS
Incident Command Training
Pandemics
H1N1 Flu Information
Bioterror
Surge Capacity
Special Needs Populations
Natural Disasters
Position Papers
Print this Page
Email this Page
Home → Emergency Preparedness
Emergency Preparedness
APIC offers educational and reference materials for infection preventionists, healthcare and public health professionals, and disaster planners related to infection prevention and emergency preparedness enabling members to collaborate with local, state, and federal agencies in areas related to preventing infection transmission during disasters.
You may also visit the Emergency Preparedness Conference Education Library to purchase recordings of emergency preparedness sessions from the 2009 APIC Annual Conference.
Incident Command Training
Links to FEMA's Incident Command... more
Pandemics
Infection prevention information for the novel pathogenic organism... more
Bioterror
Critical information on bioterrorism agents... more
Surge Capacity
Addressing infection prevention challenges created by a patient surge... more


APIC
Spreading knowledge.
Preventing infection.™
MyAPIC Store Join APIC APICList Careers Site Map Contact Us
login / logoff
+ text
SEARCH
About APIC
Member Services
Education & Certification
Research
Guidelines & Standards
Public Policy
Resources/ Information Services
Emergency Preparedness
GUIDELINES & STANDARDS
Construction Issues
Definitions & Surveillance
Environmental Services
Guidelines & Standards
HAI Compendium
Healthcare Workers & Infection Prevention
Multi Drug Resistance Organisms
Targeting Zero
I Believe in Zero CLABSIs Campaign
APIC Elimination Guides
Topics
Surveillance Technology Resources
National Patient Safety Goals
Hand Hygiene
Position Statements
Home → Guidelines & Standards → Targeting Zero
Targeting Zero
Targeting Zero
Targeting Zero is the philosophy that every healthcare institution should be working toward a goal of zero healthcare-associated infections (HAIs). While HAI prevention is challenging and complex, APIC believes that all organizations should set the aspirational goal of elimination and strive for zero infections. Every HAI impacts the life of a patient and a family, and even one should be considered too many. Further, unsafe behaviors and practices that place patients and healthcare workers at risk for HAIs should never be tolerated.
Read APIC's Position Statement on Targeting Zero HAIs.
Enacting change is difficult and cannot occur overnight. To help you in Targeting Zero, APIC has created a comprehensive set of resources featuring webinars, conferences and practical tools.
Learn more about the newest Targeting Zero initiative: I Believe in Zero CLABSIs

Spreading knowledge.
Preventing infection.™

Contact: Liz Garman, 202/454-2604
egarman@apic.org

APIC INTRODUCES "GUIDE TO ELIMINATION OF INFECTIONS IN HEMODIALYSIS"

Washington, June16, 2010 – In an effort to establish universally accepted best practices for protecting patients undergoing hemodialysis, the Association for Professionals in Infection Control and Epidemiology (APIC) has published a *Guide to the Elimination of Infections in Hemodialysis*. Following cardiovascular disease, infection is the second largest cause of death for hemodialysis patients.

The *Guide* was developed by healthcare professionals responsible for infection prevention in acute care, ambulatory and home hemodialysis settings. With input from infection preventionists, dialysis technicians and dialysis nurses, the *Guide* presents effective and actionable steps to reduce the risk of infection and protect patients.

Hemodialysis technologies have greatly improved the health and quality of life for millions of people across the globe suffering from kidney disease and kidney failure. However, patients receiving these treatments are particularly vulnerable to healthcare-associated infections (HAIs) due to their weakened immune systems and the invasive nature of hemodialysis devices. For the majority of hemodialysis patients, treatments are administered three times weekly, further increasing the risk of being infected by pathogens present in healthcare settings. Hospitalizations due to infections in hemodialysis patients rose 34% between 1993 and 2006.

"The morbidity, mortality and cost of these infections are significant and increasing, yet preventable," said APIC CEO Kathy Warye. "APIC's new *Guide to the Elimination of Infections in Hemodialysis* provides evidence-based strategies to help protect this particularly vulnerable patient population from an increasingly prevalent problem. The new *Guide* will also prepare our members to meet future priorities from the federal government, which will focus on infection control in dialysis and ambulatory surgical

CDC has Dialysis Resources also



Home | About CDC | Press Room | A-Z Index | Contact Us

Department of Health and Human Services
Centers for Disease Control and Prevention

CDC en Español

Search: GO

Infection Control Home > Protecting Patients > Device & Procedure-associated Conditions >

Dialysis Patient Safety

Prevention & Control | Outbreaks | Data & Statistics | Emergency Preparedness

Prevention & Control of Dialysis-Associated Infections

These resources may be of use to healthcare professionals

Guidelines / Recommendations

- Infection Control Requirements for Dialysis Facilities and Clarification Regarding Guidance on Parenteral Medication Vials
  The rule establishes new conditions dialysis facilities must meet to be certified under the Medicare program and is intended to update CMS standards for delivery of quality care to dialysis patients.
  August 15, 2008 MMWR / 57(32);875-876
- Guideline for Isolation Precautions: Preventing Transmission of Infectious Agents in Healthcare Settings 2007
- Guideline for Environmental Infection Control in Health-Care Facilities, 2003
- Guidelines for the Prevention of Intravascular Catheter-Related Infections HICPAC 2002
- Guideline for Hand Hygiene in Healthcare Settings HICPAC 2002
- Recommendations for Preventing Transmission of Infections Among Chronic Hemodialysis Patients
  MMWR April 27, 2001/50(RR05);1-43

Infection Control Topics

- Infection Control Home
- Healthcare-Associated Infections
- **Protecting Patients**
- Protecting Healthcare Workers
- Infection Control Guidelines
- Infection Control A-Z
- About DHQP

Related Resources

Centers for Medicare & Medicaid Services (CMS)

- CMS ESRD General Information
- ESRD Network Organizations
- ESRD Core Indicators
- Fistula First
- Medicare Dialysis Facility Compare

National Institutes of Health (NIH)

- National Kidney and Urologic Diseases Clearing House

www.apic.org/Content/NavigationMenu/PracticeGuidance/APICEliminationGuides/Hemodialysis_Resourc.htm



PreventInfection.org

Prevention Central | eNewsletters | Resources | Partners | Events

Did you know your own skin could be the source of deadly infections that could kill you? Currently, 271 people a day die on average from healthcare-associated infections (HAIs). Many people, possibly including you,

1.7 million
people are impacted by these healthcare-associated infections

aren't even aware of these deadly infections let alone how to prevent them. It all starts at their source — your skin. To prevent these infections, it is important to understand why "skin is the source." Click to learn more.

h1n1 flu

H1N1 flu

Learn the basic facts and what you can do to protect yourself and your

how aware are you?

Be a Good Visitor

Be a good visitor. Read APIC's tips on being a good visitor to a healthcare

outpatient care

Infection Prevention in Outpatient Care

As outpatient surgery has become a

www.who.int/en/

World Health Organization

Search

All WHO This site only

Home

- About WHO
- Countries
- Health topics
- Publications
- Data and statistics
- Programmes and projects

HIGHLIGHTS

WHO/C. Gaggero

Essential medicines fact sheet

Haiti situation update

Access to medicines: the role of procurement policies

16 July 2010 -- WHO, WIPO and WTO are jointly exploring ways to improve access to medicines through better procurement policies. In her address, Dr Margaret Chan explains why the price of medicines is such a critical and emotive issue for public health.

Read the Director-General's speech on improving the access to medicines

Insuring a billion people in India

KEY WHO INFORMATION

Director-General
Director-General and senior management

Governance of WHO
WHO Constitution, Executive Board and World Health Assembly

Media centre
News, events, fact sheets, multimedia and contacts

International travel and health
Publication on travel risks, precautions and vaccination requirements

World Health Report
Annual report on global public health and key statistics

World Health Organization
عربي | 中文 | English | Français
All WHO
Home
About WHO
Countries
Health topics
Publications
Data and statistics
Programmes and projects
WHO > Health topics
Infection control
This page provides links to descriptions of activities, reports, news and events, as well as contacts and cooperating partners in the various WHO programmes and offices working on this topic. Also shown are links to related web sites and topics.
RELATED SITES
- Global Patient Safety Challenge
- Patient safety
- Infection prevention and control in health care
- Injection safety
- Infection control (Western Pacific Region)

http://www.his.org.uk/



www.theific.org/



Source: www.idsociety.org

Policy and Advocacy

- Policy Focus Areas
  - Access to Quality Health Care
  - Federal Funding
  - Medical Education and Research
  - Physician Practice and Payment
  - Product Research and Development
  - Public Health and Bioemergencies
  - Special Areas of Focus
    - Adult Immunization
    - Avian/Pandemic Flu
    - **Bad Bugs, No Drugs**
    - Home Infusion Therapy
    - Positions on Health Reform
    - STAAR Act
- Advocacy Center
- Clinical Practice
- Government ID Resources
- Global ID Center

PRINT THIS PAGE

EMAIL A FRIEND

Home → Policy and Advocacy → Policy Focus Areas → Special Areas of Focus → Bad Bugs, No Drugs

Bad Bugs, No Drugs... 10 New Antibiotics by 2020

IDSA Testimony on Antibiotic Resistance, delivered before the House Energy and Commerce Health Subcommittee (pdf) June 9, 2010
Brad Spellberg, MD, FIDSA, presented IDSA's statement at a hearing on "Promoting the Development of Antibiotics and Ensuring Judicious Use in Humans." FDA's and other statements and the hearing video are located here.

The 10 x '20 Initiative Inaugural Statement (pdf) April 2010
IDSA and other organizations address the dry antibiotic pipeline and call for 10 new antibiotics by 2020.

10 x '20 Endorsing Organizations (pdf) June 29, 2010

IDSA Report: Bad Bugs, No Drugs: As Antibiotic Discovery Stagnates, a Public Health Crisis Brews (pdf) July 2004
Learn about IDSA's concerns about the lack of new antibiotics to fight drug-

Practice Guidelines

- Standards, Practice Guidelines
- Guidelines for Mobile Devices
- HIVMA Guidelines
- Pocketcards
- Resources

PRINT THIS PAGE

EMAIL A FRIEND

Standards, Practice Guidelines, and Statements Developed and/or Endorsed by IDSA

About Practice Guidelines

Practice guidelines are systematically developed statements to assist practitioners and patients in making decisions about appropriate health care for specific clinical circumstances. [Institute of Medicine Committee to Advise the Public Health Service on Clinical Practice Guidelines, 1990] Attributes of good guidelines include validity, reliability, reproducibility, clinical applicability, clinical flexibility, clarity, multidisciplinary process, review of evidence, and documentation. [Institute of Medicine Committee to Advise the Public Health Service on Clinical Practice Guidelines, 1990]

It is important to realize that guidelines cannot always account for individual variation among patients. They are not intended to supplant physician judgment with respect to particular patients or special clinical situations. IDSA considers adherence to the guidelines listed below to be voluntary, with the ultimate determination regarding their application to be made by the physician in the light of each patient's individual circumstances.

IDSA Policy on its Clinical Practice Guidelines in Development

---

Standards, Practice Guidelines Summary Table

Clinical Practice Guidelines by Category

Antimicrobial Agent Use

PA Patient Safety Authority

Source: http://www.patientsafetyauthority.org/Pages/Default.aspx

Catheter-Associated Urinary Tract Infections (UTI)

Prevention & Control of Catheter-Associated Urinary Tract Infections (UTI)

This guideline updates and expands the original Centers for Disease Control and Prevention (CDC) Guideline for Prevention of Catheter-associated Urinary Tract Infections (CAUTI) published in 1981. Several developments necessitated revision of the 1981 guideline, including new research and technological advancements for preventing CAUTI, increasing need to address patients in non-acute care settings and patients requiring long-term urinary catheterization, and greater emphasis on prevention initiatives as well as better defined goals and metrics for outcomes and process measures. In addition to updating the previous guideline, this revised guideline reviews the available evidence on CAUTI prevention for patients requiring chronic indwelling catheters and individuals who can be managed with alternative methods of urinary drainage (e.g., intermittent catheterization).

2009 Guidelines

These resources may be of use to healthcare professionals

- Guideline for Prevention of Catheter-associated Urinary Tract Infections, 2009 (407 KB / 67 pages)
- Guideline for Prevention of CAUTI 2009 Appendices (4.41 MB / 268 pages)

Infection Control Topics

- Infection Control Home
- Healthcare-Associated Infections
- **Protecting Patients**
- Protecting Healthcare Workers
- Infection Control Guidelines
- Infection Control A-Z
- About DHQP

www.cdc.gov/hicpac/CAUTI_fastFacts.html



www.cdc.gov/hicpac/Disinfection_Sterilization/acknowledg.html



Healthcare Infection Control Practices Advisory Committee (HICPAC)

HICPAC
- About HICPAC
- Member Roster
- Charter
- Event Calendar
- Methodology Guideline
- Projects in Progress
- Publications
- **2008 Disinfection & Sterilization Guideline**
  - Table of Contents
  - Executive Summary
  - Approach to Disinfection and Sterilization
  - Factors Affecting

HICPAC > Publications

Guideline for Disinfection and Sterilization in Healthcare Facilities, 2008

The complete Guideline for Disinfection and Sterilization in Healthcare Facilities, 2008 is available for download in PDF format (948 KB / 158 pages).

William A. Rutala, Ph.D., M.P.H.[1,2], David J. Weber, M.D., M.P.H.[1,2], and the Healthcare Infection Control Practices Advisory Committee (HICPAC)[3]

[1]Hospital Epidemiology
University of North Carolina Health Care System
Chapel Hill, NC 27514

[2]Division of Infectious Diseases
University of North Carolina School of Medicine
Chapel Hill, NC 27599-7030

[3]HICPAC Members

Text size: S M L XL
Email page
Print page
Bookmark and share

Download page:
PDF [107 KB]

Contact HICPAC:

Centers for Disease Control and Prevention
Healthcare Infection Control Practices Advisory Committee (HICPAC)
MS A07
1600 Clifton Rd
Atlanta, GA 30333

800-CDC-INFO

http://www.cdc.gov/hicpac/pubs.html

- Fast Facts
- Podcast: Dr. Sanjay Saint discusses CAUTIs
- Management of MDRO's in Healthcare Settings
- Guidance on Public Reporting of HAIs 2003
- Proceedings & Presentations
- Contact HICPAC

2007 Guideline for Isolation Precautions: Preventing Transmission of Infectious Agents in Healthcare Settings This document is intended for use by infection control staff, healthcare epidemiologists, healthcare administrators, nurses, other healthcare providers, and persons responsible for developing, implementing, and evaluating infection control programs for healthcare settings across the continuum of care.
Complete PDF version available for download Guideline for Isolation Precautions: Preventing Transmission of Infectious Agents in Healthcare Settings 2007 PDF (3.80 MB / 225 pages)



Guidelines for Environmental Infection Control in Health-Care Facilities June 6, 2003 / 52(RR10);1-42 The *Guidelines for Environmental Infection Control in Health-Care Facilities* is a compilation of recommendations for the prevention and control of infectious diseases that are associated with healthcare environments.
Available for download [PDF - 1.4 MB]

Guidelines for Hand Hygiene in Healthcare Settings Published 2002 October 25, 2002 / Vol. 51 / No. RR-16 (PDF 496 KB / 56 pages)
Recent developments in the field have stimulated a review of the scientific data regarding hand hygiene and the development of new guidelines designed to improve hand-hygiene practices in

800-CDC-INFO
(800-232-4636)
TTY: (888) 232-6348
24 Hours/Every Day
hicpac@cdc.gov

This presentation is intended solely to provide general information and does not constitute legal advice. Attendance at the presentation or later review of these printed materials does not create an attorney-client relationship with the presenter(s). You should not take any action based upon any information in this presentation without first consulting legal counsel familiar with your particular circumstances.

Thanks for Attending!

- Sue Dill Calloway RN, Esq. CPHRM, CCMSCP
- AD, BA, BSN, MSN, JD
- President of Patient Safety and Education Consulting
- 5447 Fawnbrook Lane
- Dublin, Ohio 43017
- 614 791-1468
- sdill1@columbus.rr.com
- (Call with questions, No emails)