

The Latest and Greatest CMS Nursing CoPs



The information provided in AHC Media Webinars does not, and is not intended to constitute medical or legal advice. Opinions, references and links provided by our speakers are provided for your convenience and do not represent our endorsement of such opinions, products or services.

Speaker



- Sue Dill Calloway RN, Esq.
CPHRM, CCMSCP
- AD, BA, BSN, MSN, JD
- President of Patient Safety and
Education Consulting
- 5447 Fawnbrook Lane
- Dublin, Ohio 43017
- 614 791-1468 (Call with Questions, No emails)
- sdill1@columbus.rr.com
- Email questions to CMS at hospitalscg@cms.hhs.gov


Objectives

- Discuss frequently cited areas within the Nursing CoPs.
- Recall why all medications must be administered within three different time frames.
- Review why all verbal orders must be signed off with a corresponding date and time.
- Explain new and revised standards, regulations, and laws put forth by CMS, TJC and the federal government.
- Evaluate compliance requirements and penalties.

Introduction to the CMS Hospital Conditions of Participation (CoPs)



You Don't Want One of These



The Conditions of Participation (CoPs)

- Regulations first published in 1986
 - Manual updated more frequently now
 - Many changes since 1986
- First regulations are published in the **Federal Register** then CMS publishes the **Interpretive Guidelines** and some have **survey procedures** ²
 - Hospitals should check this website once a month for changes

¹www.gpoaccess.gov/fr/index.html ²www.cms.hhs.gov/SurveyCertificationGenInfo/PMSR/list.asp

Location of CMS Hospital CoP Manuals

Medicare State Operations Manual Appendix

Email questions to CMS at hospitalscg@cms.hhs.gov

- Each Appendix is a separate file that can be accessed directly from the SOM Appendices Table of Contents, as applicable.
- The appendices are in PDF format, which is the format generally used in the IOM to display files. Click on the red button in the "Download" column to see any available file in PDF.
- To return to this page after opening a PDF file on your desktop, use the browser "back" button. This is because closing the file usually will also close most browsers

CMS Hospital CoP Manuals **new** address

www.cms.hhs.gov/manuals/downloads/som107_Appendixtoc.pdf

App. No.	Description	PDF File
A	Hospitals	 2,185 KB
AA	Psychiatric Hospitals	 606 KB

CoP Manual Also Called SOM

State Operations Manual Appendix A - Survey Protocol, Regulations and Interpretive Guidelines for Hospitals

Table of Contents
(Rev. 151, 11-20-15)

www.cms.hhs.gov/manuals/downloads/som107Appendixtoc.p

Transmittals for Appendix A

Survey Protocol

Introduction

Task 1 - Off-Site Survey Preparation

Task 2 - Entrance Activities

Task 3 - Information Gathering/Investigation

Task 4 - Preliminary Decision Making and Analysis of Findings

Task 5 - Exit Conference

Task 6 - Post-Survey Activities

Psychiatric Hospital Survey Module


Psychiatric Unit Survey Module

Rehabilitation Hospital Survey Module

Inpatient Rehabilitation Unit Survey Module

Hospital Swing-Bed Survey Module

Email questions
hospitalscg@cms.hhs.gov



CMS Survey and Certification Website

The screenshot shows the CMS.gov website interface. At the top, there is a navigation bar with links for Home, About CMS, Careers, Newsroom, FAQ, Archive, and social media icons for RSS, Facebook, and Twitter. Below this is the CMS.gov logo and the text 'Centers for Medicare & Medicaid Services'. A search bar is located on the right side of the header. A horizontal menu below the header contains eight categories: Medicare, Medicaid/CHIP, Medicare-Medicaid Coordination, Insurance Oversight, Innovation Center, Regulations, Guidance & Standards, Research, Statistics, Data & Systems, and Outreach & Education. The breadcrumb trail reads: CMS Home > Medicare > Survey & Certification - General Information > Policy & Memos to States and Regions. On the left, a sidebar titled 'Survey & Certification - General Information' lists various topics, with 'Policy & Memos to States and Regions' highlighted. The main content area is titled 'Policy & Memos to States and Regions' and contains a description: 'CMS Survey and Certification memoranda, guidance, clarifications and instructions to State Survey Agencies and CMS Regional Offices.' Below the description, there are filter options: 'Select From The Following Options:' with radio buttons for 'Show all items' (selected) and 'Show only (select one or more options):'. The 'Show only' options include: 'Show only items whose [] is within the past []', 'Show only items whose Fiscal Year is []', and 'Show only items containing the following word: []'. A 'Show Items' button is at the bottom of the filter section. Below the button, it says 'There are 455 items in this list.' A URL is overlaid on the right side of the page: www.cms.gov/SurveyCertificationGenInfo/PMSR/list.asp#TopOfPage.

CMS Survey Memos

Policy & Memos to States and Regions

CMS Survey and Certification memoranda, guidance, clarifications and instructions to State Survey Agencies and CMS Regional Offices.

Title 	Memo # 	Posting Date 	Fiscal Year 
Certification Number (CCN) State Codes –State Operations Manual (SOM) Section 2779A Revisions	16-09-ALL	2016-03-11	2016
Advance Copy – Interpretive Guidelines for the Organ Transplant Conditions of Participation (CoPs) at 42 Code of Federal Regulations (CFR) §§ 482.68 through 482.104	16-10-Transplant	2016-03-11	2016
National Downtime of the Quality Improvement and Evaluation System (QIES)	16-12-ALL	2016-03-11	2016
Critical Access Hospital (CAH) Recertification Checklist for Evaluation of Compliance with the Location and Distance Requirements	16-08-CAH	2016-02-12	2016
FY 2015 Report to Congress (RTC): Review of Medicare’s Program Oversight of Accrediting Organizations (AOs) and the Clinical Laboratory Improvement Amendments of 1988 (CLIA) Validation Program	16-07-AO	2016-01-29	2016
Medicare Learning Network (MLN) Infection Control Courses	16-06-ALL	2016-01-22	2016

Example of a CMS Survey Memo

DEPARTMENT OF HEALTH & HUMAN SERVICES
Centers for Medicare & Medicaid Services
7500 Security Boulevard, Mail Stop C2-21-16
Baltimore, Maryland 21244-1850



Center for Clinical Standards and Quality/Survey & Certification Group

Ref: S&C: 14-15-Hospital

DATE: March 14, 2014

TO: State Survey Agency Directors

FROM: Director
Survey and Certification Group

SUBJECT: Requirements for Hospital Medication Administration, Particularly Intravenous (IV) Medications and Post-Operative Care of Patients Receiving IV Opioids

Memorandum Summary

- **Medication Administration:** We are updating our guidance for the hospital medication administration requirements to:
 - Make clear that the medication administration requirements under the nursing services condition of participation (CoP) are related to only some components of the overall hospital medication process, but that hospitals are expected, through this and the related requirements under the pharmaceutical services and quality assessment/performance improvement CoPs, to take a comprehensive approach to the medication process.
 - Update our guidance for IV medications and blood transfusions in general; and
 - Reflect the need for patient risk assessment and appropriate monitoring during and after medication administration, particularly for post-operative patients receiving IV opioid medications, in order to prevent adverse events.
- **Immediate Post-operative Care:** Clarification is also being made to the guidance for the surgical services CoP requirement for hospitals to have adequate provisions for immediate post-operative care, to emphasize the need for post-operative monitoring of patients

CAH Memo Makes Many Similar Regulations

DEPARTMENT OF HEALTH & HUMAN SERVICES
Centers for Medicare & Medicaid Services
7500 Security Boulevard, Mail Stop C2-21-16
Baltimore, Maryland 21244-1850



Center for Clinical Standards and Quality / Survey & Certification Group

Ref: S&C: 15-19-CAH

DATE: January 16, 2015
TO: State Survey Agency Directors
FROM: Director
Survey and Certification Group
SUBJECT: Revised State Operations Manual (SOM) Appendix W, Critical Access Hospitals (CAHs)

Memorandum Summary

The Centers for Medicare & Medicaid Services (CMS) CAH Conditions of Participation (CoPs) Changed in Two Final Rules:

- CMS-3267-F was published on May 12, 2014 and portions related to CAHs became effective July 11, 2014. Among other provisions, this final rule revised the CAH Conditions of Participation (CoP) requirements related to the responsibilities of doctors of medicine (MDs) and doctors of osteopathy (DOs).
- CMS-1599-F was published August 19, 2013 and became effective October 1, 2013. This final rule revised the CAH CoP requirements related to provision of inpatient acute care services.

SOM Appendix W Updated:

- We are updating the pertinent portions of the CAH interpretive guidelines, found in SOM Appendix W, to reflect these rule changes.

The Conditions of Participation (CoPs)

- The manual is known as the conditions of participation or the CoPs for short
- The CoP sections are called tag numbers
- They go from Tag 0001 to 1164 and nursing starts at tag 385 in Appendix A
 - All the sections contain a tag number so it is easy to go back and look up that section if you want to read more about it
 - CAH has separate manual – Appendix W

How to Keep Up with Changes

- First, periodically check to see you have the most current CoP manual¹
- Once a month go out and check the survey and certification website²
- Once a month check the CMS transmittal page³
- Have one person in your facility who has this responsibility

■ ¹ http://www.cms.hhs.gov/manuals/downloads/som107_Appendicestoc.pdf

■ ² <http://www.cms.gov/SurveyCertificationGenInfo/PMSR/list.asp#TopOfPage>

■ ³ <http://www.cms.gov/Transmittals>

Transmittals

CMS.gov

Centers for Medicare & Medicaid Services

Learn about [your healthcare options](#)

- Medicare
- Medicaid/CHIP
- Medicare-Medicaid Coordination
- Private Insurance
- Innovation Center
- Regulations & Guidance**
- Research, Statistics Data & Systems

Home > Regulations and Guidance > Transmittals > 2016 Transmittals

www.cms.gov/Regulations-and-Guidance/Guidance/Transmittals/2016-Transmittals.html

Transmittals

[2016 Transmittals](#)

[2015 Transmittals](#)

[2014 Transmittals](#)

[2013 Transmittals](#)

[2012 Transmittals](#)

[2011 Transmittals](#)

[2010 Transmittals](#)

[2009 Transmittals](#)

[2008 Transmittals](#)

[2007 Transmittals](#)

[2006 Transmittals](#)

[2005 Transmittals](#)

[2004 Transmittals](#)

2016 Transmittals

Show entries: 10

Filter On:

Transmittal #	Issue Date	Subject	Implementation Date	CR #	MM Article #
-------------------------------	----------------------------	-------------------------	-------------------------------------	----------------------	------------------------------

SE1606		Guidance on the Physician Quality Reporting System (PQRS) 2014 Reporting Year and 2016 Payment Adjustment for Rural Health Clinics (RHCs), Federally Qualified Health Centers (FQHCs), and Critical Access Hospitals (CAHs)			SE1606
------------------------	--	---	--	--	--------

CMS Proposed Changes to the CoPs on Nursing



CMS Proposes New Changes

- CMS proposes new changes to the hospital CoPs which address infection control
 - Called the Improvement in Patient Care Act
- Published in the Federal Register on June 16, 2016
- Would require every hospital to have an antimicrobial stewardship program
- Change title to Infection Prevention & Control and Antibiotic Stewardship
- IP must be qualified and appointed by board

Antibiotic Stewardship Program

The screenshot displays the Federal Register website interface. At the top, there is a navigation bar with links for Sections, Browse, Search, Policy, Learn, Blog, and My FR, along with a search box for documents. The Federal Register logo and the text "The Daily Journal of the United States Government" are visible. A blue banner highlights the "Proposed Rule" section. The main heading is "Medicare and Medicaid Programs; Hospital and Critical Access Hospital (CAH) Changes To Promote Innovation, Flexibility, and Improvement in Patient Care". Below this, it states "A Proposed Rule by the Centers for Medicare & Medicaid Services on 06/16/2016". There are social media icons for flagging, email, Twitter, and Facebook. A green button says "SUBMIT A FORMAL COMMENT". A grey box indicates "Comments on this document are being accepted at Regulations.gov". A link says "Read the 8 submitted public comments". Navigation arrows for "Previous Document" and "Next Document" are present. A red box contains a "LEGAL DISCLAIMER". A "Font Controls" section includes plus, minus, and text size icons. A utility section offers "PDF", "DEV", and "PRINT" options, along with a "PUBLIC INSPECTION" link. The "Publication Date" is listed as "Thursday, June 16, 2016", and the "Agencies" section is partially visible. A vertical "Site Feedback" button is on the right side.

Sections ▾ Browse ▾ Search ▾ Policy ▾ Learn ▾ Blog ▾ My FR ▾ Search Documents

ARCHIVES AND RECORDS ADMINISTRATION
1985

FEDERAL REGISTER
The Daily Journal of the United States Government

SCRIPTA MANET
Sign in Sign up

Site Feedback

Proposed Rule

Medicare and Medicaid Programs; Hospital and Critical Access Hospital (CAH) Changes To Promote Innovation, Flexibility, and Improvement in Patient Care

A Proposed Rule by the [Centers for Medicare & Medicaid Services](#) on 06/16/2016

Comments on this document are being accepted at [Regulations.gov](#)

SUBMIT A FORMAL COMMENT

Read the **8** submitted public comments

← Previous Document
Next Document →

LEGAL DISCLAIMER

Font Controls + - A A

PDF DEV PRINT
PUBLIC INSPECTION

Publication Date:
Thursday, June 16, 2016

Agencies:

ACTION Proposed Rule.

SUMMARY This proposed rule would update the requirements that hospitals and critical access hospitals (CAHs) must meet to participate in the Medicare and Medicaid programs. These proposals are intended to conform the requirements to current standards of practice and support improvements in quality of care, reduce barriers to care, and reduce some issues that may exacerbate workforce shortage concerns.

UNIFIED AGENDA [Hospital and Critical Access Hospital \(CAH\) Changes to](#)



FEDERAL REGISTER

Vol. 81

Thursday,

No. 116

June 16, 2016

www.gpo.gov/fdsys/pkg/FR-2016-06-16/pdf/2016-13925.pdf

Part IV

Department of Health and Human Services

Centers for Medicare & Medicaid Services

42 CFR Parts 482 and 485

Medicare and Medicaid Programs; Hospital and Critical Access Hospital (CAH) Changes To Promote Innovation, Flexibility, and Improvement in Patient Care; Proposed Rule

CMS Proposes New Changes to Nursing

- Would change LIP to LP so PA can order restraints and assess patient
- Would clarify the need for adequate staff on each unit and need supervision to make sure immediate availability of RN at the bedside when needed
- Must have enough staff to respond to patient needs
- Must have P&P as to which outpatient areas you need an RN and review every 3 years
- This policy must be approved by the MEC
- Keep up to date nursing care plan

CMS Proposes New Changes to Nursing

- All nurses, including agency nurses, must follow all P&Ps
- Must provide for supervision of all nurses, including agency nurses, and evaluation of all clinical activities
 - CNO must provide this
- Document all complications
- Want to refer to CNO as nursing leadership
- Final diagnosis in chart in outpatient within 7 days

CMS Three Worksheets Discharge Planning, Infection Control and QAPI



CMS Hospital Worksheets History

- October 14, 2011 CMS issues a 137 page memo in the survey and certification section and it was pilot tested in hospitals in 11 states
- Memo discusses surveyor worksheets for hospitals by CMS during a hospital survey
- Addresses discharge planning, infection control, and QAPI (performance improvement)
 - Had 3 pilots
 - Discharge planning one will be revised in 2016 when proposed discharge planning standards are final
- Final ones issued November 26, 2014

Final 3 Worksheets QAPI

DEPARTMENT OF HEALTH & HUMAN SERVICES
Centers for Medicare & Medicaid Services
7500 Security Boulevard, Mail Stop C2-21-16
Baltimore, Maryland 21244-1850



Center for Clinical Standards and Quality/Survey & Certification Group

REF: S&C: 15-12-Hospital

DATE: November 26, 2014

TO: State Survey Agency Directors

www.cms.gov/SurveyCertificationGroupInfo/PMSR/list.asp#TopOfPage

FROM: Director
Survey and Certification Group

SUBJECT: Public Release of Three Hospital Surveyor Worksheets

Memorandum Summary

- **Three Hospital Surveyor Worksheets Finalized:** The Centers for Medicare & Medicaid Services (CMS) has finalized surveyor worksheets for assessing compliance with three Medicare hospital Conditions of Participation (CoPs): Quality Assessment and Performance Improvement (QAPI), Infection Control, and Discharge Planning. The worksheets are used by State and Federal surveyors on all survey activity in hospitals when assessing compliance with any of these three CoPs.
- **Final Worksheets Made Public:** Via this memorandum we are making the worksheets publicly available. The hospital industry is encouraged, but not required, to use the worksheets as part of their self-assessment tools to promote quality and patient safety.

CMS Hospital Worksheets

- Hospitals should be familiar with the three worksheets
- Will use whenever a validation survey or certification survey is done at a hospital by CMS
- CMS says worksheets are used by State and federal surveyors on all survey activity in assessing compliance with any of the three CoPs
- Hospitals are encouraged by CMS to use the worksheet as part of their self assessment tools which can help promote quality and patient safety

CMS Hospital Worksheets

- Some of the questions asked might not be apparent from a reading of the CoPs
- So the worksheets are a good communication device
- It helps to clearly communicate to hospitals what is going to be asked in these 3 important areas
- Hospitals might want to consider putting together a team to review the 3 worksheets and complete the form in advance as a self assessment
- Hospitals should consider attaching the documentation and P&P to the worksheet

CMS Survey Memos

ISMP IV Push Guidelines



CMS Survey Memos

- CMS has many recent memos of interest
 - Privacy and confidentiality, CRE and ERCPs
 - Luer misconnections, IV and blood and blood products
 - Use of insulin pens issue, EMTALA and EBOLA
 - Single dose vials and safe injection practices
 - Humidity in the OR, Infection control 4 breaches
 - Discharge planning, Safe Injection Practices
 - Complaint manual and reporting to AO
 - Deficiencies of hospitals, Equipment Maintenance
 - OPO, Medication and Safe Opioid Use, Compounding

CMS Memo on Insulin Pens

- CMS issues memo on insulin pens on May 18, 2012
- Insulin pens are intended to be used on one patient only
- CMS notes that some healthcare providers are not aware of this
- Insulin pens were used on more than one patient which is like sharing needles
- Every patient must have their own insulin pen
- Insulin pens must be marked with the patient's name

Insulin Pens

DEPARTMENT OF HEALTH & HUMAN SERVICES
Centers for Medicare & Medicaid Services
7000 Security Boulevard, Mail Stop C2-21-16
Baltimore, Maryland 21244-1882



Office of Clinical Standards and Quality/Survey & Certification Group

Ref: S&C: 12-30-ALL

DATE: May 18, 2012

TO: State Survey Agency Directors

[www.cms.gov/Medicare/Provider-Enrollment-and-](http://www.cms.gov/Medicare/Provider-Enrollment-and-Certification/SurveyCertificationGenInfo/Policy-and-Memos-to-States-and-Regions.html)

FROM: Director
Survey and Certification Group

[Certification/SurveyCertificationGenInfo/Policy-and-Memos-to-States-and-Regions.html](http://www.cms.gov/Medicare/Provider-Enrollment-and-Certification/SurveyCertificationGenInfo/Policy-and-Memos-to-States-and-Regions.html)

SUBJECT: Use of Insulin Pens in Health Care Facilities

Memorandum Summary

Insulin Pen devices: The Centers for Medicare & Medicaid Services (CMS) has recently received reports of use of insulin pens for more than one patient, with at least one 2011 episode resulting in the need for post-exposure patient notification. These reports indicate that some healthcare personnel do not adhere to safe practices and may be unaware of the risks these unsafe practices pose to patients. Insulin pens are meant for use by a single patient only. Each patient/resident must have his/her own. Sharing of insulin pens is essentially the same as sharing needles or syringes, and must be cited, consistent with the applicable provider/supplier specific survey guidance, in the same manner as re-use of needles or syringes.

Background

Insulin pens are pen-shaped injector devices that contain a reservoir for insulin or an insulin cartridge. These devices are designed to permit self-injection and are intended for single-person use. In healthcare settings, these devices are often used by healthcare personnel to administer insulin to patients. Insulin pens are designed to be used multiple times by a single patient/resident, using a new needle for each injection. Insulin pens must never be used for more than one patient/resident. The reuse of insulin pens, such as the reuse of insulin pens, needles, or syringes, is prohibited.

CMS Memo on Insulin Pens

- Regurgitation of blood into the insulin cartridge after injection can occur creating a risk if used on more than one patient
- Hospital needs to have a policy and procedure
- Staff should be educated regarding the safe use of insulin pens
- More than 2,000 patients were notified in 2011 because an insulin pen was used on more than one patient
- CDC issues reminder on same and has free flier

CDC Reminder on Insulin Pens

Injection Safety [www.cdc.gov/injectionsafety/clinical-reminders/insulin-](http://www.cdc.gov/injectionsafety/clinical-reminders/insulin-pens.html)

Injection Safety

CDC's Role

CDC Statement

Information for Providers

Information for Patients

Preventing Unsafe Injection Practices

Infection Prevention during Blood Glucose Monitoring and Insulin Administration

FAQs regarding Assisted Blood Glucose Monitoring and Insulin Administration

CDC Clinical Reminder: Fingerstick Devices

► **Clinical Reminder: Insulin Pens**

Recent Publications

Recent Meetings

The One & Only Campaign

[Injection Safety](#)

> [Infection Prevention during Blood Glucose Monitoring and Insulin Administration](#)

[Recommend](#) [Tweet](#) 40 [Share](#)

CDC Clinical Reminder: Insulin Pens Must Never Be Used for More than One Person

Available for download [Clinical Reminder: Insulin Pens](#) [PDF - 182 KB]

Summary

The Centers for Disease Control and Prevention (CDC) has become increasingly aware of reports of improper use of insulin pens, which places individuals at risk of infection with pathogens including hepatitis viruses and human immunodeficiency virus (HIV). This notice serves as a reminder that insulin pens must **never** be used on more than one person.

Background

Insulin pens are pen-shaped injector devices that contain a reservoir for insulin or an insulin cartridge. These devices are designed to permit self-injection and are intended for single-person use. In healthcare settings, these devices are often used by healthcare personnel to administer insulin to patients. Insulin pens are designed to be used multiple times, for a single person, using a new needle for each injection. Insulin pens must **never** be used for more than one person.

[Email page link](#)

[Print page](#)

Get email updates

To receive email updates about this page, enter your email address:

[What's this?](#)

Contact Us:

 Centers for Disease Control and Prevention
1600 Clifton Rd
Atlanta, GA 30333

 800-CDC-INFO
(800-232-4636)
TTY: (888) 232-6348

[Contact CDC-INFO](#)

Related Links

[One & Only Campaign](#)

CDC Has Flier for Hospitals on Insulin Pens

CDC CLINICAL REMINDER

Insulin Pens Must Never Be Used for More than One Person

Summary

The Centers for Disease Control and Prevention (CDC) has become increasingly aware of reports of improper use of insulin pens, which places individuals at risk of infection with pathogens including hepatitis viruses and human immunodeficiency virus (HIV). This notice serves as a reminder that insulin pens must **never** be used on more than one person.

Background

Insulin pens are pen-shaped injector devices that contain a reservoir for insulin or an insulin cartridge. These devices are designed to permit self-injection and are intended for single-person use. In healthcare settings, these devices are often used by healthcare personnel to administer insulin to patients. Insulin pens are designed to be used multiple times, for a single person, using a new needle for each injection. Insulin pens must **never** be used for more than one person. Regurgitation of blood into the insulin cartridge can occur after injection [1] creating a risk of bloodborne pathogen transmission if the pen is used for more than one person, even when the needle is changed.

In 2009, in response to reports of improper use of insulin pens in hospitals, the Food and Drug Administration (FDA) issued an alert for healthcare professionals reminding them that insulin pens are meant for use on a single patient only and are not to be shared between patients [2]. In spite of this alert, there have been continuing reports of patients placed at risk through inappropriate reuse and sharing of insulin pens, including an incident in 2011 that required notification of more than 2,000 potentially exposed patients [3]. These events indicate that some healthcare personnel do not adhere to safe practices and may be unaware of the risks these unsafe practices pose to patients.

Recommendations



Insulin Pen Posters and Brochures Available



About the Campaign

Safe Injection Practices

Healthcare Provider Information

Patient Information

Campaign Resources

News

Contact Us



Insulin Pen Safety – One Insulin Pen, One Person

BE AWARE
DON'T SHARE



**ONE INSULIN PEN,
ONLY ONE PERSON**

[www.oneandonlycampaign.org
/content/insulin-pen-safety](http://www.oneandonlycampaign.org/content/insulin-pen-safety)

The Safe Injection Practices Coalition created an insulin pen poster and brochure for healthcare providers as a reminder that insulin pens and other injectable medications are meant for one person and should never be shared. PDFs of these educational materials are linked below:

Specific Materials for Safe Use of Insulin Pens – for Clinicians and Patients


- [Poster](#)
- [Brochure](#)

[Click here](#) to order free copies of these materials from the Centers for Disease Control and Prevention (CDC) (publication numbers 22-1501 and 22-1503).

Additional Resources

- [VA Patient Safety Alert: Multi-Dose Pen Injectors](#) (Department of Veterans Affairs, January 2013)

BE AWARE DON'T SHARE



Insulin pens that contain more than one dose of insulin are only meant for one person.

They *should never be used for more than one person*, even when the needle is changed.

**ONE INSULIN PEN,
ONLY ONE PERSON**

The One & Only Campaign is a public health campaign aimed at raising awareness among the general public and healthcare providers about safe injection practices.

For more information,
please visit:
www.ONEandONLYcampaign.org

Brochure

DON'T DO IT

Sharing Insulin Pens and Other Injection Equipment Jeopardizes Patients

In 2009, in response to reports of improper use of insulin pens in hospitals, the Food and Drug Administration issued an alert for healthcare professionals reminding them that insulin pens are meant for use on a single person only and are not to be shared. Unfortunately, there have been continuing reports of persons placed at risk of bloodborne and bacterial pathogen transmission through sharing of insulin pens.

A SIMPLE RULE

Injection equipment (e.g., insulin pens, needles and syringes) should **never** be used for more than one person.



About the Safe Injection Practices Coalition

The Safe Injection Practices Coalition (SIPC) is a partnership of healthcare-related organizations led by the Centers for Disease Control and Prevention that was formed to promote safe injection practices in all U.S. healthcare settings. The SIPC has developed the One & Only Campaign – a public health education and awareness campaign – aimed at both healthcare providers and patients to advance and promote safe injection practices.

For more information,
please visit:

www.ONEandONLYcampaign.org

BE AWARE DON'T SHARE



ONE INSULIN PEN, ONLY ONE PERSON



What Every
Healthcare Professional
Needs To Know

Recommendations for Safe Insulin Pen Use

Protection from infection is a basic expectation anywhere healthcare is delivered. Use of insulin pens and other injection equipment for more than one person poses unacceptable risks and should be considered a "never" event.

- Insulin pens and other injection equipment containing multiple doses of medication are meant for use on a single person only, and should never be used for more than one person, even when the needle is changed.
- Insulin pens and other injection equipment should be clearly labeled with the person's name or other identifying information to ensure that the correct pen is used only on the correct individual.
- Hospitals and other facilities should review their policies and educate their staff regarding safe use of insulin pens and similar devices.
- If reuse is identified, exposed persons should be promptly notified and offered appropriate follow-up including bloodborne pathogen testing.


These recommendations apply to any setting where insulin pens and other injection equipment are used, including assisted living or residential care facilities, skilled nursing facilities, clinics, health fairs, shelters, detention facilities, senior centers, schools, and camps as well as licensed healthcare facilities.



ONE INSULIN PEN, ONLY ONE PERSON

Insulin Administration

Insulin pens are pen-shaped injector devices that contain a reservoir for insulin or an insulin cartridge. These devices are designed to permit self-injection. They are intended for single-person use.




In healthcare settings, these devices are often used by healthcare personnel to administer insulin to patients. Insulin pens are designed to be used for a single person multiple times, using a new needle for each injection.

Back flow of blood into the insulin reservoir can occur during an injection. This creates a risk of bloodborne and bacterial pathogen transmission if the pen is used for more than one person, even when the needle is changed.

The Safe Injection Practices Coalition created an easy to use check list for facilities. Similar to a risk assessment, the list contains the necessary components of injection safety for facilities to quickly assess their practices.

A copy of the checklist can be found at www.cdc.gov/injectionsafety/Checklist



CMS Memo May 30, 2014

- CMS publishes 4 page memo on infection control breaches and when they warrant referral to the public health authorities
- This includes a finding by the state agency (SA), like the Department of Health, or an accreditation organization
 - TJC, DNV Healthcare, CIHQ, or AOA HFAP
- CMS has a list and any breaches should be referred
- Referral is to the state authority such as the state epidemiologist or State HAI Prevention Coordinator

Infection Control Breaches

DEPARTMENT OF HEALTH & HUMAN SERVICES
Centers for Medicare & Medicaid Services
7500 Security Boulevard, Mail Stop C2-21-16
Baltimore, Maryland 21244-1850



Center for Clinical Standards and Quality/Survey & Certification Group

Ref: S&C: 14-36-All

DATE: May 30, 2014

TO: State Survey Agency Directors

FROM: Director
Survey and Certification Group

SUBJECT: Infection Control Breaches Which Warrant Referral to Public Health Authorities

Memorandum Summary

- ***Infection Control Breaches Warranting Referral to Public Health Authorities:*** If State Survey Agencies (SAs) or Accrediting Organizations (AOs) identify any of the breaches of generally accepted infection control standards listed in this memorandum, they should refer them to appropriate State authorities for public health assessment and management.
- ***Identification of Public Health Contact:*** SAs should consult with their State's Healthcare Associated Infections (HAI) Prevention Coordinator or State Epidemiologist on the preferred referral process. Since AOs operate in multiple States, they do not have to confer with State public health officials to set up referral processes, but are expected to refer identified breaches to the appropriate State public health contact identified at: <http://www.cdc.gov/HAI/state-based/index.html>

CMS Memo Infection Control Breaches

- If any of the listed breaches are observed, then will take appropriate enforcement action
- And will make the public health authority aware
 - Includes LTC, ASCs, hospice, hospitals, home health agencies, CAH, rural health clinics and dialysis facilities
- CDC is working closely with SA on HAI prevention
- List of breaches to be referred include:
- Using the same needle for more than one individual;

CMS Memo Infection Control Breaches

- Using the same (pre-filled/manufactured/insulin or any other) syringe, pen or injection device for more than one individual
- Re-using a needle or syringe which has already been used to administer medication to an individual to subsequently enter a medication container (e.g., vial, bag), and then using contents from that medication container for another individual
- Using the same lancing/fingerstick device for more than one individual, even if the lancet is changed
- CMS also issued EBOLA and CRE memos

CRE and ERCP's

DEPARTMENT OF HEALTH & HUMAN SERVICES
Centers for Medicare & Medicaid Services
7500 Security Boulevard, Mail Stop C2-21-16
Baltimore, Maryland 21244-1850



Center for Clinical Standards and Quality/Survey & Certification Group

Ref: S&C-15-32 Hospitals/CAHs/ASCs

DATE: April 3, 2015

TO: State Survey Agency Directors

FROM: Director
Survey and Certification Group

SUBJECT: Alert Related to Outbreaks of Carbapenem-Resistant *Enterobacteriaceae* (CRE) during gastrointestinal endoscopy, particularly Endoscopic Retrograde Cholangiopancreatography (ERCP)

Memorandum Summary

- **Situation:** Recent newspaper articles, medical publications, and adverse event reports associate multidrug-resistant bacterial infections caused by CRE with patients who have undergone ERCP. Duodenoscopes used to perform ERCP are difficult to clean and disinfect, even when manufacturer reprocessing instructions are followed correctly, and have been implicated in these outbreaks. The U.S. Food and Drug Administration (FDA) has issued a Safety Communication warning, with related updates, that the design of duodenoscopes may impede effective cleaning.
- **Expectations for Reprocessing Duodenoscopes:** Hospitals, critical access hospitals (CAHs), and ambulatory surgical centers (ASCs) are expected to meticulously follow the manufacturer's instructions for reprocessing duodenoscopes, as well as adhere to the nationally recognized Multisociety consensus guidelines developed by multiple expert organizations and issued in 2011.

3 EBOLA Memos Issued

DEPARTMENT OF HEALTH & HUMAN SERVICES
Centers for Medicare & Medicaid Services
7500 Security Boulevard, Mail Stop C2-21-16
Baltimore, Maryland 21244-1850



Center for Clinical Standards and Quality/Survey & Certification Group

Ref: S&C: 15-24-Hospitals

DATE: February 13, 2015
TO: State Survey Agency Directors
FROM: Director
Survey and Certification Group
SUBJECT: Emergency Medical Treatment and Labor Act (EMTALA) and Ebola Virus Disease (EVD) – Questions and Answers (Q+A)

Memorandum Summary

EMTALA & Ebola Requirements:

- On November 21, 2014 the Centers for Medicare & Medicaid Services (CMS) Survey & Certification Group released SC 15-10-Hospitals concerning EMTALA Requirements and Implications Related to the EVD.
- The CMS has received follow-up questions regarding EMTALA and Ebola and has produced a Q+A document in response.

The CMS released S&C 15-10 on November 21, 2014 to provide guidance to hospitals and critical access hospitals (CAHs) regarding meeting EMTALA requirements in the case of individuals potentially exposed to Ebola. The memo is available via the following link:

<http://www.cms.gov/Medicare/Provider-Enrollment-and->

CMS Memo on Safe Injection Practices

- CMS issues a 7 page memo on safe injection practices
- Discusses the safe use of single dose medication to prevent healthcare associated infections (HAI)
- Notes new exception which is important especially in medications shortages
- General rule is that single dose vial (SDV) can only be used on one patient
- Will allow SDV to be used on multiple patients if prepared by pharmacist under laminar hood following USP 797 guidelines

Single Dose Safe Injection Practices

DEPARTMENT OF HEALTH & HUMAN SERVICES
Center for Medicare & Medicaid Services
1100 Security Boulevard, Mail Stop C2-21-18
Baltimore, Maryland 21244-1880



Office of Clinical Standards and Quality/Survey & Certification Group

Ref: S&C: 12-35-ALL

DATE: June 15, 2012

TO: State Survey Agency Directors

FROM: Director
Survey and Certification Group

SUBJECT: Safe Use of Single Dose/Single Use Medications to Prevent Healthcare-associated Infections

Memorandum Summary

- *Under certain conditions, it is permissible to repackage single-dose vials or single use vials (collectively referred to in this memorandum as "SDV's") into smaller doses, each intended for a single patient.* The United States Pharmacopeia (USP) has established standards for compounding which, to the extent such practices are also subject to regulation by the Food and Drug Administration (FDA), may also be recognized and enforced under §§501 and 502 of the Federal Food, Drug and Cosmetics Act (FDCA). These USP compounding standards include USP General Chapter 797, *Pharmaceutical Compounding - Sterile Preparations* ("USP <797>"). Under USP <797>, healthcare facilities may repackage SDVs into smaller doses, each intended for use with one patient. Among other things, these standards currently require that:
 - The facility doing the repackaging must use qualified, trained personnel to do so, under International Organization for Standardization (ISO) Class 5 air quality conditions within an ISO Class 7 buffer area. All entries into a SDV for purposes of repackaging under these conditions must be completed within 6 hours of the initial needle puncture.
 - All repackaged doses prepared under these conditions must be assigned and labeled with a beyond use date (BUD), based on an appropriate determination of contamination risk level in accordance with USP <797>, by the licensed healthcare professional supervising the repackaging process.
- *Administering doses from one SDV to multiple patients without adhering to USP <797>*

CMS Memo on Safe Injection Practices

- All entries into a SDV for purposes of repackaging must be completed with 6 hours of the initial puncture in pharmacy following USP guidelines
- Only exception of when SDV can be used on multiple patients
- Otherwise using a single dose vial on multiple patients is a violation of CDC standards
- CMS will cite hospital under the hospital CoP infection control standards since must provide sanitary environment
 - Also includes ASCs, hospice, LTC, home health, CAH, dialysis, etc.

CMS Memo on Safe Injection Practices

- Bottom line is you can not use a single dose vial on multiple patients
- CMS requires hospitals to follow nationally recognized standards of care like the CDC guidelines which has 10 practices
- SDV typically lack an antimicrobial preservative
- Once the vial is entered the contents can support the growth of microorganisms
- The vials must have a beyond use date (BUD) and storage conditions on the label

CMS Memo on Safe Injection Practices

- Make sure pharmacist has a copy of this memo
- If medication is repackaged under an arrangement with an off site vendor or compounding facility ask for evidence they have adhered to 797 standards
- ASHP Foundation has a tool for assessing contractors who provide sterile products
- Go to
www.ashpfoundation.org/MainMenuCategories/PracticeTools/SterileProductsTool.aspx
- Click on starting using sterile products outsourcing tool now

CMS Memo on Safe Injection Practices

- So if make it in a single dose vial then you need to buy it in a single dose vial
 - If they only make it in a multi-dose vial then try and use it as a single dose vial
 - If not then try and use it only on one patient
- Do not take multi-dose vial into patient room or into OR
 - Unless in OR you treat it as a single dose vial and discard
 - Mark multi-dose vial expires in 28 days unless sooner by manufacturer
 - Clean off lid even if new vial for 10-15 seconds and let dry or as per manufacturer instructions (new one is 5 seconds)

Print this page Email this page

Advancing Practice

Optimizing Antithrombotic Management: An Assessment Tool

Bar Code Guide

My Medicine List™

Outsourcing Sterile Products Preparation: Contractor Assessment Tool

Pharmacy Practice Model Initiative

Outsourcing Sterile Products Preparation: Contractor Assessment Tool

Developed with support from PharMEDium Services, LLC
Now available!

Preparation of sterile parenteral products is a critical component of health-system pharmacy practice. For departments that choose to outsource the preparation of parenteral medications, this web-based tool can be used to evaluate proposals during the selection of an external organization that would provide parenteral product preparation services.

The assessment tool helps you evaluate each of these areas:

- Regulatory compliance
- Quality and patient safety measures
- Medication administration safety features
- Service excellence

Start using the Sterile Products Outsourcing Tool now!



www.ashpfoundation.org/MainMenuCategories/PracticeTools/SterileProductsTool.aspx

Safe Injection Practices



EMERGENCY
MEDICINE
PATIENT SAFETY
FOUNDATION

Safe Injection Practices Patient Safety Brief
Emergency Medicine Patient Safety Foundation


By: Sue Dill Calloway RN MSN JD CPHRM
Ruth Carrico PhD RN FSHEA CIC

July 2012



Unsafe Injection Practices and Disease Transmission

Reuse of syringes combined with the use of single-dose vials for multiple patients undergoing anesthesia can transmit infectious diseases. The syringe does not have to be used on multiple patients for this to occur.



Source: www.southernnevadahealthdistrict.org

Not All Vials Are Created Equal

SINGLE-DOSE OR MULTI-DOSE?

NOT ALL VIALS ARE CREATED EQUAL.

Dozens of recent outbreaks have been associated with reuse of single-dose vials and misuse of multiple-dose vials. As a result of these incidents, patients have suffered significant harms, including death. CDC and the One & Only Campaign urge healthcare providers to recognize the differences between single-dose and multiple-dose vials and to understand appropriate use of each container type.

This information can literally save a life.



ONEANDONLYCAMPAIGN.ORG

DO YOU PROVIDE TREATMENT FOR PATIENTS WITH CANCER?

PROTECT YOUR PATIENTS, YOURSELF, AND YOUR BUSINESS

Since 2002, at least nine serious infectious disease outbreaks have occurred in cancer clinics. These outbreaks involved unsafe injection practices, including the reuse of syringes. As a result, hundreds of patients became infected and thousands more required notification and testing for bloodborne pathogens.



REMEMBER! WHEN PREPARING MEDICATIONS AND INJECTIONS...

NEVER reuse these items:



Needles or syringes that have been used for any purpose



Vials with "single-dose vial" printed on the label



Saline bags



Intravenous tubing

ALWAYS follow aseptic technique* when:



Preparing any medication



Disinfecting a vial's septum



Accessing a central line



Injecting any medications

*Aseptic technique is used by health care workers to prevent the contamination of clean areas, equipment, and sterile medications. This will help prevent the spread of infection. Please refer to [CDC's Basic Infection Control and Prevention Plan for Outpatient Oncology Settings](#) for more information.

LEARN MORE ABOUT WAYS YOU CAN KEEP YOUR PATIENTS

1 ONE NEEDLE,
ONE SYRINGE,
ONLY ONE TIME.



ISMP IV Push Medications Guidelines

- ISMP has published a 26 page document called “ISMP Safe Practice Guidelines for Adult IV Push Medications
- The document is organized into factors that increase the risk of IV push medications in adults,
 - Current practices with IV injectible medications
 - Developing consensus guidelines for adult IV push medication and
 - Safe practice guidelines
 - About 90% of all hospitalized patients have some form of infusion therapy

IV Push Medicine Guidelines

ISMP Safe Practice Guidelines for Adult IV Push Medications

A compilation of safe practices from the
ISMP Adult IV Push Medication Safety Summit

Remember; CMS says you have to follow standards of care and specifically mentions the ISMP so surveyor can cite you if you do not follow this.

Prepared by the Institute for
Safe Medication Practices (ISMP)



IV Push Medications Guidelines

- Provide IV push medications in a ready to administer form
- Use only commercially available or pharmacy prepared prefilled syringes of IV solutions to flush and lock vascular access devices
- If available in a single dose vial then need to buy in single dose vial
- Aseptic technique should be used when preparing and administering IV medication
 - This includes hand hygiene before and after administration

IV Push Medications Guidelines

- The diaphragm on the vial should be disinfected even if newly opened
 - The top should be cleaned using friction and a sterile 70% isopropyl alcohol, ethyl alcohol, iodophor, or other approved antiseptic swab for at least ten seconds to it dry
 - Medication from glass ampules should be used with a filter needle unless the specific drug precludes this
- Medication should only be diluted when recommended by the manufacturer or in accordance with evidence based practice or approved hospital policies

IV Push Medications Guidelines

- If IV push medication needs to be diluted or reconstituted these should be performed in a clean, uncluttered, and separate location
- Medication should not be withdrawn from a commercially available, cartridge type syringe into another syringe for administration
- It is also important that medication not be drawn up into the commercially prepared and prefilled 0.9% saline flushes
 - This are to flush an IV line and are not approved to use to dilute medication

3.6 Do NOT dilute or reconstitute IV push medications by drawing up the contents into a commercially-available, prefilled flush syringe of 0.9% sodium chloride.

Discussion: Commercially available prefilled syringes of saline and heparin are regulated by the US Food and Drug Administration as *devices*, not as medications. These devices have been approved for the flushing of vascular access devices, but have NOT been approved for the reconstitution, dilution, and/or subsequent administration of IV push medications. Such use would be considered “off label” and not how manufacturers intended these products to be used, nor have prefilled flush syringes been tested for product safety when used in this manner.

Warnings intended to limit the use of prefilled syringes for medication preparation and administration appear on some syringe barrels, clearly stating “IV flush only.” Some manufacturers have also limited or removed the gradation markings on the prefilled flush syringes in order to prevent measurement of a secondary medication in the flush syringe. When prefilled syringes are used in an off-label manner, the practitioner and employer bear the legal liability for any adverse events occurring from this practice.³¹

The mislabeling that occurs when medications are added to a prefilled syringe and a secondary label is not applied creates significant risk for errors. In many cases, the manufacturer’s label is permanently affixed to the syringe barrel and contains product codes and a barcode as well as specific information about the fluid and its volume. When another medication is added to this syringe, there is no adequate method to amend the manufacturer’s label, without covering the current information.³¹ Thus, the syringe frequently remains labeled as 0.9% sodium chloride, when it also contains the diluted or reconstituted medication.

Although this unsafe practice is widespread, and many who use it mistakenly believe the risk of an error is insignificant—a belief clearly reinforced during public comment regarding this guidance statement—summit participants arrived at a consensus that the practice must be eliminated.

3.7 When necessary to prepare more than one medication in a single syringe for IV push administration

IV Push Medications Guidelines

- Combination of more than one medication in a single syringe is seldom necessary and could result in unwanted changes in the medication
- Never use IV solution or mini bags as a common source to flush an IV as to dilute for more than one patient
- Label syringes of IVP medication unless prepared and immediately given with no break
- Administer IV push medication at rate recommended by manufacturer or supported by evidenced based practices and often given too fast

CMS Memo 1 of 2 on RH

- CMS issues memo related to the relative humidity (RH)
- AORN use to say temperature maintained between 68-73 degrees and humidity between 30-60% in OR, PACU, cath lab, endoscopy rooms and instrument processing areas
- CMS says if no state law can write policy or procedure or process to implement the waiver
- Waiver allows RH between 20-60%
- In anesthetizing locations- see definition in memo

Humidity in Anesthetizing Areas

DEPARTMENT OF HEALTH & HUMAN SERVICES
Centers for Medicare & Medicaid Services
7500 Security Boulevard, Mail Stop C2-21-16
Baltimore, Maryland 21244-1820



Center for Clinical Standards and Quality / Survey & Certification Group

Ref: S&C: 13-25-LSC & ASC

DATE: April 19, 2013
TO: State Survey Agency Directors
FROM: Director
Survey and Certification Group
SUBJECT: Relative Humidity (RH): Waiver of Life Safety Code (LSC) Anesthetizing Location Requirements; Discussion of Ambulatory Surgical Center (ASC) Operating Room Requirements

Memorandum Summary

- ***RH of ≥20 Percent Permitted in Anesthetizing Locations:*** The Centers for Medicare & Medicaid Services (CMS) is issuing a categorical LSC waiver permitting new and existing ventilation systems supplying hospital and critical access hospital (CAH) anesthetizing locations to operate with a RH of ≥20 percent, instead of ≥35 percent. We are also recommending that RH not exceed 60 percent in these locations.
- ***This Waiver Does Not Apply:***
 - When more stringent RH control levels are required by State or local laws and regulations; or
 - Where reduction in RH would negatively affect ventilation system performance.
- ***Hospitals & CAHs Must Elect to Use the Categorical Waiver:***
 - Individual waiver applications are not required, but facilities are expected to have written documentation that they have elected to use the waiver.
 - At the entrance conference for any survey assessing LSC compliance, a facility that has elected to use this waiver must notify the survey team.
- ***Ongoing Requirements:***
 - Facilities must monitor RH in anesthetizing locations and take corrective actions when needed to ensure RH remains at or above 20 percent.
- ***ASCs:*** ASCs are not subject to all of the same LSC requirements as hospitals, but are required, consistent with 42 CFR 416.44(a)(1), to maintain RH in operating rooms in accordance with nationally accepted guidelines.
- ***State Operations Manual (SOM) Appendices A, E, F, & H*** are being updated accordingly.

Impact of Lowering the Humidity

- Lowering humidity can impact some equipment and supplies
- Can affect shelf life and product integrity of some sterile supplies including EKG electrodes
- Some electro-medical equipment may be affected by electrostatic discharge especially older equipment
 - Can cause erratic behavior of software and premature failure of the equipment
 - It can affect calibration of the equipment
- Follow the manufacturers instructions for use that explains any RH requirements

CMS Memo on Low Relative Humidity

DEPARTMENT OF HEALTH & HUMAN SERVICES
Centers for Medicare & Medicaid Services
7500 Security Boulevard, Mail Stop C2-21-16
Baltimore, Maryland 21244-1850



Center for Clinical Standards and Quality / Survey & Certification Group

Ref: S&C: 15-27-Hospital, CAH & ASC

DATE: February 20, 2015

TO: State Survey Agency Directors

FROM: Director
Survey and Certification Group

SUBJECT: Potential Adverse Impact of Lower Relative Humidity (RH) in Operating Rooms (ORs)

Memorandum Summary

- **Information on OR RH** is provided for Ambulatory Surgical Centers (ASCs) & Supplemental Information for Hospitals & Critical Access Hospitals (CAHs) Using the Categorical Waiver of Life Safety Code (LSC) Anesthetizing Location RH Requirements
 - The Association for the Advancement of Medical Instrumentation (AAMI) coordinated the release on January 5, 2015 of a Joint Communication of multiple healthcare-related organizations on how a RH of <30% in ORs may affect the performance of some sterile supplies and electro-medical equipment.
- **S&C 13-25-LSC & ASC** permits hospitals and CAHs to use a LSC categorical waiver to establish an RH level <35% in anesthetizing locations. Before electing or continuing to use this categorical waiver, hospitals and CAHs are expected to ensure that the humidity levels in their ORs are compatible with the manufacturers' instructions for use (IFUs) for the supplies and equipment used in that setting.
- **ASCs do not require a categorical waiver** in order to use a lower RH level in their ORs but also need to ensure they comply with the IFUs for their OR supplies and equipment.

Impact of Lowering the Humidity



American Hospital Association®



Quality Advisory

January 21, 2015

01-21-2015 Accessed ; https://www.magnetmail.net/actions/email_web_version.cfm?recipient_id=1331564405&message_id=8663272&user_id=AHA_8&group_id=1105177&jobid=25267573

NEW GUIDANCE ON HUMIDITY LEVELS IN THE OPERATING ROOM

THE ISSUE

A change in the standards regulating a hospital's physical environment in the operating room (OR) may conflict with the instructions for use on some equipment and supplies routinely used in surgery. To ensure patient safety during surgery, the AHA in collaboration with its personal membership groups, the American Society for Healthcare Engineering (ASHE) and the Association for Healthcare Resource & Materials Management (AHRMM), urge hospitals to examine their humidity levels in the OR and consider the effects on equipment and products used during surgery. This advisory and associated attachments will assist in your assessment.

BACKGROUND

Many safety codes and standards regulating the health care physical environment now require relative humidity levels in ORs (not other areas of the facility) to be at least 20 percent, a change from the 30 percent minimum humidity required by some previous editions of codes. The 20 percent threshold provides hospitals with flexibility during

Lowering Humidity Can Have Other Effects

RELATIVE HUMIDITY LEVELS IN THE OPERATING ROOM JOINT COMMUNICATION TO HEALTHCARE DELIVERY ORGANIZATIONS January 2015



American Hospital
Association®



This is an important communication to the multiple stakeholders in healthcare whose work touches sterile supplies and electro-medical equipment used in delivering care to patients. The subject is about how relative humidity (RH) levels lower than 30% can impact the integrity and functionality of some of these products, with a special emphasis on RH levels in the operating room (OR). The following professional organizations have collaborated in the development of this communication: Ambulatory Surgery Center Association (ASCA), American College of Clinical Engineering (ACCE), American Hospital Association (AHA), American Society for Healthcare Engineering (ASHE), American Society of Anesthesiologists (ASA), American Society of Heating, Refrigeration and Air Conditioning Engineers (ASHRAE), Association for Healthcare Resource & Materials Management (AHRMM), Association for the Advancement of Medical Instrumentation (AAMI), Association of periOperative Registered Nurses (AORN), Association of Surgical Technologists (AST), Health Industry Distributors Association (HIDA), and the International Association of Healthcare Central Service Materials Management (IAHCSMM).¹

Complaint Manual Update

- CMS issues memo
- CMS updates the Complaint Manual
- Hospital found to be in immediate jeopardy could have a full validation survey if the RO requests it
 - Regional office has discretion
- GAO emphasized need to share complaint information and SA survey finding with the applicable accreditation agency and CMS agrees
 - TJC, DNV, AOA, or CIHQ

Complaint Manual Update

DEPARTMENT OF HEALTH & HUMAN SERVICES
Centers for Medicare & Medicaid Services
7500 Security Boulevard, Mail Stop C2-21-16
Baltimore, Maryland 21244-1850



Center for Clinical Standards and Quality/Survey & Certification Group

Ref: S&C: 13-27-Deemed Providers/Suppliers & Hospitals

DATE: April 19, 2013

TO: State Survey Agency Directors

FROM: Director
Survey and Certification Group

SUBJECT: Update of State Operations Manual (SOM) Chapter 5, Complaint Investigation

Memorandum Summary

Post-Complaint Survey Procedure - Deemed Providers/Suppliers:

- A full survey of a deemed provider/supplier after a complaint survey with condition-level findings will be made on a selective rather than an automatic basis.
- All survey reports and related correspondence must be shared promptly with a deemed provider/supplier's accrediting organization (AO).

Hospital Restraint/Seclusion Death Reporting: This section is being moved, to reflect the fact that the procedures therein apply to all hospitals, not just deemed hospitals. We are also streamlining the procedure for making disclosures to State Protection and Advocacy (P&A) agencies, to reduce burden.

A. Full Survey After Complaint for Deemed Providers/Suppliers

Luer Misconnections Memo

- CMS issues memo on luer misconnections
- This has been a patient safety issues for many years
- Staff can connect two things together that do not belong together because the ends match
- For example, a patient had the blood pressure cuff connected to the IV and died of an air embolism
- Luer connections easily link many medical components, accessories and delivery devices

Luer Misconnections Memo

DEPARTMENT OF HEALTH & HUMAN SERVICES
Center for Medicare & Medicaid Services
7500 Security Boulevard, Mail Stop C2-23-18
Baltimore, Maryland 21244-1890



Center for Clinical Standards and Quality/Survey & Certification Group

Re: S&C: 13-14-ALL

DATE: March 8, 2013
TO: State Survey Agency Directors
FROM: Director
Survey and Certification Group
SUBJECT: Luer Misconnection Adverse Events

Memorandum Summary

- **Luer Misconnections continue to result in adverse events and deaths –** Luer connectors easily link many medical components, accessories, and delivery systems. Clinicians, in any type of provider or supplier setting, can mistakenly connect the wrong devices and deliver substances through the wrong route. Despite numerous alerts and warnings, a patient's blood pressure tubing was recently misconnected to an intravenous (IV) line in an ambulatory surgery center (ASC) resulting in a patient death.
- **Adverse Event Complaint Investigation:** During a complaint investigation for an adverse event involving delivery of an incorrect substance or utilization of an incorrect delivery route, surveyors must be alert to whether the event involved misconnection of a Luer device. If so, surveyors must determine whether the facility has taken actions to ensure systems are in place to prevent recurrence of this type of adverse event.
- **Facility Reporting to Food & Drug Administration (FDA):** Surveyors should encourage health care facilities to report problems with Luer misconnections to the FDA, even if no adverse event occurred.

PA Patient Safety Authority Article

Table. Tubing Misconnections Reported to the Pennsylvania Patient Safety Authority, January 2008 to September 2009

MISCONNECTION	NUMBER OF REPORTS
Secondary intravenous (IV) infusion connected to lower "Y" port of primary IV tubing set	8
Hemodialysis arterial and venous tubing lines reversed	5
G-tube and J-tube lines reversed	3
Incorrect tubing connection (no further explanation provided in reports)	3
Epidural and patient-controlled analgesia (PCA) tubing sets reversed	2
Nonhemodialysis arterial and venous tubing lines reversed	2
Cell saver tubing connected to cell saver reservoir	1
Feeding tube set connected to Broviac®	1
Feeding tube set connected to peripherally inserted central catheter (PICC) line	1
Feeding tube set connected to suction port	1
Imaging contrast tubing set connected to tracheostomy cuff	1
IV tubing set connected to dialysis catheter	1
IV tubing set connected to PICC line	1
IV tubing set connected to tracheostomy cuff	1
Knee irrigation connected to peripheral IV tubing	1
Miscommunication (arterial line noted in medical record as peripheral IV)	1
Oral medication delivered through peripheral IV line	1
Suction line connected to water seal	1
Suction and feeding tubing sets reversed	1
Total	36

June 2010 Pa Patient Safety Authority

Tubing Mismatches: Making the Connection to Patient Safety

ABSTRACT

Some patients may have multiple tubing lines connected to them for reasons such as delivery of medication and nutrition therapy. With these multiple lines, the potential for tubing mismatches becomes more prevalent. Tubing mismatches can occur with intravenous catheters, feeding tubes, peritoneal dialysis tubes, and tracheostomy cuffs, among other devices. One of the main reasons for tubing mismatches is that many types of tubing for different types of medical devices incorporate lock connectors. These connectors contribute to mismatches because they often have normally dissimilar tubes or catheters to be connected together. Between January 2008 and September 2009, 38 events of tubing mismatches were reported to the Pennsylvania Patient Safety Authority involving various types of mismatches. Methods for reducing the likelihood of tubing mismatches include equipment design solutions and education. New controls (guidance and lock protection), equipment design solutions either prevent the user from making a mismatch or prompt the user to make the correct connection. Administrative controls are policies and practices that reduce the risk of mismatches such as tracing lines back to their source. (Pa Patient Saf Act 2010 Jun;7[2]:41-5.)

Introduction

Depending on acuity level, patients may have multiple lines connecting them to medical devices used for delivering medication or nutrition therapy. Medical devices connected to patients may also have tubing lines connecting the devices with other medical devices. Under these circumstances, tubing mismatches can occur with potentially fatal results. Mismatches have been recognized as a serious problem for many years. One of the earliest published reports of mismatches was the tragic incident delivery of breast milk via intravenous (IV) administration in 1972.¹ However, mismatches have garnered more attention in recent years, especially in the United States, due in part to the tubing mismatched Neutralizer Alarm issued by the Joint Commission in April 2006.²

for IV delivery connected to nasogastric tubes.³ The Alert calls for risk reduction strategies and recommendations, which are included in the overall risk reduction strategies below.

There are many types of mismatches; however, this article will focus on liquid-to-liquid and liquid-to-gas mismatches because these mismatches can pose the most serious harm to patients and are the most frequently reported to the Pennsylvania Patient Safety Authority. Liquid lines are typically those that administer medications or nutrient but can also include solution lines such as flush lines. Medical gas lines are typically used for respiratory support or to power pneumatic medical devices. Liquid-to-liquid mismatches can result in a liquid substance entering the wrong body part or the wrong substance entering the patient. Liquid-to-gas mismatches are incorrect connections that can result in gas entering liquid line patients' blood vessels or liquid entering patients' respiratory tracts.⁴

A common cause for tubing mismatches, whether liquid-to-liquid or liquid-to-gas, is that many types of tubing lines for different medical devices incorporate common lock connectors. The International Unit system for Standardization (ISO) designates a line connector as a central fitting with a 1/8" barrel taper that accepts needles, needles, and other medical equipment.⁵ The line connector system consists of male and female counterparts that are joined together either by push (they slip or screw-in threaded their lock) fittings. Lock connectors contribute to mismatches because they easily allow functionally dissimilar tubes or catheters to be connected together.⁶

Mismatches in Pennsylvania

Between January 2008 and September 2009, 38 tubing mismatch events were reported to the Authority. 35 liquid-to-liquid events and 3 liquid-to-gas events. (See the Table for a breakdown of the types of mismatches reported.) Examples of the Serious Events and Incidents involving mismatches reported to the Authority include the following:

The patient is a 60-year-old female admitted . . . to intensive care for surgery. The physician ordered a 75 ml bolus of 20% parenteral nutrition. A . . . nurse connected the bag of 20% to the patient's lower "Y" on

ISMP Tubing Misconnections

www.ismp.org

ISMP Medication Safety Alert! Acute Care

PREVENTING CATHETER/TUBING MISCONNECTIONS: MUCH NEEDED HELP IS ON THE WAY

From the July 15, 2010 issue

Catheter/tubing misconnections remain a serious problem in healthcare. Just a few weeks ago we learned of another fatal event. Over Memorial Day weekend, a 19-month-old child, who was receiving treatment for a chronic gastrointestinal disorder, died at a pediatric care center. A suspension of **QUESTRAN** (cholestyramine) was accidentally given via a central line intravenous catheter instead of through an enteral feeding tube.

In May 2010, another report was published about barium sulfate being administered via the superior vena cava during upper gastrointestinal study (Soghoian S, Hoffman RS, Nelson L. Unintentional IV injection of barium sulfate in a child. *AJ Health-Syst Pharm.* 2010;67:734-36). The patient, a 17-month-old child, had a central venous catheter (CVC) in place for antibiotic therapy. As the procedure began, approximately 3 mL of barium sulfate was injected into the CVC, which was mistaken as the child's gastrostomy tube. Fortunately, no respiratory distress developed and the child was discharged 3 days later.

Luer connector systems, common to many healthcare catheters, tubes, administration sets, extension sets, and syringe have been at the heart of many catheter/tubing misconnections. At the center of one of the most commonly reported problems is the fact that some manufactured enteral catheters still have ports that only accept parenteral administration sets and syringes. So, even if a liquid medication is prepared in an oral syringe, the medication must be transferred to a parenteral syringe for administration via this type of enteral catheter port, risking the accidental administration of the drug via a parenteral line.

Below are examples of the type of reports we have received associated with catheter/tubing misconnections, all of which we've described in this newsletter since publication began in 1996:

- IV infusions connected to epidural lines, and epidural solutions connected to IV lines
- Syringe containing IV medication given via an intrathecal catheter
- IV tubing connected to inflation balloon port of endotracheal tube or tracheostomy tube
- Sequential compression device tubing or pneumatic blood pressure cuff tubing attached to port of IV administration set
- Oxygen tubing connected to port of IV administration set
- Breast milk intravenously infused into neonates
- Bladder irrigation solutions given IV, or TPN solutions administered via foley catheter port
- IV administration set spiked into enteral nutrition container, resulting in enteral nutrition administered IV.

TJC Sentinel Event Alert #36

www.jointcommission.org

The screenshot shows the homepage of The Joint Commission website. At the top left is the logo for The Joint Commission. To the right of the logo are links for "Log In | Request Guest Access" and "Forgot password? | Log in Help". Further right are links for "Contact Us | Careers | JCR Web Store | Press Room". Below these links is a search bar with the text "Search" and a "Go" button. A navigation menu below the search bar includes "Accreditation", "Certification", "Standards", "Measurement", "Topics", "About Us", and "Daily Update". Below the navigation menu is a breadcrumb trail "Home > Topic Details" and a date "Monday, 9/14/2014, April 23". A sidebar on the left contains a "Sign up for News and Alerts" box with a "Sign up here" button. The main content area is titled "Topic Library Item" and features the heading "Sentinel Event Alert, Issue 36: Tubing misconnections—a persistent and potentially deadly occurrence" with a date of "April 3, 2014". Below the heading is a "Download This File" button. The text below the button reads: "Tubing and airway misconnection errors are an important and under-reported problem in health care organizations. In addition, these errors are often caught and corrected before any injury to the patient occurs. Given the reality of and potential for life-threatening consequences, increased awareness and analysis of these errors—including avoided errors—can lead to dramatic improvement in patient safety." To the right of the text is an image of a tangled piece of clear medical tubing. On the far right of the page is a vertical sidebar with social media icons for Twitter, Facebook, LinkedIn, and a "More" button.

Managing Risk During the Transition

Sentinel Alert *Event*

A complimentary publication of The Joint Commission
Issue 53, August 20, 2014

Published for Joint Commission accredited organizations and interested health care professionals, *Sentinel Event Alert* identifies specific types of sentinel and adverse events and high risk conditions, describes their common underlying causes, and recommends steps to reduce risk and prevent future occurrences.

Accredited organizations should consider information in a *Sentinel Event Alert* when designing or redesigning processes and consider implementing relevant suggestions contained in the alert or reasonable alternatives.

Managing risk during transition to new ISO tubing connector standards

Tubing misconnections continue to cause severe patient injury and death, since tubes with different functions can easily be connected using luer connectors, or connections can be “rigged” (constructed) using adapters, tubing or catheters. This is why new ISO (International Organization for Standardization) tubing connector standards are being developed for manufacturers. Through an international consensus process, the standards are being developed, tested and approved to assure reliable designs and processes. The phased implementation of redesigned tubing connectors that are the result of these new ISO connector standards begins now. The Joint Commission urges health care organizations to be vigilant and begin planning for the upcoming period of transition, which will introduce changes and new risks into the health care environment. Under the new ISO connector standards, small-bore (less than 8.5 mm inner diameter) connectors will be engineered to make it nearly impossible to connect one delivery system to another delivery system that serves a completely different function^{1,2,3,4,5} – for example, accidentally connecting a feeding administration set to a tracheostomy tube, or an intravenous (IV) tube to an epidural site.

The first new ISO connector standard (ANSI/AAMI/ISO 80369-1) has been adopted and others are expected to be introduced and adopted through 2014 and 2015. Health care organizations should begin preparing for changes in connectors and do everything possible during the transitional period to avoid

New Standards Prevent Tubing Misconnections

- New and unique international standards being developed for connectors for gas and liquid delivery systems
- To make it impossible to connect unrelated systems
- Includes new connectors for enteral, respiratory, limb cuff inflation neuraxial, and intravascular systems
- Phase in period for product development, market release and implementation guided by the FDA and national organizations and state legislatures
 - FAQ on small bore connector initiative

New standards to prevent tubing misconnections will have unprecedented impact on supply chain and patient safety



What if you could no longer connect any of the equipment that you have in stock to give enteral feedings (e.g., feeding sets, tubes, oral syringes). That is the likely scenario - once new standards to prevent tubing misconnections are released - without a carefully crafted implementation plan across all settings where care is delivered.

The very simple and universal design of most *connectors* in all of health care creates a serious risk that tubes from totally unrelated systems can be inadvertently connected leading to patient death or serious injury. This means that an enteral feeding tube could be accidentally connected to an IV line, delivering formula into a vein with fatal consequences. An international group of stakeholders are working together to solve this problem by developing unique design standards for every delivery system so that unrelated systems can never be mistakenly connected together.

What do these new standards mean for healthcare

New and unique international standards are being developed for connectors for each gas and liquid delivery system in healthcare to make it virtually impossible to connect unrelated systems¹.

These new connector standards will include new designs for connectors of enteral, respiratory, limb cuff inflation, neuraxial, and intravascular systems. It is anticipated that the standards for enteral connectors will be the first to be released in 2014. There will be a phase-in period for product development, market release and implementation guided by the FDA, existing state legislation, suppliers, and national organizations working together.



www.premierinc.com/tubingmisconnections/

Hospital CoPs for QAPI

- CMS issued new hospital COPs for QAPI
- CMS issues Memo March 15, 2013 on AHRQ Common Formats
 - Hospitals are required to track adverse events for PI
- Starts with tag number 0263
- Short section because the hospital compare program is not part of the CMS CoP
 - Hospital compare is the indicators that must be sent to CMS to receive full reimbursement rates

Report Adverse Events to PI

DEPARTMENT OF HEALTH & HUMAN SERVICES
Centers for Medicare & Medicaid Services
7500 Security Boulevard, Mail Stop C2-21-16
Baltimore, Maryland 21244-1850



Center for Clinical Standards and Quality / Survey & Certification Group

Ref: S&C: 13-19-HOSPITALS

DATE: March 15, 2013
TO: State Survey Agency Directors
FROM: Director
Survey and Certification Group
SUBJECT: AHRQ Common Formats - Information for Hospitals and State Survey Agencies (SAs) - Comprehensive Patient Safety Reporting Using AHRQ Common Formats

Memorandum Summary

Hospitals are Required to Track Adverse Events: The Condition of Participation (CoP) for Quality Assessment and Performance Improvement (QAPI) at 42 CFR 482.21(a)(2) requires hospitals to track adverse patient events. However, several recent reports completed by the Department of Health and Human Services Office of the Inspector General (OIG) indicated that hospitals fail to identify most adverse events.

Use of the Common Formats May Help Hospitals Improve Tracking. The OIG suggested staff failure to understand what events need to be reported to the hospital's QAPI program contributes to the problems with internal tracking systems. The OIG recommended that the Agency for Healthcare Research and Quality (AHRQ) and the Centers for Medicare and Medicaid Services (CMS) could help hospitals improve their ability to track adverse patient safety events by disseminating information on AHRQ's Common Formats. The Common Formats define a systematic process for reporting adverse events, near misses, and unsafe conditions, and allow a hospital to report harm from all causes. Hospital use of the AHRQ Common Formats is voluntary, but a hospital that uses them and is adept at the analysis that they permit will be in a better position

Adverse Event Reporting

- Hospitals are required to track AE
- Several reports show that nurses and others were not reporting adverse events and not getting into the PI system
- OIG recommends using the AHRQ common formats to help with the tracking
- States could help hospitals improve the reporting process
- Encouraged all surveyors to develop an understanding of this tool



- Home
- About the PSOPPC
- Data Submission
- AHRQ Common Formats
- Technical Assistance Resources
- Questions and Answers
- News and Alerts

Welcome to the PSO Privacy Protection Center

The Patient Safety Organization Privacy Protection Center (PSOPPC) was created by the Agency for Healthcare Research and Quality (AHRQ) to support the implementation of the Patient Safety and Quality Improvement Act (PL-109-41), passed by the United States Congress in July, 2005. The PPC provides technical assistance to PSOs by ensuring patient safety event data is non-identifiable for data submission and reporting to the NPSD, and provides technical assistance on use of Common Formats. [Read more about the PPC.](#)



www.psoppc.org/web/patientsafety

Common Formats

General Information

- [About AHRQ Common Formats](#)

Active for Reporting

Hospital Common Formats - v1.2

- [Event Descriptions, Sample Reports, Forms and User's Guide](#)
- [Technical Specifications](#)

Hospital Common Formats - v1.1

- [Event Descriptions, Sample Reports, Forms and User's Guide](#)
- [Technical Specifications](#)

Open for Public Review/Comment

Readmissions Common Format - v0.1 Beta

- [Event Descriptions, Sample Reports, Forms and User's Guide](#)

Skilled Nursing Facility Common Formats - v0.1 Beta

- [Event Descriptions, Sample Reports, Forms and User's Guide](#)

PSOPPC Services

Services Available to PSOs

- [Patient Safety Resources](#)
- [PSO Resource List](#)
- [PPC Contact](#)

Common Formats Support

- [Frequently Asked Questions](#)
- [Release Schedule](#)

Submitting Data

- [PSO Data Submission to the PSOPPC](#)

Spotlight

- [New PPC Contact, Hospital Common Formats, v1.1.1.2](#)
- [Hospital Common Formats, Version 1.2 Now Available \(04/23/2012\)](#)
- To provide feedback on the Common Formats, go to the [QD National Quality Forum \(NQF\)](#).

Hospital Common Formats



PSO Privacy
Protection Center

[Sign In](#) [How do I get an account?](#)

Support [Home](#) [Contact Us](#)

Search

[Home](#)

[About the PSOPPC](#)

[Data Submission](#)

AHRQ Common Formats

[Technical Assistance
Resources](#)

[Questions and Answers](#)

[News and Alerts](#)

Hospital Common Formats

Through a contract with the Agency for Healthcare Research and Quality (AHRQ), the National Quality Forum (NQF) solicited feedback on the formats from private sector organizations and individuals. The NQF, a nonprofit organization that focuses on healthcare quality, then convened an expert panel to review the comments received, and provide feedback to AHRQ. Based on the expert panel's feedback, AHRQ further revised and refined the Common Formats that are now available as Hospital Common Formats Version 1.2 & 1.1.

The following Hospital Common Formats are active for reporting and available for implementation and use by healthcare providers and Patient Safety Organizations (PSOs). These versions of the Common Formats are also accepted by the PSOPPC for national reporting.

Hospital Common Formats - Version 1.2

- [Event Descriptors, Sample Reports, & Forms](#)
- [Technical Specifications](#)
- [Users Guide](#)

Hospital Common Formats - Version 1.1

- [Event Descriptors, Sample Reports, & Forms](#)
- [Technical Specifications](#)
- [Users Guide](#)

ActionNet

[Terms Of Use](#) [Privacy Policy](#) [Account Info](#) [Patient Information Act \(HIPAA\)](#)

https://www.psoapp.org/web/patientinfo/versions/1.2_documents

Deficiency Reports



Access to Hospital Complaint Data

- CMS issued Survey and Certification memo on March 22, 2013 regarding access to hospital complaint data
- Includes acute care and CAH hospitals
 - Does not include the plan of correction but can request
 - Questions to bettercare@cms.hhs.com
- This is the CMS 2567 deficiency data and lists the tag numbers
- Updating quarterly
 - Available under downloads on the hospital website at www.cms.gov

Access to Hospital Complaint Data

- There is a list that includes the hospital's name and the different tag numbers that were found to be out of compliance
 - Many on restraints and seclusion, EMTALA, infection control, patient rights including consent, advance directives and grievances
- Two websites by private entities also publish the CMS nursing home survey data and hospitals
- The ProPublica website
- The Association for Health Care Journalist (AHCJ) websites

Access to Hospital Complaint Data

DEPARTMENT OF HEALTH & HUMAN SERVICES
Centers for Medicare & Medicaid Services
2000 Roanoke Boulevard, Mail Stop C3-21-16
Baltimore, Maryland 21244-1800



Center for Clinical Standards and Quality/Survey & Certification Group

Re: S&C: 13-21- ALL

DATE: March 21, 2013
TO: State Survey Agency Directors
FROM: Director
Survey and Certification Group
SUBJECT: Access to Statements of Deficiencies (CMS-2567) on the Web for Skilled Nursing Facilities, Nursing Facilities, Hospitals, & Critical Access Hospitals

Memorandum Summary

- **Survey Findings Posted on www.cms.gov:** In July 2012, the Centers for Medicare & Medicaid Services (CMS) began posting redacted Statements of Deficiencies (CMS-2567s) for skilled nursing facilities and nursing facilities on *Nursing Home Compare*. In March 2013, CMS began posting CMS-2567s for short-term acute care hospitals and critical access hospitals (CAHs) for surveys based on complaint investigations. This memorandum describes the contents and location of these files.
- **Other Web-based Tools Based on These Data:** At least two additional websites, provided by private parties (*ProPublica* and the Association for Health Care Journalists), publish information based on the CMS-2567 data. These websites are independent of CMS. CMS does not endorse or sponsor any particular private party application.
- **Plans of Correction (POC):** The posted CMS data do not contain any POC information. State Survey Agencies (SAs) and CMS Regional Offices (RO) may see an increase in requests for both the CMS-2567 and any associated POCs.
- **Questions & Answers:** We plan to issue an update to this memorandum that will include an attachment of frequently asked questions in order to provide answers to other queries that may arise.

Background – Nursing Home Survey Findings

In July 2012, CMS began posting nursing home statements of deficiencies, derived from the Form

Updated Deficiency Data Reports

The screenshot shows the CMS.gov website interface. At the top, there is a navigation bar with links for Home, About CMS, Newsroom Center, FAQs, Archive, Share, Help, Email, and Print. Below this is the CMS.gov logo and the text 'Centers for Medicare & Medicaid Services'. A search bar is also present. A horizontal menu contains categories: Medicare, Medicaid/CHIP, Medicare-Medicaid Coordination, Private Insurance, Innovation Center, Regulations and Guidance, Research, Statistics, Data and Systems, and Outreach and Education. The breadcrumb trail reads: Home > Medicare > Survey & Certification - Certification & Compliance > Hospitals. On the left, a sidebar lists various provider types, with 'Hospitals' selected. The main content area is titled 'Hospitals' and contains the following text:

This page provides basic information about being certified as a Medicare and/or Medicaid hospital provider and includes links to applicable laws, regulations, and compliance information.

A hospital is an institution primarily engaged in providing, by or under the supervision of physicians, inpatient diagnostic and therapeutic services or rehabilitation services. Critical access hospitals are certified under separate standards. Psychiatric hospitals are subject to additional regulations beyond basic hospital conditions of participation. The State Survey Agency evaluates and certifies each participating hospital as a whole for compliance with the Medicare requirements and certifies it as a single provider institution.

Under the Medicare provider-based rules it is possible for 'one' hospital to have multiple inpatient campuses and outpatient locations. It is not permissible to certify only part of a participating hospital. Psychiatric hospitals that participate in Medicare as a Distinct Part Psychiatric hospital are not required to participate in their entirety.

However, the following are not considered parts of the hospital and are not to be included in the evaluation of the hospital's compliance:

- Components appropriately certified as other kinds of providers or suppliers. i.e., a distinct part Skilled Nursing Facility and/or distinct part Nursing Facility, Home Health Agency, Rural Health Clinic, or Hospice; Excluded residential, custodial, and non-service units not meeting certain definitions in the Social Security Act; and,
- Physician offices located in space owned by the hospital but not functioning as hospital outpatient services departments

Accredited Hospitals - A hospital accredited by a CMS-approved accreditation program may substitute accreditation under that program for survey by the State Survey Agency. Surveyors assess the hospital's compliance with the Medicare Conditions of Participation (CoP) for all services, areas and locations covered by the hospital's provider agreement under its CMS Certification Number (CCN).

Although the survey generally occurs during daytime working hours (Monday through Friday), surveyors may conduct

Deficiency reports available at www.cms.gov/Medicare/Provider-Enrollment-and-Certification/CertificationandCompliance/Hospitals.html

CMS Hospital Nursing CoPs



TJC Revised Requirements

- TJC has published many changes over the past two years
 - Many of the changes reflected in their standards is to be in compliance with the CMS CoP
 - Standards are for hospitals that use them to get deemed status to allow payment for M/M patients
 - This means hospitals do not have to have a survey by CMS every 3 years
 - Can still get a complaint or validation survey
 - So now TJC standards crosswalk closer to the CMS CoPs (not called JCAHO any more)

Mandatory Compliance

- Hospitals that participate in Medicare or Medicaid must meet the COPs for all patients in the facilities and not just those patients who are Medicare or Medicaid
- Hospitals accredited by TJC, CIHQ, AOA, or DNV Healthcare have what is called deemed status
 - These are the only ones that CMS has given deemed status to for hospitals
- This means you can get reimbursed without going through a state agency survey
 - States can still institute a survey and be more restrictive

Survey Procedure

- Step one is publication in Federal Register
- Step two is where CMS publishes the interpretive guidelines
- The interpretive guidelines provide instructions to the surveyors on how to survey the CoPs
 - These are called survey procedure
 - Not all the standards have survey procedures
 - Questions such as “Ask patients to tell you if the hospital told them about their rights”

Nursing Services 385

- Standard: Must have an organized nursing service that provides 24 hour nursing services
 - Must have at least one RN furnishing or supervising 24 hours
 - You must have a RN on duty at all times
- Survey procedures determine nursing services is integrated into hospital QAPI
- Make sure there is adequate staffing

Survey Procedure 385

- Surveyor is suppose to interview the chief nursing officer (CNO) which CMS calls the DON or Director of Nursing
- Surveyor is to request a copy of the organizational chart
- Will look at job descriptions including the CNO
- Surveyor to select at least one patient from every inpatient nursing unit
 - Suppose to observe nursing care
 - To make sure there is adequate staffing
 - Will also look at patient medical records and care plans to make sure up to date, incident reports and P&P and talk to patients

Director of Nursing Service 0386

- Standard; Hospital must have a well organized service of administrative authority and delineations of responsibilities for patient care
- CNO must be a current licensed RN
- CNO is responsible for operating the nursing department
- CNO is responsible for determining types and numbers of nursing personnel and staff
 - Includes nurses, supervisors, assistant director, unit clerks, orderlies, nurse aides etc.

Nursing Service 386

- Must be one nursing service hospital wide
- Operation of nursing services includes the quality of care provided by nursing
- Survey Procedure
 - To verify CNO approves patient care P&P's
 - Verify CNO develops the nursing service staffing P&P
 - Will review the organization chart and look at lines of authority in the nursing department
 - Will read the job description for the CNO to make sure it specified duties and responsibilities of nursing services

Nurse Staffing 392

- Standard; Nursing service must have adequate number of nurses and personnel to care for patients
- Must have qualified nursing supervisory personnel
- Every department or unit must have a RN present
 - Not available if working on two units at same time
 - Must ensure RN for the bedside care of any patient
 - Must revise as needed for nursing staff absenteeism

Survey Procedure 392

- Surveyor to look at staffing schedules that correlate number and acuity of patients
- Surveyor to take into account the number of patient, intensity of illness and nursing needs
- Surveyor to take into consideration the training and experience of the personnel
- Also to consider the physical layout and size of the hospital
- Surveyor is to review the medical records of the patients to makes sure care is provided as ordered

Nurse Staffing 392

- There are 3 evidenced based studies that show the importance of having adequate staffing which results in better outcomes
- Study said patients who want to survive their new hospital visit should look for low nurse-patient ratio
- First evidenced based study on impact of nursing staff on patient outcomes
- Nurse Staffing and Quality of Patient Care, AHRQ, Evidence Report/Technology Report Number 151, March 2007, AHRQ Publication No. 07-E005¹

¹<http://www.ahrq.gov/downloads/pub/evidence/pdf/nursestaff/nursestaff.pdf>

AHRQ Nurse Staffing and Quality

Evidence Report/Technology Assessment
Number 151

Nurse Staffing and Quality of Patient Care

Prepared for:
Agency for Healthcare Research and Quality
U.S. Department of Health and Human Services
540 Gaither Road
Rockville, MD 20850
www.ahrq.gov

Contract No. 290-02-0009

Prepared by:
Minnesota Evidence-based Practice Center, Minneapolis, Minnesota

Investigators
Robert L. Kane, M.D.
Tatyana Shamliyan, M.D., M.S.
Christine Mueller, Ph.D., R.N.
Sue Duval, Ph.D.
Timothy J. Wilt, M.D., M.P.H.

www.ahrq.gov/downloads/pub/evidence/pdf/nursestaff/nursestaff.pdf
www.ahrq.gov/downloads/pub/evidence/pdf/nursestaff/nursestaff.pdf

Nursing Linked to Safety

- IOM (National Academy of Medicine) study also linked adequate staffing levels to patient outcomes
- Limit the number of hours worked to prevent fatigue
- Suggests no mandatory overtime for nurses
- Never work a nurse over 12 hours or 60 hours in one week
 - Or will have 38% or 3 times the error rate

Nursing Linked to Safety

- Also showed medication error rate, falls, pressure ulcers, UTI, surgery site infections, gastric ulcers, codes, readmission rate, LOS (length of stay), etc. are linked to staffing
- Redesigning the work force
- See Keeping Patients Safe: Transforming the Work Environment of Nurses 2004¹

¹www.nap.edu/openbook/0309090679/html/23/html

Nursing Staffing Linked to Safety

- AHRQ 2008 has published 3 volume, 51 chapter handbook for nurses at no cost
- Great resource that every hospital should have
- Nurse Staffing and Patient Care Quality and Safety
- Again shows that patient safety and quality is affected by short staffing
- Patient Safety and Quality: An Evidence-Based Handbook for Nurses, 2008¹
- ¹<http://www.ahrq.gov/qual/nursesfdbk>

You Are Here: [AHRQ Home](#) > [Quality & Patient Safety](#) > [Medical Errors & Patient Safety](#) > [Patient Safety and Quality: An Evidence-Based Handbook for Nurses](#)

Patient Safety and Quality

An Evidence-Based Handbook for Nurses

Nurses play a vital role in improving the safety and quality of patient care—not only in the hospital or ambulatory treatment facility, but also of community-based care and the care performed by family members. Nurses need know what proven techniques and interventions they can use to enhance patient outcomes.

To address this need, the Agency for Healthcare Research and Quality (AHRQ), with additional funding from the Robert Wood Johnson Foundation, has prepared this comprehensive, 1,400-page, handbook for nurses on patient safety and quality—*Patient Safety and Quality: An Evidence-Based Handbook for Nurses*. (AHRQ Publication No. 08-0043).

Experts in the field reviewed the literature, and their contributions are grouped into these sections:

[Patient Safety and Quality](#) / [Evidence-based Practice](#) / [Patient-centered Care](#) / [Working Conditions—Work Environment](#) / [Critical Opportunities for Patient Safety and Quality](#) / [Tools](#)

Select to download the [Entire Volume](#) as a PDF file (10 MB) or [individual chapters](#) (below). All PDF files are accessible. [PDF Help](#).

Select for [Ordering Information](#).

Edited by Ronda G. Hughes, Ph.D., M.H.S., R.N., of AHRQ

Contents

[Introduction](#)

www.ahrq.gov/qual/nursesfdbk

RN 24 Hours a Day A-0393

- **Standard:** Must have 24 hour nursing services provided or supervised by a RN
 - Exception for rural hospitals that have a nursing waiver granted for temporary shortage of nurses
 - Will make sure salary offered is comparable to three nearest hospitals
- Rural hospitals with 50 or fewer beds may be granted a temporary waiver of 24 hour RN requirement by the regional office
- Surveyor is to verify that there is at least one RN on each unit 24 hours a day

Verify Licensure 394

- **Standard:** Must have procedure to ensure nursing personnel have a valid and current license
 - Hospital procedure must ensure all nurses have a current and valid license
 - Must assure that all staff meet standards such as continuing education and certification and training
 - Surveyor will look at license verification P&P
 - Will look at HR records to make sure licensed

Verification of Nursing License

- Most state boards of nursing have an online verification process
 - Considered primary source verification
 - Can print out information for employee file
 - Don't forget to check the OIG list of excluded individuals (LEIE) and document it in the HR file for nurses

LEIE Downloadable Database

The screenshot shows the website for the Office of Inspector General, U.S. Department of Health & Human Services. The page is titled "LEIE Downloadable Databases". It features a navigation menu with links for "About OIG", "Reports & Publications", "Fraud", "Compliance", "Recovery Act Oversight", "Exclusions", and "Newsroom". A search bar is located in the top right corner. The main content area includes a "Download the LEIE Database" section with a "Download" button and a "Follow this page" notification. A "Search the OIG LEE Database" button is also present. On the right side, there is a "I'm looking for" section with a dropdown menu and a list of links: "Online Searchable Database", "LEIE Downloadable Databases", "Monthly Supplements Archive", "Quick Tips", and "Background Information".

Office of Inspector General
U.S. Department of Health & Human Services

Home / Exclusions / LEIE Downloadable Database

LEIE Downloadable Databases

Subscribe to receive email when this page is updated

Download the LEIE Database

Follow this page updated: 12-12-2012

LEIE Database

11-2012 Updated LEE Database EXE (2P)

Current Monthly Supplements

Search the OIG LEE Database

I'm looking for

Let's start by choosing a topic:

Select One

- Online Searchable Database
- LEIE Downloadable Databases
- Monthly Supplements Archive
- Quick Tips
- Background Information

[www.oig.hhs.gov/
exclusions/exclusions_list.asp](http://www.oig.hhs.gov/exclusions/exclusions_list.asp)

RN for Every Patient 395

- Standard; A RN must supervise and evaluate the nursing care for every patient
- RN must do admission assessment
- Must use acceptable standard of care
- Must follow hospital P&P
- Evaluation would include assessing each patient's needs, health status and response to interventions

Nursing Care Plan 396

- **Standard:** Hospital must ensure that nursing staff develop and keeps a current, nursing care plan for each patient
 - The nurse plan may be part of an interdisciplinary plan
- Frequent problematic standard
- Starts upon admission, includes discharge planning, physiological and psychosocial factors
- Assessment considers goals, physiological and psychosocial factors and discharge planning

Nursing Care Plan 396

- Based on assessing the patient's needs
- The interdisciplinary POC does not eliminate the need for a nursing plan of care
- You have to have one but it can be part of the larger, coordinated interdisciplinary POC
 - Interdisciplinary plan of care serve to promote collaboration between members of the team
- Care plan is part of the patient's medical records and must be initiated soon after admission, revised and implemented
 - Will look at **6-12** care plans

RN Assigns Care of Patient 397

- **Standard:** RN must assign the nursing care of each patient to other nursing staff in accordance with patient needs
 - Based on the qualifications and competence of the staff available
- RN must make all patient care assignments
- CNO will ensure all staff have proper education, experience, competence and qualifications for each patient
- Surveyor will review nursing assignments to make sure staff are qualified

Agency Nurses 398

- **Standard:** Agency nurses must adhere to P&P's
 - CMS calls them non-employee nurses
- CNO must provide adequate supervision and evaluate activities of agency nurses (once a year)
 - Includes other personnel such as volunteers
 - Must be supervised by RN who is a hospital employee
- CNO must make sure agency nurses know the hospital P&Ps
- Must include orientation to hospital and to specific unit, emergency procedures, and safety P&P's

Tag 405 Revised in Pharmacy Memo 2016

DEPARTMENT OF HEALTH & HUMAN SERVICES
Centers for Medicare & Medicaid Services
7500 Security Boulevard, Mail Stop C2-21-16
Baltimore, Maryland 21244-1850



Center for Clinical Standards and Quality/Survey & Certification Group

Ref: S&C: 16-01-Hospital

DATE: October 30, 2015

TO: State Survey Agency Directors

FROM: Director
Survey and Certification Group

SUBJECT: Revised Hospital Guidance for Pharmaceutical Services and Expanded Guidance Related to Compounding of Medications

Memorandum Summary

Hospital Appendix A Updated: The Centers for Medicare & Medicaid Services (CMS) has updated the State Operations Manual (SOM) Appendix A with respect to both the hospital survey process and the interpretive guidelines for the pharmaceutical services Condition of Participation (CoP). The update includes the following features:

- ***Pharmaceutical Services:*** Revisions were made to portions of the pharmaceutical services CoP to bring them into alignment with current accepted standards of practice. To improve clarity, the revised guidance addresses: accepted professional pharmacy principles, including United States Pharmacopeia (USP) standards; compounding of medications, particularly compounded sterile preparations (CSPs); determining

Preparation/Admin of Drugs 405 2016

- **Standard:** Drugs must be prepared and administered according to state and federal law (Effective Nov. 20, 2015)
- **Standard:** Need an practitioner's **order**
 - Important issue with CMS to have an order for all medications administered or standing order
 - Make sure order is documented in the medical record
 - Surveyor will observe nurse prepare and pass medications

Drugs & Biologicals 405

- Drugs and biologicals may be administered on orders of other practitioners:
 - Allowed by state law
 - State scope of practice act
 - Hospital P&P and
 - MS bylaws and R/R (Rules and Regulations)
- Must not only be within acceptable standards of practice (SOP) but done under the supervision of nursing
- CMS has blue box advisories which are not to be cited

Pharmacy Should Prepare Piggybacks & IVs

For Information – Not Required/Not to be Cited

Although the regulation addresses both preparation and administration of drugs and biologicals and does not prohibit preparation of drugs by nursing staff, to improve patient safety it is generally preferable for hospitals to avoid nurse preparation of drugs in patient care areas, and instead rely upon pharmacy IV admixture systems and/or commercially available unit dose products.

Preparation/Administration of Drugs 405

- **Standard:** Medications must be prepared and administered with acceptable national standards of practice and mentions specific organizations
 - National Coordinating Council for Medication Error Reporting and Prevention
 - Institute for Healthcare Improvement
 - U.S Pharmacopeia
 - Institute for Safe Medication Practices
 - Infusion Nurses Society
 - CDC at www.cdc.gov
 - Also according to the TJC MM chapter, manufacturer's directions and hospital policy

Timing of Medication Administration Tag 405

- What are acceptable standards of care?
 - National organizations that are recognized in the field issue written statements and policies that direct patient care
- The hospital's P&Ps must be consistent with SOC
- Standards of care can be set by state pharmacy boards and national organizations like the ones mentioned by CMS
- Others include:
 - ASHP (American Society of Healthcare System Pharmacist), American Nurses Association (ANA), American Pharmacy Association (APA), APIC, etc.

ISMP Institute for Safe Medication Practices



Institute for Safe Medication Practices

A Nonprofit Organization Educating the Healthcare Community and Consumers About Safe Medication Practices



[Home](#) [Support ISMP](#) [Newsletters](#) [Webinars](#) [Report Errors](#) [Educational](#) [Store](#) [Consulting](#) [FAQ](#) [Tools](#) [About Us](#) [Contact Us](#)



www.ismp.org

Google™ Custom Search

Search

ISMP ANNUAL FUND

Looking forward to next 20 years of advancing medication safety

[FIND OUT MORE](#)

2014 Medication Safety Intensive

October 2 and 3, 2014
Nashville, TN

December 5 and 6, 2014
Anaheim, CA

[Click here for details](#)

UPCOMING WEBINARS

The Pharmacist's Role in Protocol-Driven Care for Pain Management, Nutritional Support, Anticoagulation, and More!

Wednesday, July 16, 2014
from 1:30pm - 3:00pm ET

[Click here for more information](#)

Education & Awareness

- [Newsletters](#)
- [Consulting Services](#)
- [Educational Programs](#)
- [Let ISMP be your PSO](#)
- [Professional Development](#)
- [Self Assessments](#)
- [Consumers](#)

Medication Safety Tools & Resources

Featured Tools

- [New standards for healthcare connectors – the "Stay Connected" program](#)
- [The Root Cause Analysis Workbook for Community/ Ambulatory Pharmacy](#)
- [Special Error Alerts](#)
- [2014-15 Targeted Medication Safety Best Practices for Hospitals](#)
- [ISMP Guidelines](#)
- [High-Alert Medications](#)
- [Confused Drug Name List](#)
- [Community Pharmacy Medication Safety Tools and Resources](#)
- [Error-Prone Abbreviation List](#)

Report Medication Errors or Safety Concerns

QuarterWatch
Perspectives on Hypersensitivity
[READ NEW ISSUE](#)

Research-based Medication Safety Tools **FREE**
 Consumer medication leaflets
 Risk-reduction scorecards software
 Bar-coding readiness assessment

Long-Term Care AdviseERR
NEW medication safety newsletter for LTC facilities.
[SUBSCRIBE](#)

Infusion Nurses Society INS



[Home](#) [Member Login](#) [Not a Member? Join Our Mailing List](#) [Renew/Rejoin](#) [Site Map](#)



[Online Store](#) | [What's New](#) | [Infusion-Related Links](#) | [Position Papers](#) | [Career Center](#) | [Certification \(INCC\)](#)

[Knowledge Center](#) [About INS](#) [Membership](#) [Chapters/Affiliates](#) [Meetings/Events](#) [Exhibiting and Advertising](#) [Publications](#) [Gardner Foundation](#)

Signed Up for the CRNI® Exam?

Challenging infusion nurses since 1984



The two most recommended study aids by CRNI® passers

Get the Study Aid Bundle Today

Funded through an educational grant by



"I have gained so much from belonging to INS. Infusion therapy touches nearly every patient, and I feel I make a real difference for them because of the knowledge I have obtained through INS." --- Linda Resler, BSN, RN, CRNI®

For more than 40 years, INS has set the standard for infusion care, providing education to nurses and health care professionals who practice infusion therapy. Through membership, meetings, resources, and scholarships, the opportunities for lifelong learning never stop.

[Member Login](#)

[Join INS Now!](#)

[Learn More](#)

Upcoming Meeting

Register today for the Fall National Academy!

www.ins1.org



National Coordinating Council

Home
About NCC MERP
About Medication Errors
Council Recommendations
Report a Medication Error
NAN Alert
Council Communiqué
Members Page
For Consumers
Contact
Sitemap


Upcoming Meetings:

WebEx, July 17, 2014 1PM-4PM

Rockville, October 22, 2014 10AM-4PM

In-person meetings are held at USP headquarters in Rockville, MD

[NCC MERP 15 Year Anniversary Report](#)



► **Welcome to the NCC MERP web site**
National Coordinating Council for Medication Error

National Coordinating Council for Medication Error Reporting and Prevention

The National Coordinating Council for Medication Error Reporting and Prevention (NCC MERP) is an independent body comprised of [27 national organizations](#).

In 1995, USP spearheaded the formation of the National Coordinating Council for Medication Error Reporting and Prevention: [ABOUT NCC MERP](#)

Leading national health care organizations are meeting, collaborating, and cooperating to address the interdisciplinary causes of errors and to promote the safe use of medications. USP is a founding member and the Secretariat for NCC MERP. For a history on NCC MERP activity, see [Council Communiqué](#)

MEDICATION ERRORS:

Definition: NCC MERP defines a Medication Error

Category Index: Our Medication Error Index classifies an error according to the severity of the outcome, shown by chart ([Color / Black & White](#)) and algorithm ([Color / Black & White](#))

Dangerous Abbreviations: See table for intended meaning and common errors

Taxonomy: NCC MERP provides a standard taxonomy of medication errors to provide a standard language and structure when analyzing medication error reports.

Are you receiving the NAN Alert? The National Alert Network (NAN) publishes incident driven reports of medication errors; lessons learned can be used to increase the safety of the medication use system. Click on [NAN Alert](#) to subscribe and see previous editions!

Is your organization interested in membership? [Find out more.](#)

© 2014 National Coordinating Council for Medication Error Reporting and Prevention. All Rights Reserved. *Permission is hereby granted to reproduce information contained herein provided that such reproduction shall not modify the text and shall include the copyright notice appearing on the pages from which it was copied. This copyright statement will change to the

<http://www.nccmerp.org/pdf/algorColor2014-06-12.pdf>

www.nccmerp.org

Council Recommendations
Report a Medication Error
NAN Alert
Council Communiqué
Members Page
For Consumers
Contact
Sitemap

Upcoming Meetings:

WebEx, July 17, 2014 1PM-4PM

Rockville, October 22, 2014 10AM-4PM

In-person meetings are held at USP headquarters in Rockville, MD

[NCC MERP 15 Year Anniversary Report](#)

NATIONAL ALERT NETWORK (NAN)

 [Print](#)

The National Alert Network (NAN) is a coalition of members of the National Coordinating Council for Medication Error Reporting and Prevention (NCC MERP). The Institute for Safe Medication Practices (ISMP) and the American Society of Health-System Pharmacists (ASHP) publish the alerts from the National Medication Errors Reporting Program, operated by ISMP. The alerts are incident driven. The NCC MERP, ISMP and the ASHP encourage the sharing and reporting of medication errors, so that lessons learned can be used to increase the safety of the medication use system.

[February 18, 2014](#)

Potential inaccuracy of electronically transmitted medication history information used for **medication reconciliation**

[June 10, 2013](#)

Important Change with **Heparin Labels**

[April 17, 2013](#)

Confusion regarding the generic name of the HER2-targeted drug **KADCYLA (ado-trastuzumab emtansine)**

[January 23, 2013](#)

Severe burns and permanent scarring after **glacial acetic acid** ($\geq 99.5\%$) mistakenly applied topically

[April 25, 2012](#)

Proper disposal of **fentaNYL patches** is critical to prevent accidental exposure

[March 18, 2012](#)

Potential for wrong route errors with Exparel (**bupivacaine liposome injectable suspension**)

[Jun 2011](#)

Risk of potentially fatal overdose with **colistimethate**

[June 2010](#)

EPINEPHrine pre-filled syringe shortage

[Apr 2010](#)

Another child is victim of **heparin** error

Institute for Healthcare Improvement IHI



Improving Health and Health Care Worldwide

www.ihl.org



ABOUT US

TOPICS

EDUCATION

RESOURCES

REGIONS

ENGAGE WITH IHI



Effectively preventing *C. difficile* requires true multidisciplinary teamwork, says IHI faculty Dr. Brian Koll, and infection prevention staff are not solely responsible for this work »

FOCUS AREAS

- Improvement Capability »
- Person- and Family-Centered Care »
- Patient Safety »
- Quality, Cost, and Value »
- Triple Aim for Populations »

OPEN SCHOOL

The IHI Open School is transforming health care education around the world »



Free Audio Program

AUDIO PROGRAM



WIHI: From Prehospital to In-Hospital: The Continuum for Time-Sensitive Care July 24 | 2-3pm ET »

WEB-BASED TRAINING



Behavioral Health Integration: A Key Step Towards the Triple Aim Begins August 14 »

WEB-BASED TRAINING



Appropriate Use of Blood Products Begins August 19 | An IHI Expedition »

USP U.S. Pharmacopeial

The screenshot shows the homepage of the U.S. Pharmacopeial Convention website. At the top, there are language selection buttons for English, Español, 简体中文, and Português. A 'Log-In' section includes a dropdown menu for 'Select an Account' and a 'Go' button, along with a 'Cart' icon. The USP logo and 'U.S. Pharmacopeial Convention' text are prominently displayed. A search bar is located to the right of the logo, with a 'Search' button and a 'Go' button. Below the search bar are links for 'Calendar', 'Support', and 'A to Z Reference Standards Index'. A navigation menu contains links for 'About USP', 'USP-NF', 'Dietary Supplements', 'Food Ingredients', 'Reference Standards', 'Global', 'Meetings & Courses', 'News', and 'Store'. A large banner for the 'Call for 2015-2020 Candidates' is featured, with the text 'USP Council of Experts • Expert Committees' and a 'Make a Difference—Become a USP Volunteer' button. A 'CONVENTION 2015' logo is also present. Below the banner, the 'Standards Updates' section is active, showing updates to the USP-NF. A list of updates includes: 'Four New Intent to Revise Notices (27-Jun-2014)', 'Methyphenidate Hydrochloride Extended-Release Tablets Revision Bulletin Updated (27-Jun-2014)', 'Additional Feedback Sought on Proposed Storage and Distribution General Chapters (posted 13-Jun-2014)', and 'USP 37-NF 32, Second Supplement Commentary (02-Jun-2014)'. The 'Find information for...' section lists categories: 'Healthcare Professionals', 'Manufacturers', 'Delegates/Experts/Trustees', 'Patients/Consumers', and 'Regulators'. The 'Connect with USP' section includes social media icons for Facebook, Twitter, YouTube, LinkedIn, and RSS. The 'Featured Highlights' section features a 'Call for 2015 Resolutions' banner. The 'Press Releases' section mentions 'The National Alliance for Hispanic Health and the U.S. Pharmacopeial Convention partner to raise awareness about the safe use of vitamins and other...'. The 'Key Issues' section lists 'USP Medicare Model Guidelines' and 'Compounding'. A 'Careers at' section is partially visible at the bottom right.

English Español 简体中文 Português

Log-In: Select an Account Go Cart

USP U.S. Pharmacopeial Convention

Search Entire Site

Calendar Support A to Z Reference Standards Index

About USP USP-NF Dietary Supplements Food Ingredients Reference Standards Global Meetings & Courses News Store

Our Mission
USP's mission is to improve global health through public standards and related programs that help ensure the quality, safety, and benefit of medicines and foods.

Call for 2015-2020 Candidates
USP Council of Experts • Expert Committees
Make a Difference—Become a USP Volunteer APPLY NOW

CONVENTION 2015

Standards Updates

USP-NF Reference Standards Food Chemicals Codex

Review these updates to the USP-NF.

- Four New Intent to Revise Notices (27-Jun-2014)
- Methyphenidate Hydrochloride Extended-Release Tablets Revision Bulletin Updated (27-Jun-2014)
- Additional Feedback Sought on Proposed Storage and Distribution General Chapters (posted 13-Jun-2014)
- USP 37-NF 32, Second Supplement Commentary (02-Jun-2014)

Find information for...

- Healthcare Professionals
- Manufacturers
- Delegates/Experts/Trustees
- Patients/Consumers
- Regulators

Connect with USP

Featured Highlights

Press Releases

The National Alliance for Hispanic Health and the U.S. Pharmacopeial Convention partner to raise awareness about the safe use of vitamins and other...

Key Issues

USP Medicare Model Guidelines

Compounding

Careers at

www.usp.org

Centers for Disease Control & Prevention CDC



Centers for Disease Control and Prevention
CDC 24/7: Saving Lives, Protecting People™

SEARCH



CDC A-Z INDEX ▾

Diseases & Conditions ▾

Healthy Living ▾

Travelers' Health ▾

More CDC Topics ▾

National HIV Testing Day

is June 27 - Get Tested!



Recommend Tweet Share

What's New

Healthy Eating & Lifestyle Resource Center
Million Hearts®
millionhearts.hhs.gov

News

Million Hearts
Looking for low-sodium recipes? Checkout Million Heart's new resource center.

Feature

Caribbean Travel
Traveling to the Caribbean? Stay healthy and safe.

Feature



Carbon Monoxide (CO) Poisoning
Learn to protect yourself and your family as CO can be deadly.

TIPS FROM FORMER SMOKERS

News

Campaign Preview 2014
CDC's Tips From Former Smokers Campaign: New Stories to be released on July 7, 2014.

CDC IV Guidelines



- Every hospital should have the 2011 CDC Guidelines for the Prevention of Intravascular Catheter Related Infections
 - How to prep the skin for the peripheral IV
 - How to secure the needle
 - How long to change the dressing
 - How long do you change the IV tubing



www.cdc.gov/hicpac/pdf/guidelines/bsi-guidelines-2011.pdf

Guidelines for the Prevention of Intravascular Catheter-Related Infections, 2011

Naomi P. O'Grady, M.D.¹, Mary Alexander, R.N.², Lillian A. Burns, M.T., M.P.H., C.I.C.³, E Patchen Dellinger, M.D.⁴, Jeffery Garland, M.D., S.M.⁵, Stephen D. Heard, M.D.⁶, Pamela A. Lipsett, M.D.⁷, Henry Masur, M.D.¹, Leonard A. Mermel, D.O., Sc.M.⁸, Michele L. Pearson, M.D.⁹, Issam I. Raad, M.D.¹⁰, Adrienne Randolph, M.D., M.Sc.¹¹, Mark E. Rupp, M.D.¹², Sanjay Saint, M.D., M.P.H.¹³ and the Healthcare Infection Control Practices Advisory Committee (HICPAC)¹⁴.

*1*National Institutes of Health, Bethesda, Maryland

*2*Infusion Nurses Society, Norwood, Massachusetts

*3*Greenich Hospital, Greenwich, Connecticut

*4*University of Washington, Seattle, Washington

*5*Wheaton Franciscan Healthcare-St Joseph, Milwaukee, Wisconsin

*6*University of Massachusetts Medical School, Worcester, Massachusetts

*7*Johns Hopkins University School of Medicine, Baltimore, Maryland

*8*Warren Alpert Medical School of Brown University and Rhode Island Hospital, Providence, Rhode Island

*9*Office of Infectious Diseases, CDC, Atlanta, Georgia

*10*MD Anderson Cancer Center, Houston, Texas

*11*The Children's Hospital, Boston, Massachusetts

*12*University of Nebraska Medical Center, Omaha, Nebraska

*13*Ann Arbor VA Medical Center and University of Michigan, Ann Arbor, Michigan

NOTICE TO READERS:	7
Introduction	8
Summary Of Recommendations.....	9
Education, Training and Staffing	9
Selection of Catheters and Sites	10
Peripheral Catheters and Midline Catheters	10
Central Venous Catheters	11
Hand Hygiene and Aseptic Technique	12
Maximal Sterile Barrier Precautions	12
Skin Preparation	13
Catheter Site Dressing Regimens.....	13
Patient Cleansing	15
Catheter Securement Devices	15
Antimicrobial/Antiseptic Impregnated Catheters and Cuffs	15
Systemic Antibiotic Prophylaxis	15
Antibiotic/Antiseptic Ointments.....	15
Antibiotic Lock Prophylaxis, Antimicrobial Catheter Flush and Catheter Lock Prophylaxis.....	16
Anticoagulants	16
Replacement of Peripheral and Midline Catheters	16
Replacement of CVCs, Including PICCs and Hemodialysis Catheters	16
Umbilical Catheters.....	17
Peripheral Arterial Catheters and Pressure Monitoring Devices for Adult and Pediatric Patients	18
Replacement of Administration Sets	19
Needleless Intravascular Catheter Systems.....	19
Performance Improvement	20
Background Information	20
Terminology and Estimates of Risk	20
Epidemiology and Microbiology in Adult and Pediatric Patients	23
Pathogenesis	23
Strategies for Prevention of Catheter-Related Infections in Adult and Pediatric Patients.....	25
Education, Training and Staffing	25
Selection of Catheters and Sites	26
Peripheral and Midline Catheter Recommendations.....	26
Central Venous Catheters Recommendations.....	27

CDC 10 Recommendations

- The CDC has a page on Injection Safety that contains the excerpts from the Guideline for Isolation Precautions: Preventing Transmission of Infectious Agents in Healthcare Settings
- Summarizes their 10 recommendations for safe injection practices
- CMS expects hospitals to follow the CDC guidelines
 - Available at <http://www.cdc.gov/ncidod/dhqp/injectionSafetyPractices.html>

10 CDC Standards Safe Injection Practices

Injection Safety

Injection Safety

CDC's Role

CDC Statement

Information for Providers

Information for Patients

Preventing Unsafe Injection Practices

► Safe Injection Practices

CDC Clinical Reminder: Spinal Injection Procedures

Infection Prevention during Blood Glucose Monitoring and Insulin Administration

Recent Publications

Recent Meetings

The One & Only Campaign

Related Links

[One & Only Campaign](#)

[HICPAC](#)

[2007 Guideline for Isolation](#)

[Injection Safety](#) > [Preventing Unsafe Injection Practices](#)

[Recommend](#) 7

[Tweet](#)

[Share](#)

Safe Injection Practices to Prevent Transmission of Infections to Patients

Download the complete [2007 Guideline for Isolation Precautions: Preventing Transmission of Infectious Agents in Healthcare Settings](#) [PDF - 3.80 MB]

III.A.1.b. Safe Injection Practices The investigation of four large outbreaks of HBV and HCV among patients in ambulatory care facilities in the United States identified a need to define and reinforce safe injection practices 453. The four outbreaks occurred in a private medical practice, a pain clinic, an endoscopy clinic, and a hematology/oncology clinic. The primary breaches in infection control practice that contributed to these outbreaks were 1) reinsertion of used needles into a multiple-dose vial or solution container (e.g., saline bag) and 2) use of a single needle/syringe to administer intravenous medication to multiple patients. In one of these outbreaks, preparation of medications in the same workspace where used needle/syringes were dismantled also may have been a contributing factor. These and other outbreaks of viral hepatitis could have been prevented by adherence to basic principles of aseptic technique for the preparation and administration of parenteral medications 453, 454. These include the use of a sterile, single-use, disposable needle and syringe for each injection given and prevention of contamination of injection equipment and medication



[Email page link](#)

[Print page](#)

Contact Us:

[Centers for Disease Control and Prevention](#)
1600 Clifton Rd
Atlanta, GA 30333

[800-CDC-INFO](#)
(800-232-4636)
TTY: (888) 232-6348

[Contact CDC-INFO](#)

Medication Errors Tag 405

- CMS talks about the studies that show the large number of medication errors in hospitals
- Institute of Medicine said drug related adverse outcomes in 1.9 million inpatient hospital stays
- This is 4.7% of all patient stays
- There are 838,000 patient who are treated and released for drug related AE
- This is 0.8% of all visits
- Despite CPOE, ePHI, scanning and other technologies

Drugs & Biologicals 405

- CMS would allow them to document and sign the order
- For example, the above practitioners would be permitted as allowed by the state scope of practice such as by the state pharmacy board and if the hospital has granted them privileges
 - A PharmD manages the Anticoagulant Clinic or works with diabetic patients in managing their insulin
 - The MS approved the INR chart for patients on warfarin (coumadin)
 - Pharmacists changes dose and writes and signs off order

Drugs and Biologicals 405

- CMS calls them drugs and biologicals
 - Joint Commission calls them medications
 - Each state law differs on scope of practice on what PA, NP, CRNA, Pharm.D etc. can do so be aware of your state specific law
 - MS can C&P certain non-physician providers like the dietician or pharmacist
- Drugs and biologicals must be administered by or under the supervision of nursing or other personnel as allowed by law, P&Ps, and MS bylaws and R/Rs

Standing Orders and Outpatient Orders

- Drugs must be administered in response to an order from a practitioner or based on standing orders
- This includes ordering outpatient services for practitioners who are not privileged but are permitted by hospital & MS P&P to order
- Exception is for flu and Pneumovac
 - Need physician approved protocol after assessment of contraindications

CMS Changes to Medication Administration

- CMS issued a survey and certification memo with changes to Tag 405 on December 22, 2011, June 7, 2013 and March 14, 2014 memo effective June 6, 2014 and 2016
 - Tag 405 use to say that all medications must be given within 30 minutes of the scheduled time
 - Now three blocks of time to give medications
 - Thanks to the efforts of the ISMP
 - Included section on standing orders all but one sentence moved to tag 457

CMS Memo Med & Safe Opioid Use

DEPARTMENT OF HEALTH & HUMAN SERVICES
Centers for Medicare & Medicaid Services
7500 Security Boulevard, Mail Stop C2-21-16
Baltimore, Maryland 21244-1850



Center for Clinical Standards and Quality/Survey & Certification Group

Ref: S&C: 14-15-Hospital

DATE: March 14, 2014

TO: State Survey Agency Directors

FROM: Director
Survey and Certification Group

SUBJECT: Requirements for Hospital Medication Administration, Particularly Intravenous (IV) Medications and Post-Operative Care of Patients Receiving IV Opioids

Memorandum Summary

- **Medication Administration:** We are updating our guidance for the hospital medication administration requirements to:
 - Make clear that the medication administration requirements under the nursing services condition of participation (CoP) are related to only some components of the overall hospital medication process, but that hospitals are expected, through this and the related requirements under the pharmaceutical services and quality assessment/performance improvement CoPs, to take a comprehensive approach to the medication process.
 - Update our guidance for IV medications and blood transfusions in general; and
 - Reflect the need for patient risk assessment and appropriate monitoring during and after medication administration, particularly for post-operative patients receiving IV opioid medications, in order to prevent adverse events.
- **Immediate Post-operative Care:** Clarification is also being made to the guidance for the surgical services CoP requirement for hospitals to have adequate provisions for immediate post-operative care, to emphasize the need for post-operative monitoring of patients

Final Transmittal Issued June 6, 2014

CMS Manual System

Pub. 100-07 State Operations
Provider Certification

Transmittal 116

Department of Health &
Human Services (DHHS)
Centers for Medicare &
Medicaid Services (CMS)

Date: June 6, 2014

SUBJECT: Revised State Operations Manual (SOM), Appendix A, Survey Protocol, Regulations and Interpretive Guidelines for Hospitals

I. SUMMARY OF CHANGES: Clarification is provided for various provisions of 42 CFR 482.23(c), concerning medication administration, and 42 CFR 482.51(b)(4), concerning post-operative patient care.

NEW/REVISED MATERIAL - EFFECTIVE DATE: June 6, 2014
IMPLEMENTATION DATE: June 6, 2014

The revision date and transmittal number apply to the red italicized material only. Any other material was previously published and remains unchanged. However, if this revision contains a table of contents, you will receive the new/revised information only, and not the entire table of contents.

II. CHANGES IN MANUAL INSTRUCTIONS: (N/A if manual not updated.)
(R = REVISED, N = NEW, D = DELETED) – (Only One Per Row.)

R/N/D	CHAPTER/SECTION/SUBSECTION/TITLE
R	Appendix A/Survey Protocol, Regulations and Interpretive Guidelines for Hospitals/A-0405/§482.23(c)Standard: Preparation and Administration of Drugs
R	Appendix A/Survey Protocol, Regulations and Interpretive Guidelines for Hospitals/A-0409/§482.23(c)(4)/Blood transfusions and intravenous medications must be administered in accordance with State law and approved medical staff policies and procedures.
R	Appendix A/Survey Protocol, Regulations and Interpretive Guidelines for Hospitals/A-0412/§482.23(c)(6)/The hospital may allow a patient (or his or her caregiver/support person where appropriate) to self-administer both hospital-issued medications and the patient's own medications brought into the hospital, as defined and specified in the hospital's policies and procedures.
R	Appendix A/Survey Protocol, Regulations and Interpretive Guidelines for Hospitals/A-0957/§482.51(b)(4)/There must be adequate provisions for immediate post-operative care.

www.cms.gov/Regulations-and-Guidance/Guidance/Transmittals/Downloads/R116SOMA.pdf

ISMP New Guideline www.ismp.org



ISMP Acute Care Guidelines for Timely Administration of Scheduled Medications

Background

The Institute for Safe Medication Practices (ISMP) developed these Acute Care Guidelines for Timely Administration of Scheduled Medications after conducting an extensive survey in late-2010 involving almost 10,000 nurses regarding the requirement in the Centers for Medicare & Medicaid Services (CMS) conditions of participation (COP) to administer medications within 30 minutes before or after the scheduled time. The nurses who responded to the survey made it clear that changes to drug delivery methods and gradual increases in the complexity of care, number of prescribed medications per patient, and number of patients assigned to each nurse have made the long-standing CMS “30-minute rule” error prone.

Many nurses reported feeling great pressure to take shortcuts to comply with the rule, which have led to errors, some harmful. While delays in administering certain time-sensitive medications can also result in harm, some size-fit-all, inflexible requirements to administer all scheduled medications within 30 minutes of the scheduled time is a precarious substitute given that relatively few medications truly require exact timing of doses.

CMS staff have requested a copy of the final guidelines, and based on our conversations with them, we are optimistic that positive changes will be made to the current “30-minute rule.” For now, hospitals will still be held accountable for the “30-minute rule” in the CMS Interpretive Guidelines. However, given widespread support for these more reasonable and clinically appropriate guidelines, we hope CMS reviewers will allow hospitals to justify their carefully considered policies and procedures regarding timely medication administration using these guidelines to anchor the process.

Definitions

1 Scheduled medications include all maintenance doses administered

How to Use the Guidelines

These guidelines are applicable **ONLY** to scheduled medications (see definition below).

The guidelines are intended to be used as a resource when acute care organizations develop or revise policies and procedures related to timely administration of scheduled medications. The guidelines are not standards or evidence-based practices that have been proven by scientific studies, but they have been vetted by hundreds of medication and patient safety experts, hospital medication safety teams, professional nursing, pharmacy, and respiratory therapy organizations, the Joint Commission, hospital pharmacists, and frontline nurses who bear ultimate responsibility for administering medications in a timely manner.

An interdisciplinary team with adequate nursing representation needs to translate the guidelines into facility-specific policies and procedures. In general, the guidelines represent a safe, effective, and efficient approach to timely administration of scheduled medications. However, the details may differ from one organization to another based on differing patient populations and medication systems, including available technology.

Please keep in mind that the policies and procedures developed by acute care organizations using these guidelines will require flexibility of the goals for timely administration, or appropriate, to accommodate the additional time needed to learn to operate new medication-related technologies.

Advisory Group

A list of advisory group professionals who provided input during development of these guidelines can be found at: www.ismp.org/tools/guidelines/acutecareadvisory_group.pdf

2 Time-critical scheduled medications are those where death or

Practitioner Order Requirements

- Name of the patient
- Age and weight of the patients to facilitate dose calculation requirements
 - Must have P&P to address for children and use only Kg or Grams for newborns
 - Other circumstances like as weight on elderly patient with history of renal failure and is being prescribed antibiotics
 - Hospitals must specify a unified approach
- Date and time of the order

Use Kg and Not Pounds for Children

Acetaminophen Dosing Chart

Acetaminophen (Tylenol) Dose every 4 to 6 hours <i>Maximum 5 doses in 24 hours</i>		Infants' Concentrated Drops 80 mg/ 0.8 mL Dropperful (Use only the dropper provided)	Children's Suspension Liquid 160 mg/ 5 mL Teaspoon (tsp)	Children's Soft Chews Chewable 80 mg each Tablet	Junior Strength Chewable 160 mg each Tablet	Adult Regular Strength 325 mg each Tablet
Weight	Age					
6-11 lbs	0-3 mos	½ = 0.4 mL				
12-17 lbs	4-11 mos	1 = 0.8 mL	½ tsp			
18-23 lbs	12-23 mos	1 ½ = 0.8 + 0.4 mL	¾ tsp			
24-35 lbs	2-3 yrs	2 = 0.8 + 0.8 mL	1 tsp	2	1	
36-47 lbs	4-5 yrs		1 ½ tsp	3	1 ½	
48-59 lbs	6-8 yrs		2 tsp	4	2	
60-71 lbs	9-10 yrs		2 ½ tsp	5	2 ½	1
72-95 lbs	11 yrs		3 tsp	6	3	1 ½
96 lbs +	12 yrs +		4 tsp	8	4	2

Ibuprofen Dosing Chart

Ibuprofen (Motrin, Advil) Dose every 6 to 8 hours <i>Maximum 4 doses in 24 hours</i>		Infants' Concentrated Drops 50 mg/ 1.25 mL Dropperful (Use only the dropper provided)	Children's Suspension Liquid 100 mg/ 5 mL Teaspoon (tsp)	Children's Chews Chewable 50 mg each Tablet	Junior Strength 100 mg each Tablet	Adult Regular Strength 200 mg each Tablet
<i>Under 6 mos</i>		<i>Consult Your Child's Provider</i>				
Weight	Age					
12-17 lbs	6-11 mos	1 = 1.25 mL				
18-23 lbs	12-23 mos	1 ½ = 1.875 mL				
24-35 lbs	2-3 yrs		1 tsp	2		
36-47 lbs	4-5 yrs		1 ½ tsp	3		
48-59 lbs	6-8 yrs		2 tsp	4	2	
60-71 lbs	9-10 yrs		2 ½ tsp	5	2 ½	
72-95 lbs	11 yrs		3 tsp	6	3	
96 lbs +	12 yrs +		4 tsp	8	4	2

Practitioner Order Requirements

- Drug name
- Dose, frequency, and route
- Dose calculation requirements
- Exact strength or concentration, when applicable
- Quantity and/or duration, when applicable
- Specific instructions for use, when applicable and
- Name of the prescriber

Medical Staff Approved P&P

- MS must approve the P&P for medication administration
 - Should be part of QAPI process
 - Should be done in consultation with nurses and pharmacists
 - Drugs must be administered under supervision of nursing or other personnel
- CMS has many specifics which must be included in this MS approved P&P
- Needs to be consistent with state law and the scope of practice

P&P Requirements

- Must identify the categories of licensed personnel who can prepare and administer
 - For example, Ohio allows RNs and LPNs who have passed a pharmacy course to prepare and administer
- Must include the types of medications they are allowed to prepare and administration
 - For example, the Ohio Board of Nursing does not allow a LPN to hang blood or give certain IV medications
- Must address education or training requirements and CMS has some recommendations

Education Recommendation

- CMS recommend training in orientation and as part of continuing education
- Training **may** include the following;
 - Safe handling and preparation of authorized medications
 - Knowledge of the indications, side effects, drug interactions, compatibility, and dose limits of administered medications
 - Equipment, devices, special procedures, and/or techniques required for medication administration (IV pumps, PCA, tubing, etc.)

P&P Requirements

- What must be included in the training during orientation or CNE to demonstrate competence
- Training content and documentation of **competence**
- P&P must include basic safe practices for medication administration such as the following required elements
 - Patient's identity
 - To make sure it is the right patient and identifiers might include name, MR number, identification number, DOB
 - Confirmed by wrist band, patient identification card, patient statement or other things included in the hospital policy

P&P Requirements

- There must be agreement between the patient's MAR (medication administration record) and the medication's label
- Need to have culture of safety in which staff feel comfortable to ask questions
- Confirm before medication is given the following on the five rights:
 - Right medication, right patient, right dose
 - Right route (IM, PO, IV, IO, intrathecally, etc)
 - Right time to adhere to the prescribed frequency and time of administration

Medication Process 405

- Medication process has five stages
 - Ordering/prescribing
 - Transcribing and verifying
 - Dispensing and delivering
 - Administering
 - And monitoring/reporting
- CMS also mentions the recent literature regarding the nine rights of medication administration

9 Rights of Medication Administration

For Information – Not Required/Not to be Cited

Recent literature identifies up to nine “rights” of medication administration including:*

- Right patient*
- Right drug*
- Right route*
- Right time*
- Right dose*
- Right documentation*
- Right action (appropriate reason)*
- Right form*
- Right response*

However, other sources refer to 8 or 10 “rights, and some of these topics, such as right action, appear to involve prescribing and/or dispensing. Accordingly, there does not (yet) appear to be consensus about expanding beyond the 5 “rights.”

**Reference: Elliott, M. and Lis, Y. (2010). The Nine Rights of Medication Administration: An Overview. British Journal of Nursing, Vol. 19, 5, 300-305.*

Safe Injection Practices

- Must ensure staff follow SOP to prevent HAI related to medication preparation
 - References infection control worksheet
 - Assessed under infection control section
- Compounded sterile preparations (CSP) can cause HAI if proper precautions are not followed such as following the USP standards
- Nurses may prepare sterile medication for immediate use
- CMS mentions the following apply

Compounding 2016

- Must only involve simple transfer of not more than 3 commercially manufactured, sterile, nonhazardous products from the manufacturer's original container
- And not more than two entries into any one container including a vial or an IV bag
- Administration must be within one hour following the preparation
- Must follow aseptic technique during all phases of preparation

Compounding

- Must label it unless you prepare it and immediately administer it to the patient
- CSP label must include: patient identification, name and amount of ingredients, name or initial of person who prepared it, and exact one hour BUD
- Drug is outdated after its expiration date or BUD
 - BUD is December 2017 but multi-dose vial expires in 28 days when opened unless sooner by manufacturer
- Need P&P to give clear directions to staff on how to determine BUD date if not available from manufacturer

procedures must be based on accepted professional principles which are equivalent to, or more stringent than, those described in the USP/NF (USP).³

According to Chapters <795> and <797> of the USP, the BUD must be safe for patients, and determined conservatively. The section in USP <797> entitled “Determining Beyond-Use Dates,” which addresses sterile compounding, notes that “the truly valid evidence for predicting beyond-use dating can be obtained only through product-specific experimental studies.” It provides an example of testing considered more appropriate for certain types of CSPs such as “CSPs with a narrow therapeutic index, where close monitoring or dose titration is required to ensure therapeutic effectiveness and to avoid toxicity....” It also provides examples of important issues that a pharmacist must be able to critically interpret and evaluate when consulting literature sources in the process of determining a BUD; and distinguishes between reviewing literature specific to a particular drug, composition, concentration of ingredients, fill volume, container, storage conditions and duration of use, etc., versus merely reviewing available publications or tables. The former is the preferred approach, while the latter results in a “theoretical BUD,” which has an inherent likelihood of inaccuracy or error.

Timing of Medication 405

- P&P needs to include the timing of medication based on the nature of the medication and the clinical application to include:
 - Medications or categories of medications **not eligible** for scheduled dosing times
 - These are ones that require exact time based on diagnosis type, treatment requirements or therapeutic goals
 - Include definition in your P&P
 - Also looks at patient risk factors
 - Such as stat drugs, loading dose, one time dose for scheduled procedure, doses timed for serum drug level, PRN, or investigational drugs

3 Time Frames for Administering Medication

Time Critical Medicine

1 hour before or after

2 hours before or after

Timing of Medication P&P

- Medications that are **eligible** for scheduled dosing times
 - These are those prescribed on a repeated cycle of frequency, such as once a day, BID (twice a day), TID (three times a day), hourly intervals (every 1, 2, 3 or more hours), etc.
 - Goal is to achieve a therapeutic blood level
 - BID meds might be given at 9am/9 pm or 8am/8pm
 - Policy has the standardized times so pharmacy knows when to send to unit and nurse can assess VS if needed (such as pulse rate if dig) or review blood work (like a serum K level, INR, or dig level)

Timing of Medication P&P

- Medications that are **eligible** for scheduled dosing times (continued)
 - P&P on first dose of medication, using judgment regarding next dose, retiming of missed or omitted doses
- Medications that can be given outside of their scheduled dosing time
- Evaluation of the medication timing policy and including adherence rate
 - Must track medication errors related to timing of medications and include in the PI process

Timing of Medication P&P

- Time-critical scheduled medications (30 minute or 1 hour total window)
 - These are ones in which an early or late administration of greater than thirty minutes might cause harm or have significant, negative impact on the intended therapeutic or pharmacological effect
 - P&P must include whether these drugs are always time critical
 - Examples include: Antibiotics, Anticoagulants, Insulin, Anticonvulsants, Immunosuppressive agents, **Non-IV** Pain medication, medication more frequently than every 4 hours, and administered within a specified period of time in the order

Timing of Medication P&P

- Non-time-critical scheduled medications
 - These are medications for which a longer or shorter interval of time since the prior dose does not significantly change the medication's therapeutic effect or otherwise cause harm
 - Greater flexibility is given
 - Medications given once daily, weekly, or monthly
 - May be given within **2 hours** before or after but can not exceed a total window of 4 hours (such as Allegra once a day)
 - Med scheduled more frequently than daily but less than every 4 hours (such as bid or tid) can be given **1 hour** before or after for window not to exceed 2 hours

Timing of Medication P&P

- Missed or late administration of medications
 - Policy must include what action to take if missed or not given in permitted window of time
 - Missed dose may be due from patient who is out of the department, patient refusal, problems related to medication being available or other reasons
 - Policy needs to include parameters of when nursing staff are allowed to use their own judgment on the rescheduling of late or missed doses
 - Missed or late doses must be reported to the attending physician

Assessment & Monitoring of Patients 2014

- Patients on medications needed to be carefully monitored
 - May need clinical and lab data to evaluate medication
 - Monitor respiratory status, pulse ox, BP, end tidal CO₂ with patients on **opioids**
 - Evaluate clinical signs such as confusion, agitation, unsteady gait, itching etc.
 - Know **high risk medications** policy and safe practices
 - Know risk factors for ADE such as patient has liver or kidney failure, history of sleep apnea, obesity, smoking, drug-drug interaction and first time medication use

Assessment/Monitoring of Patients Receiving Medications

Observing the effects medications have on the patient is part of the multi-faceted medication administration process. Patients must be carefully monitored to determine whether the medication results in the therapeutically intended benefit, and to allow for early identification of adverse effects and timely initiation of appropriate corrective action. Depending on the medication and route/delivery mode, monitoring may need to include assessment of:

- Clinical and laboratory data to evaluate the efficacy of medication therapy, to anticipate or evaluate toxicity and adverse effects. For some medications, including opioids, this may include clinical data such as respiratory status, blood pressure, and oxygenation and carbon dioxide levels;*
- Physical signs and clinical symptoms relevant to the patient's medication therapy, including but not limited to, somnolence, confusion, agitation, unsteady gait, pruritus, etc.*

Certain types of medications are considered inherently high risk for adverse drug events. Although mistakes may or may not be more common with these drugs, the consequences of errors are often harmful, sometimes fatal, to patients. (See also the discussion of high-risk medications (typically referred to as “high-alert” medications) in the guidance for §482.25(b))

For Information – Not Required/Not to be Cited

The Institute for Safe Medication Practices (ISMP) makes available a list of high alert medications, which it defines as those medications that bear a heightened risk of causing significant patient harm when they are used in error. The current list may be found at: <http://www.ismp.org/Tools/highAlertMedicationLists.asp>

ISMP List of High Alert Medication

ISMP's List of *High-Alert Medications*

 [Printer friendly version](#)

High-alert medications are drugs that bear a heightened risk of causing significant patient harm when they are used in error. Although mistakes may or may not be more common with these drugs, the consequences of an error are clearly more devastating to patients. We hope you will use this list to determine which medications require special safeguards to reduce the risk of errors. This may include strategies like improving access to information about these drugs; limiting access to high-alert medications; using auxiliary labels and automated alerts; standardizing the ordering, storage, preparation, and administration of these products; and employing redundancies such as automated or independent doublechecks when necessary. (Note: manual independent double-checks are not always the optimal error-reduction strategy and may not be practical for all of the medications on the list).

Classes/Categories of Medications	Specific Medications
adrenergic agonists, IV (e.g., EPINEPH rine, phenylephrine, norepinephrine)	epoprostenol (Flolan), IV
adrenergic antagonists, IV (e.g., propranolol, metoprolol, labetalol)	magnesium sulfate injection
anesthetic agents, general, inhaled and IV (e.g., propofol, ketamine)	methotrexate, oral, non-oncologic use
antiarrhythmics, IV (e.g., lidocaine, amiodarone)	opium tincture
antithrombotic agents, including: <ul style="list-style-type: none"> ■ anticoagulants (e.g., warfarin, low-molecular-weight heparin, IV unfractionated heparin) ■ Factor Xa inhibitors (e.g., fondaparinux) ■ direct thrombin inhibitors (e.g., argatroban, bivalirudin, dabigatran etexilate, lepirudin) 	oxytocin, IV
	nitroprusside sodium for injection
	potassium chloride for injection concentrate
	potassium phosphates injection
	promethazine, IV
	vasopressin, IV or intraosseous

High Alert How to Guide IHI

10/01/2008



Getting Started Kit: Prevent Harm from High-Alert Medications

How-to Guide

A national initiative led by IHI, the 5 Million Lives Campaign aims to dramatically improve the quality of American health care by protecting patients from five million incidents of medical harm between December 2006 and December 2008. The How-to

www.ihl.org/NR/rdonlyres/8B2475CD-56C7-4D9B-B359-801F3CC3A8D5/0/HighAlertMedicationsHowToGuide.doc

So What's In Your Policy?



**WISCONSIN
PATIENT SAFETY
INSTITUTE**

330 E Lakeside St
PO Box 1551
Madison, WI 53701
Voice: 608.442.3789
800.762.8976
Fax: 608.283.5402

www.WPSI.org

MODEL HIGH-ALERT MEDICATIONS POLICY & PROCEDURES

PURPOSE

- To provide guidance to acute care organizations for the safe handling and administration of medications designated as High Alert Medications.
- To increase awareness of High Alert Medications, thereby improving patient safety.

DEFINITION

High Alert Medications are drugs that bear a higher risk of causing significant patient harm when they are used in error.¹

POLICY

- A. The following medications are appropriate for inclusion in a High Alert Medications policy.
- Epidural infusions
 - Fentanyl
 - Heparin (>100 units, flushes exempt)
 - Insulin (including regular, aspart, NPH, and glargine)
 - Lidocaine with epinephrine vials
 - Neuromuscular blocking agents (atracurium, cisatracurium, mivacurium, pancuronium, rapacuronium, rocuronium, succinylcholine, vecuronium, etc)
 - Patient Controlled Analgesia (PCA) infusions of any medication
 - Total Parenteral Nutrition (TPN) and Total Nutrient Admixture (TNA) solutions
 - Oncologic agents
 - Moderate sedation agents (e.g., midazolam)

- B. The following medications may also be appropriate for inclusion in a High Alert Medication policy in addition to the medications above.

- Glycoprotein IIb/IIIa inhibitors (eptifibatide, abciximab, tirofiban)
- Iron Dextran
- Adrenergic antagonists agents (e.g., esmolol)
- Anticonvulsants

- C. Concentrated electrolyte vials (e.g., potassium chloride) should not be stocked in patient care areas.

PROCEDURES

Safety procedures during the ordering, preparation, dispensing and administration of High Alert Medications include:

Prescribing

- A. Verbal orders for High Alert Medications should be discouraged.
- B. If possible, prescribing for High Alert Medications should be standardized using preprinted orders.

Preparation and dispensing

- A. All storage locations should be clearly labeled and separated from regular stock. If High Alert Medications must be kept in patient care areas, locked storage areas should be used with a distinct High Alert Medication warning label visibly placed on the storage bin.

POLICY

A. The following medications are appropriate for inclusion in a High Alert Medications policy.

- Epidural infusions
- Fentanyl
- Heparin (>100 units, flushes exempt)
- Insulin (including regular, aspart, NPH, and glargine)
- Lidocaine with epinephrine vials
- Neuromuscular blocking agents (atracurium, cisatracurium, mivacurium, pancuronium, rapacuronium, rocuronium, succinylcholine, vecuronium, etc)
- Patient Controlled Analgesia (PCA) infusions of any medication
- Total Parenteral Nutrition (TPN) and Total Nutrient Admixture (TNA) solutions
- Oncologic agents
- Moderate sedation agents (e.g., midazolam)
- Anesthetic agents (e.g., propofol)
- Adrenergic agonists (phenylephrine)

C. Concentrated electrolyte vials (e.g., potassium chloride) should not be stocked in patient care areas.

PROCEDURES

Safety procedures during the ordering, preparation, dispensing and administration of High Alert Medications include:

Prescribing

- A. Verbal orders for High Alert Medications should be discouraged.
- B. If possible, prescribing for High Alert Medications should be standardized using preprinted orders.

Preparation and dispensing

- A. All storage locations should be clearly labeled and separated from regular stock. If High Alert Medications must be kept in patient care areas, locked storage areas should be used with a distinct High Alert Medication warning label visibly placed on the storage bin.

The Wisconsin Patient Safety Institute enhances and promotes patient safety by advocating for the adoption of safe practices in health care organizations throughout Wisconsin.

Assessment & Monitoring of Patients

- ADE, such as anaphylaxis or opioid-induced respiratory depression may require timely and appropriate
- Post-medication monitoring in case of a high alert medication may include regular assessment of VS, pulse ox, and sedation levels of post surgery patient on PCA
- Such as Richmond agitation sedation scale (RASS) or the Pasero Opioid-Induced sedation scale (POSS), Inova Sedation Scale (ISS), Ramsey scale, Aldrete Scoring system

Pasero Opioid-induced Sedation Scale POSS

Pasero Opioid-induced Sedation Scale (POSS)

<https://secure.tha.com/surveys/files/pasero-opioid-induced-sedation-scale-poss.pdf>

S = Sleep, easy to arouse

Acceptable; no action necessary; may increase opioid dose if needed

1. Awake and alert

Acceptable; no action necessary; may increase opioid dose if needed

2. Slightly drowsy, easily aroused

Acceptable; no action necessary; may increase opioid dose if needed

3. Frequently drowsy, arousable, drifts off to sleep during conversation

Unacceptable; monitor respiratory status and sedation level closely until sedation level is stable at less than 3 and respiratory status is satisfactory; decrease opioid dose 25% to 50% or notify prescriber or anesthesiologist for orders; consider administering a non-sedating, opioid-sparing nonopioid, such as acetaminophen or an NSAID, if not contraindicated.

4. Somnolent, minimal or no response to verbal or physical stimulation

Unacceptable; stop opioid; consider administering naloxone; notify prescriber or anesthesiologist; monitor respiratory status and sedation level closely until sedation level is stable at less than 3 and respiratory status is satisfactory.

Richmond Agitation Sedation Scale RASS

Richmond Agitation Sedation Scale (RASS) *

www.icudelirium.org/docs/RASS.pdf

Score	Term	Description	
+4	Combative	Overtly combative, violent, immediate danger to staff	
+3	Very agitated	Pulls or removes tube(s) or catheter(s); aggressive	
+2	Agitated	Frequent non-purposeful movement, fights ventilator	
+1	Restless	Anxious but movements not aggressive vigorous	
0	Alert and calm		
-1	Drowsy	Not fully alert, but has sustained awakening (eye-opening/eye contact) to <i>voice</i> (≥ 10 seconds)	} Verbal Stimulation
-2	Light sedation	Briefly awakens with eye contact to <i>voice</i> (<10 seconds)	
-3	Moderate sedation	Movement or eye opening to <i>voice</i> (but no eye contact)	
-4	Deep sedation	No response to voice, but movement or eye opening to <i>physical</i> stimulation	} Physical Stimulation
-5	Unarousable	No response to <i>voice</i> or <i>physical</i> stimulation	

Procedure for RASS Assessment

1. Observe patient
 - a. Patient is alert, restless, or agitated. (score 0 to +4)
2. If not alert, state patient's name and *say* to open eyes and look at speaker.
 - b. Patient awakens with sustained eye opening and eye contact. (score -1)
 - c. Patient awakens with eye opening and eye contact, but not sustained. (score -2)
 - d. Patient has any movement in response to voice but no eye contact. (score -3)
3. When no response to verbal stimulation, physically stimulate patient by shaking shoulder and/or rubbing sternum.

Comparison of Sedation Scales Medscape

Pain Management Nursing

Comparison of Selected Sedation Scales for Reporting Opioid-Induced Sedation Assessment

Allison Theresa Nisbet, MSN, CPN, AOCNS, RN-BC, Florence Mooney-Cotter, MSN, CNS-BC, RN-BC | Disclosures
Pain Manag Nurs. 2009;10(3):154-164.

Comment



Print

- Abstract and Introduction
- Nurse Assessment of Sedation Using a Sedation Scale
- ▶ **Study Aims and Methods**
- Results
- Limitations
- Summary Recommendations
- References

Study Aims and Methods

The present research study was designed to report measures of reliability and validity of three sedation scales currently used to measure sedation as an outcome of opioid administration for pain management in non-critical care settings: the Inova Health System Acute Care Sedation Scale (ISS), the RASS, and the POSS. Reliability and validity had not been previously established for any of these scales in the non-critical care setting. The following research questions were addressed by the study:

Research question 1: Is there a significant difference in validity or reliability between three commonly used sedation scales when used by non-critical care nurses for the measurement of postopioid sedation?

Research question 2: Is there a significant difference in means observed between scales in the total correct score obtained by the nurses (sedation score and nursing actions chosen)?

Research question 3: Is there a significant difference in means observed between scales in the nurses' total combined rating of each scale's performance regarding ease of use, information provided to inform clinical decision making, and confidence (in

score obtained and actions chosen)?

The study aims had immediate organizational significance, because the scale (the ISS) used to assess opioid-induced sedation at the facility in which the research was conducted had not previously been tested for

ADVERTISEMENT

Once-daily SAMSCA provides a significant increase in serum sodium in as early as 8 hours

Experience free water clearance

SAMSCA is contraindicated in the following conditions:

- Urgent need to raise serum sodium acutely
- Inability of the patient to sense or appropriately respond to thirst
- Hypovolemic hyponatremia
- Concomitant use of strong CYP 3A inhibitors
- Anuric patients

FULL PRESCRIBING INFORMATION including Boxed WARNING

MOST POPULAR ARTICLES

According to PHYSICIANS

1. No Amount of Alcohol Is Safe
2. Alcohol and CVD: The Tipping Point
3. 10 Cool, Amazing Gadgets and Trends to Help Your Practice
4. More Evidence Musical Training May Boost Executive Function
5. Utility of a Single Early Warning Score in Patients With Sepsis in the Emergency Department

▶▶ View More

www.medscape.com/viewarticle/708387_3

Assessment & Monitoring of Patients

- Staff are expected to include patient reports of his experience with medication's effect
- Patient should be instructed to notify nurse if there is difficulty breathing or a reaction to the medication
- Hospital needs P&P to address the manner and frequency of monitoring
- P&P should include information to be communicated at shift change
- Should include patient's risk factors
- **Document** after medication administered

Surveyor Procedure Tag 405

- Surveyor to verify the established time requirements do not exceed the following:
 - 1 hour for time-critical scheduled medications
 - 2 hours for medications prescribed more frequently than daily, but no more frequently than every 4 hours and
 - 4 hours for medications prescribed for daily or longer administration intervals

Survey Procedures

- Surveyor to verify nurses are administering medications within their scope of practice
- That the MS has approved the P&P which include the timing of medications
- Verify the hospital has P&P that identify which medications are:
 - Not eligible for scheduled dosing times
 - Eligible for scheduled dosing times and are time-critical and
 - Eligible for scheduled dosing times and are not time-critical.

Survey Procedures

- Surveyor to watch a nurse pass meds and make sure patient is identified
- Make sure nurse follows policy when administering medications
- Surveyor to interview nurses and make sure they understand the hospital policy and timing of medications
- Can the nurses identify time-critical and non-time critical medications?
- Will look at standing orders to make sure they comply with these requirements

Survey Procedures 2014

- Are patients assessed by nursing and/or other staff, per hospital policy, for their risk to their prescribed medications?
- Are patients who are at higher risk and/or receiving high-alert medications monitored for adverse effects?
- Are staff knowledgeable about intervention protocols when patients experience adverse medication-related events?
- If immediate use CSP is prepared outside of pharmacy are practices consistent with USP 797?

Physician Order 406

- **Standard:** Drugs and biologicals can be prepared and administered on the orders contained in pre-printed and electronic standing orders, order sets, and protocols
 - If meets tag 457 requirements which is a new tag number where standing order section was moved to
- An exception is flu and Pneumovac which can be given by protocol approved by the MS after assessment of contraindications
 - Order does not need to be authenticated

Physician Order 406

- Orders for drugs must be documented and signed by practitioners allowed to write them
 - Or signed by practitioners as allowed by state law, state scope of practice, hospital P&P and MS bylaws and R/Rs
- Doctors can write orders and if allowed NP and PAs
- Removed section about use of rubber stamps which is in the medical record chapter anyway
- Adds a section that talks about standing orders

Standing Orders 406

- Nurses or others authorized by hospital P&P and state law may
 - Administer drugs and biologicals in accordance with pre-printed and electronic standing orders, order sets, and protocols
 - CMS collectively just refers to these as standing orders
- Need to address well defined clinical scenarios involving medication administration
- Refers to tag **457** for requirements on standing order P&Ps

Examples of Standing Orders

- Practitioner must still sign off, date, and time
- Chest pain protocol or asthma protocol with Albuterol and Atrovent are an example of initiation of orders
- Code teams gives ACLS drugs in an arrest
- Timing of orders should not be a barrier to effective emergency response
- Preprinted order
 - Should send memo so doctors and providers are aware of the standing order guidelines in tag 457

Verbal Orders 407

- **Standard;** Verbal orders, if used, are be used infrequently
- Verbal orders are a patient safety issue
- Have lead to many errors
- Joint Commission has standard and NPSG, CMS has standard in CMS hospital CoPs, QIO 7th scope of work, National Coordinating Council recommendations
- Rewrite your P&P and Medical staff by-laws to be consistent with these standards
- Repeated VO section in MR starting with tag 454 and reiterated area of verbal orders offer too much room for error

CMS Verbal Orders

- Emphasizes to be used infrequently and never for convenience of the physicians
- This means that physician should not give verbal orders in nursing station if he or she can write them
- Can be used in emergency or if surgeon is scrubbed in during surgery
- Regulation broadens category of practitioners who can sign orders off
- CMS suggests four things in hospital P&P on verbal orders

Verbal Orders P&P Should Include

- Limitations on VO such as not for chemotherapy
 - Include situations in which they may be used
- List the elements for a complete VO
 - Such as patient name, drug, dose, frequency, name of person giving and taking order, etc
- Provide guidelines for clear and effective communications
- Establish identity and authority of practitioner issuing VO

Verbal Orders 408

- **Standard:** When verbal orders are used must be accepted by persons authorized in hospital P&P
- Verbal order for drugs and biologicals may be accepted by individual permitted by state and federal law and hospital P&P to accept
 - For example, hospital allows pharmacists to accept drug orders
 - Hospital will not accept order for Vancomycin IV from an medical assistant in the physicians office
 - It must be from the physician or office nurse
 - Must document order in the chart

Signing Off Verbal Orders

- Physician must sign off a verbal order, **date**, and **time** it when signed off and sign off **ASAP**
- Any physician or practitioner on the case can sign off any VO of another
 - CMS permanently renewed
- This practice must be addressed in the hospital's P&P
- Now a NP or PA may sign off a verbal order, if within their scope (where they had authority to write order) and allowed by state law, hospital policy and delegated to this by the physician

Verbal Orders Changes in

- The 2 main changes were placed in tag 454 in the MR chapter and **not** in the nursing section
 - Regulation states that verbal orders should be authenticated based on state law
 - Some states require order to be signed off in 24 hours or 48 hour
- If no state law **use** to say you had to do this within 48 hours
 - No longer say if no state law follow your P&P but sign **off asap** such as next time the physician sees the patient
 - Need hospital P&P to reflect these guidelines
 - Write it down and repeat it back

Joint Commission Verbal Orders

- RC.02.03.03 (IM 6.50) requires that qualified staff receive and record VO
- Define in writing who can receive and record VO
- Date and document identity of who gave, received, and implemented the order
- Authenticated within time frame law/regulation
- Write it down and read back the completed order or test result

Blood Transfusions and IVs 409

- **Standard:** Blood transfusions and IV medications must be administered with state law and MS P&P
- Use to require special training for this and there was a long list of things that nurses had to be trained on
- CMS eliminated the regulations mandating training for non-physicians who administer IV medication and blood and blood products
 - CMS says because this training is already standard practice but must still be **competent** in those areas
 - Must follow your P&P and state scope of practice

Blood and IV Medication Training

- Must still follow **state law requirements**
 - In some states an LPN can not hang blood
 - Or the LPN can not push certain IV medications in some states
 - Must show they are competent
- Must still have approved **Medical Staff Policies** and Procedures in place
- Staff must follow these which have most of the things that were previously required

Blood Transfusions and IVs 2014

- Hospital P&P for blood and IV medication must be based on state law and MS P&P and must address the following:
 - Vascular access route such as central line, peripheral or implanted port and what medications can be given IV and via what type of access devices
 - Basic safety practices for medication administration
 - Tracing line and tubes prior to administration to be sure proper route
 - Verify proper programming of infusion devices

Blood Transfusions and IVs 2014

- Patient Monitoring
 - Monitor for the effects of the medication since IV medications have a more rapid effect
 - Monitoring to include assessment of risk factors that would influence type and frequency of monitoring
 - Such as patient with renal failure on Vancomycin and dose is based on lab test
- P&P expected to address
 - Monitoring for fluid and electrolyte balance
 - Monitor patients on high alert meds including **opioids** and evaluate for over-sedation and respiratory depression

Blood Transfusions and IVs

- Risk factors for patients receiving opioids include
 - Snoring or history of sleep apnea
 - No recent opioid use or first-time use of IV opioids
 - Increased opioid dose requirement or opioid habituation
 - Longer length of time receiving general anesthesia during surgery
 - Receiving other sedating drugs, such as benzodiazepines, antihistamines, sedatives, or other CNS depressants
 - Preexisting pulmonary or cardiac disease
 - Thoracic or other surgical incisions that may impair breathing

Blood Transfusions and IVs 409 2014

- Hospital P&P is expected to address:
 - Monitoring for fluid and electrolyte balance
 - Monitoring patients for high alert medications including IV opioids
 - Expected to address monitoring for over-sedation and respiratory depression for **safe opioid use**
 - Can erroneously assume patient is asleep when they are having progressive symptoms of respiratory compromise
 - Factors that put patients at high risk include snoring, history of sleep apnea, first time use of IV opioids, increased opioid dose, longer length of time receiving general anesthesia, pulmonary or cardiac disease or thoracic or surgical incisions

Assess and Monitor Patients

- Need to assess and monitor the effects of the medications
- To allow for early identification of adverse effects
- Some may need to use clinical and lab data to evaluate efficacy of medication therapy
 - For **opioids** may need to monitor respiratory status, vital signs such BP, O₂ sat, pain level, sedation scale, and carbon dioxide levels
- Evaluate symptoms such as confusion, agitation, unsteady gait, pruritus, somnolence etc.
- Be aware of high alert medications

Blood Transfusions and IVs

- P&P must include who can conduct the assessments
- The frequency and duration of the assessments
- Under what circumstances practitioners prescribing IV opioids are allowed to establish protocols that differ from hospital P&P
- Assessment includes VS (TPR and BP), pain level, respiratory status, sedation level and ETCO₂
- Also mentions APSF monitoring of opioids including ETCO₂

ISMP Use a Standard Sedation Scale

For Information – Not Required/Not to be Cited

In addition to assessing risk for respiratory depression, the Institute for Safe Medication Practices recommends hospitals use a standard sedation scale when assessing patients receiving PCA. Scales such as the Richmond Agitation Sedation Scale, Pasero, Ramsey, or Glasgow Coma Scale are useful in assessing sedation.

Institute for Safe Medication Practices (ISMP), Medication Safety Alert – Fatal PCA Adverse Events Continue to Happen...Better Patient Monitoring is Essential to Prevent Harm. May 30, 2013

For Information – Not Required/Not to be Cited

Institute for Safe Medication Practices Guidelines for PCA Monitoring

<i>Assessment of Opioid Tolerance</i>	<i>Vital Signs</i>	<i>Pain</i>	<i>Sedation</i>	<i>Respiratory</i>		
				<i>Rate</i>	<i>Quality</i>	<i>SPO₂* &/or ETCO₂**</i>
<i>Baseline Assessment before PCA</i>	<i>X</i>	<i>X</i>	<i>X</i>	<i>X</i>	<i>X</i>	<i>X</i>
<i>PCA Initiation or Change in Drug/Syringe</i> <i>Q 15 minutes x 1 hour</i> <i>Q 1 hour x 4 hours</i> <i>Then Q 2 hours</i>	<i>X</i>	<i>X</i>	<i>X</i>	<i>X</i>	<i>X</i>	<i>X</i>
<i>PCA Dose Change or Bolus</i> <i>Q 1 hour x 4 hours</i> <i>Then Q 2 hours</i>	<i>X</i>	<i>X</i>	<i>X</i>	<i>X</i>	<i>X</i>	<i>X</i>
<i>Adverse Event or Patient Deterioration (e.g., adverse change in sedation score)</i> <i>Q 15 minutes x 1 hour</i> <i>Q 1 hour x 4 hours</i> <i>Then Q 2 hours</i>	<i>X</i>	<i>X</i>	<i>X</i>	<i>X</i>	<i>X</i>	<i>X</i>
<i>Hand-offs/Shift Change</i>	<i>X</i>	<i>X</i>	<i>X</i>	<i>X</i>	<i>X</i>	<i>X</i>

Institute for Safe Medication Practices (ISMP), Medication Safety Alert – Fatal PCA Adverse Events Continue to happen...Better Patient Monitoring is Essential to Prevent Harm. May 30, 2013 ISMP adapted these recommendations from the San Diego Patient Safety Council

** SPO₂: Saturation of peripheral oxygen via pulse oximetry*

Safe Opioid Use & Safe Medication Use

- Patients at great risk for adverse events include age, liver or kidney failure, history of sleep apnea, history of smoking, drug-drug interaction, first time medication use and weight
 - Obesity could increase apnea and smaller patients could more sensitive to dose levels of medications
- Risk factors need to be considered in determining how often to monitor and what type of monitoring
- Must communicate important information in hand-offs such as change of shift

Safe Opioid Use & Safe Medication Use

- ADR, such as opioid-induced respiratory depression require timely intervention as per established hospital protocols
- Must also report to physician or LIP immediately
- High alert medications would want to check VS, O₂ sat, ETCO₂, and sedation levels to prevent respiratory depression and arrest
- **Staff are expected to include patient's reports of his experience of the medication's effects**
- **Educate the patient and family about notifying staff if difficulty breathing**

Safe Opioid Use & Safe Medication Use

- Hospital policy is expected to address the manner and frequency of monitoring
- Hospital P&P is expected to include information to be communicated at shift change
- It is important to document order, medication record, lab reports, vital signs etc.
- Document after actual administration of medication and no documentation in advance
- Surveyor will make sure staff is knowledgeable about intervention protocol if ADE occurs

Anesthesia Patient Safety Foundation

Anesthesia Patient Safety Foundation

- *APSF calls for every patient receiving postoperative opioid analgesics to be managed based on the following clinical considerations*:*
 - *Individualize the dose and infusion rate of opioid while considering the unique aspects of each patient's history and physical status.*
 - *Make continuous monitoring of oxygenation (pulse oximetry) the routine rather than the exception.*
 - *Assess the need for supplemental oxygen, especially if pulse oximetry or intermittent nurse assessment are the only methods of identifying progressive hypoventilation.*
 - *When supplemental oxygen is indicated, monitoring of ventilation may warrant the use of technology designed to assess breathing or estimate arterial carbon dioxide concentrations. Continuous monitoring is most important for the highest risk patients, but depending on clinical judgment, should be applied to other patients.*

When supplemental oxygen is indicated, monitoring of ventilation may warrant the use of technology designed to assess breathing or estimate arterial carbon dioxide concentrations. Continuous monitoring is most important for the highest risk patients, but depending on clinical judgment, should be applied to other patients.

*APSF also has issued a video on opioid induced ventilatory impairment:
http://apsf.org/resources_video4.php*

APSF Website

www.apsf.org

apsf®

Home | Share | Print

Search...

Monitoring OVI POVL POVL Informed Consent Med Safety Fire Safety

About APSF Donors Donate Initiatives Resource Center Grants Contact Us

NEWSLETTER

Click here to read the current issue.

Read Past Issues

MONTHLY POLL

If a patient in your setting has a significant adverse outcome, your group...

- Formally investigates the event within a rigorous, organized framework, such as a QI or other Investigatory (like NTSB) group.
- Notifies the Risk Management department and they investigate. No other investigations performed.
- The group's manager (Chairperson, etc.) is made aware

201

ASA Standards and Guidelines

Contact Us | Support ASA | Site Map

Member Sign-In

Search Terms...

American Society of Anesthesiologists[®]

myASA ASA CALENDAR ASAPAC EDUCATION CENTER JOIN ASA ASA RELATED ORGANIZATIONS SHOP ASA

For Members For Residents and Students For the Public and Media For Health Professionals

www.asahq.org/For-Members/Standards-Guidelines-and-Statements.aspx

Home » For Members » Standards, Guidelines, Statements and Other Documents

myASA Sign-In

UPDATE YOUR PROFILE? Login to MyASA and update your professional profile! [Login Now](#) »

In This Section

[Lifeline to Modern Medicine™](#)

[About ASA](#)

[Practice Management](#)

[About the Anesthesiology Profession](#) »

[Press Room](#) »

[ASA Videos](#)

MEETINGS / EVENTS

[View Calendar of Events](#) »

FDA MEDWATCH ALERTS

ASA FEATURED PRODUCT



[Add to Cart](#)

Anesthesiology Continuing Education (ACE) Program

Standards, Guidelines, Statements and Other Documents

ASA Standards, Guidelines and Statements provide guidance to improve decision-making and promote beneficial outcomes for the practice of anesthesiology. They are not intended as unique or exclusive indicators of appropriate care. The interpretation and application of Standards, Guidelines and Statements takes place within the context of local institutions, organizations and practice conditions. A departure from one or more recommendations may be appropriate if the facts and circumstances demonstrate that the rendered care met the physician's duty to the patient.

Standards provide rules or minimum requirements for clinical practice. They are regarded as generally accepted principles of patient management. Standards may be modified only under unusual circumstances, e.g., extreme emergencies or unavailability of equipment.

Guidelines are systematically developed recommendations that assist the practitioner and patient in making decisions about health care. These recommendations may be adopted, modified, or rejected according to clinical needs and constraints and are not intended to replace local institutional policies. In addition, practice guidelines are not intended as standards or absolute requirements, and their use cannot guarantee any specific outcome. Practice guidelines are subject to revision as warranted by the evolution of medical knowledge, technology, and practice. They provide basic recommendations that are supported by a synthesis and analysis of the current literature, expert opinion, open forum commentary, and clinical feasibility data.

Statements represent the opinions, beliefs, and best medical judgments of the House of Delegates. As such, they are not necessarily subjected to the same level of formal scientific review as ASA Standards or Guidelines. Each ASA member, institution or practice should decide individually whether to implement some, none, or all of the principles in ASA statements based on the sound medical judgment of anesthesiologists participating in that institution or practice.

See also: [Practice Parameters](#)

[Recommendations and Clinical Management Tools - ASA Committees](#)

Blood Transfusions

- Confirm correct patient
- Verify correct blood product
- Standard calls for two qualified persons, one who is administering the transfusion
 - TJC NPSG allows one person hanging blood if use bar coding and CMS confirms this is AABB policy and would be acceptable
- Document monitoring
- P&P include how frequent you monitor the patient and do vital signs
- How to identify and treat and report any adverse transfusion reaction

Blood Components and Blood Administration Procedures

According to the U.S. Department of Health and Human Services, 13,785,000 units of whole blood and red blood cells were transfused in the United States in 2011⁵. The collection, testing, preparation, and storage of blood and blood components are regulated by the Food and Drug Administration. However, administration of blood products via transfusion is governed by §482.23(c)(4). Blood transfusions can be life-saving. However, like IV medications, blood transfusions are not without risk of harm to patients. Transfusion reactions and/or errors can be fatal.

In addition to the safe practices and other safety considerations that apply to all IV medication administration, policies and procedures must address blood administration procedures that are consistent with accepted standards of transfusion practice, including but not limited to:

- *Confirming the following prior to each blood transfusion:*
 - *the patient's identity*
 - *verification of the right blood product for the right patient*

The standard of practice calls for two qualified individuals, one of whom will be administering the transfusion, to perform the confirmation.

- *Requirements for patient monitoring, including frequency and documentation of monitoring*

Blood Transfusions

- Staff must be **competent** in venipuncture
- Competent in using vascular access devices
- Trained in early detection and intervention for opioid over-sedation
- Must document competency
- So make sure nursing education is aware and staff trained in orientation periodically
- Make sure staff educated on P&P

Survey Procedure

- Interview nursing staff on different units who administer IV medications and blood transfusions. Are staff knowledgeable with respect to:
 - Venipuncture techniques
 - Safe medication administration practices, including general practices applying to all types of medications and practices concerning IV tubing and infusion pumps
 - Maintaining fluid and electrolyte balance
 - Patient assessment for risk related to IV medications and appropriate monitoring
 - Early detection and intervention

Survey Procedure

- Will look to see if any blood transfusions
- To review staff files for evidence of competency in administering IV medication and blood products
- Surveyor encouraged to watch staff hang blood or observe IV medication given
 - Were safe injection practices followed
 - Was appropriate access for IV medication
 - Are patients monitored for adverse reactions
 - Were transfused patients correctly identified and correct blood administered?

Incident Reports 410

- **Standard:** There must be procedure for reporting transfusion reactions, adverse drug reactions (ADRs) and errors in administration of drugs
- See tag 508 in the pharmacy section which affects nursing
- Survey procedure
 - Request procedure for reporting
 - They may review the incident reports or other documentation through QAPI program

Transfusion Reactions, ADEs, Drug Errors

- Establish a procedure in the case of ADEs and drug errors when nurses administer drugs or transfusions
- Refers back to tag 508 regarding reporting these into the QAPI system
 - Often done on an incident report
- Transfusion reactions can be serious and life threatening
 - Discussed the symptoms of a transfusion reaction: chills, hives, back pain, bloody urine, dizziness, fever, flank pain, skin flushing, kidney failure, anemia, shock, respiratory failure or death

Transfusion Reactions

- Must have P&P to ensure transfusion reactions are reported
- Must be reported immediately to practitioner
- Must be documented in the chart
- Must be reported to the QAPI program
- Surveyor is suppose to look at the hospital P&P and internal reports of transfusion reactions
- Will ask to see any incident reports

Self Administered Medication 409

- **Standard:** The hospital may allow a patient, or his or her caregiver/support person where appropriate,
 - To self administer medication
 - This includes both hospital-issued medications and the patient's own medications brought into the hospital
 - Must be defined and specified in the hospital's policies and procedures
 - CMS includes PCA as a self administered medication

Only Change in Tag 409 in 2014

- PCA pumps allow for the self-administration of intravenous (IV) medications to patients
- References the section in Tag 409 just discussed concerning assessment and monitoring requirements for post-surgical patients receiving IV opioids
- Including via patient-controlled analgesia (PCA) pumps, in and out of the post-anesthesia care (PACU) and intensive care units (ICU)

Self-Administer Medications

- Previously, the only section on self administered medications was in the pharmacy standard under tag 502
- **Standard:** The hospital may allow a patient or caregiver/support person to self administer medications in accordance with hospital P&P
 - This includes hospital issued medication and patient's own medication brought in
- These are very long sections so need to read

Self-Administer P&P Must Include

- Self administer P&P **must** include:
 - Need an order
 - Make sure assess capacity and document
 - Is the patient competent and not confused
 - Instruct the person on how to give safely
 - Address the security of the medication
 - Document when given in the medical record
 - Assess if receiving opioids including PCA

Self-Administer Medications

- Not required to do
 - Could be beneficial to some patients
- Generally applies to inpatients but may find appropriate situations for outpatients
 - Hospital does for observation patients on Medicare since does not pay for oral medications
 - Asthma patient has inhaler at bedside or patient has hemorrhoid cream or patient learns to give subq Heparin
- Teaching patient to use their medications could avoid readmissions or returns to the ED

Self-Administer Medications

- Some cases nurse may need to supervise
- May want to include in the P&P when supervision by the nurse is needed
- May exclude certain medications from self administration
- Medical staff, nursing and pharmacy departments must collaborate in developing P&P
- Surveyor will assess carefully to ensure these standards and policy requirements are met

Self Administer Medications 413

- Standard: The hospital may allow a patient or caregiver to self administer own medication or hospital issued medications as defined by P&P
- Must have policies to include:
 - Need an order which is consistent with P&P
 - Assess capacity of the patient and document
 - Evaluate the medicine for integrity
 - Address security of the medication
 - Document each medicine given

Other Sections That Impact Nursing

- There are many other sections that impact nursing that are located outside the nursing standards section;
 - Provide copy of patient rights to patient including visitation rights (Tag 95)
 - Review of contracted services (Tag 85)
 - Provision of emergency services in the ED (91)
 - Interpreters for patients with limited English proficiency (116)
 - Grievance process (118)

Other Sections That Impact Nursing

- Informed consent (131)
- Advance directives (132)
- Right to privacy (143)
- Freedom from abuse and neglect (145)
- Confidentiality (146)
- Restraint and seclusion (154)
- PI (274)
- H&P (358)

Other Sections That Impact Nursing

- Autopsies (364)
- Medical records (466)
- Discharge summary (468)
- Pharmacy and medications (490)
- Radiology orders (529)
- Lab services (578)
- Blood and blood components (592)
- Look back program (592)

Other Sections That Impact Nursing

- Dietary policies and assessment (618)
- Utilization review (652)
- Infection control (747)
- Discharge planning (800)
- Organ donation (884)
- Surgery and anesthesia (940,1000)
- Outpatient (1079 and amended 7-16-2012)
- Rehab and respiratory therapy (1123 and 1151)

CMS Proposed Changes



CMS Proposed Changes to Nursing

- CMS proposes changes on June 16, 2016 in the Federal Register and 34 pages long
- The CMS Hospital Improvement Act
- Would amend restraint standards and change LIP to LP so PAs can order and assess patients
- Has a section on access to medical records and QAPI
- Many revisions to infection control
- Would implement and require hospitals to have an antibiotic stewardship program

June 16, 2016 Federal Register



www.gpo.gov/fdsys/pkg/FR-2016-06-16/pdf/2016-13925.pdf

FEDERAL REGISTER

Vol. 81
No. 116

Thursday,
June 16, 2016

Part IV

Department of Health and Human Services

Centers for Medicare & Medicaid Services
42 CFR Parts 482 and 485

Medicare and Medicaid Programs; Hospital and Critical Access Hospital (CAH) Changes To Promote Innovation, Flexibility, and Improvement in Patient Care; Proposed Rule

CMS Proposed Changes to Nursing

- Would clarify that need nursing supervisor and enough staff immediate availability of RN at the bedside to respond when needed
- Must include in policy all the outpatient areas and which ones need a RN
 - Policy must be approved by the MEC
- Must have current nursing care plan
- Must have evidence of reassessment and include patient goals
- All nurses, including agency, must follow P&Ps

CMS Proposed Changes to Nursing

- CNO must evaluate all clinical activities of nurses including agency nurses and other staff
- Must have P&P prohibiting discrimination
 - Can't discriminate against race, color, orientation, disability, etc.
 - Must inform the patient in writing, and in a language they can understand, of their right to be free from discrimination
- Need an order for drugs
- Want to refer to CNO as nursing leadership

This presentation is intended solely to provide general information and does not constitute legal advice. Attendance at the presentation or later review of these printed materials does not create an attorney-client relationship with the presenter(s). You should not take any action based upon any information in this presentation without first consulting legal counsel familiar with your particular circumstances.

Thanks for attending! Questions?



Sue Dill Calloway RN, Esq.
CPHRM, CCMSCP
AD, BA, BSN, MSN, JD

President
Patient Safety and Healthcare
Education

5447 Fawnbrook Lane
Dublin, Ohio 43017
614 791-1468
sdill1@columbus.rr.com
(No email question, Call)