

Table 1**The Various Biotypes of *S. bovis***

<u>Name</u>	<u>Biotype</u>	<u>Clinical correlation</u>
<i>S. gallolyticus</i> subsp. <i>gallolyticus</i>	bovis I	Colon cancer; endocarditis; hepatobiliary disorder
<i>S. gallolyticus</i> subsp. <i>pasteurianus</i>	bovis II/2	Hepatobiliary disorder
<i>S. gallolyticus</i> subsp. <i>macedonicus</i> Synonym: <i>Streptococcus waius</i>	bovis II/2	Hepatobiliary disorder
<i>S. infantarius</i> subsp. <i>coli</i> Possible synonym : <i>Streptococcus lutetiensis</i>	bovis II/1	Colon cancer; meningitis
<i>S. infantarius</i> subsp. <i>infantarius</i>	bovis II/1	Meningitis
<i>S. equinus</i>	bovis I	Usually from animals
<i>S. alactolyticus</i>	none	

Adapted from: Beck M, et al. Comprehensive study of strains previously designated *Streptococcus bovis* consecutively isolated from human blood cultures and emended description of *Streptococcus gallolyticus* and *Streptococcus infantarius* subsp. *coli*. *J Clin Microbiol.* 2008;46:2966-2972.