

# EMERGENCY MEDICINE **REPORTS**

Practical, Evidence-Based Reviews in Emergency Care

SEPTEMBER 20, 2015

VOL. 36, NO. 20

## AUTHORS

**John W. Martel, MD, PhD,**  
Assistant Professor, Tufts University  
School of Medicine, Maine Medical  
Center, Portland, ME.

**Micah Ownbey, MD,** Maine Medical  
Center, Portland, ME.

**Calvin Simmons, MD,** Maine  
Medical Center, Portland, ME.

## PEER REVIEWER

**Todd L. Slesinger, MD, FACEP,**  
FCCM, FCCP, Program Director,  
Emergency Medicine, Aventura  
Hospital and Medical Center, FL.

## STATEMENT OF FINANCIAL DISCLOSURE

To reveal any potential bias in this publication, and in accordance with Accreditation Council for Continuing Medical Education guidelines, we disclose that Dr. Farel (CME question reviewer) owns stock in Johnson & Johnson. Dr. Stapczynski (editor) owns stock in Pfizer, Johnson & Johnson, AxoGen, Walgreens Boots Alliance Inc., and Bristol Myers Squibb. Ms. Mark's (executive editor) spouse works for a company that creates advertising for Uroplasty. Dr. Schneider (editor), Dr. Martel (author), Dr. Ownbey (author), Dr. Simmons (author), Dr. Slesinger (peer reviewer), and Mr. Landenberger (editorial and continuing education director) report no financial relationships with companies related to the field of study covered by this CME activity.

**AHC Media**

## Stroke Mimics: A Clinical Dilemma for the Emergency Physician

*The emergency department (ED) approach to a possible acute stroke used to be easy: ensure the blood pressure and glucose were not too high or low, computed tomography (CT) scan to exclude intracranial hemorrhage, and admit to hospital. Accuracy and rapidity of diagnosis in the ED were not necessarily priorities — alternatives and complications could be sorted out upstairs. Then, tissue plasminogen activator (tPA) for acute ischemic stroke came along. Now, both rapidity and accuracy of diagnosis are expected. As we are finding out, about a quarter of patients with symptoms apparently due to an acute ischemic stroke have another condition that mimics acute stroke. This issue of Emergency Medicine Reports is devoted to a discussion of these stroke mimics, providing guidance to their identification. The opposite of stroke mimics are stroke chameleons — in which acute strokes are misdiagnosed as other conditions. But that discussion is for another day.*

— J. Stephan Stapczynski, MD, FACEP, Editor

## Background and Introduction

Acute ischemic stroke (AIS) is a common and significant cause of mortality and morbidity in the United States, ranking fifth among all causes of death.<sup>1,2</sup> It affects upwards of 800,000 people per year.<sup>3</sup> Approximately 600,000 of these cases represent a first stroke,<sup>4</sup> with the remainder representing recurrent strokes. Stroke is also a leading cause of permanent cognitive and function-limiting disability.<sup>5</sup> Therefore, given the significant associated mortality and morbidity, the American Heart Association's (AHA) Get with the Guidelines initiative emphasizes rapid recognition and potentially aggressive treatment of patients presenting with possible stroke symptoms.<sup>6,7</sup>

Prompt recognition of an AIS is crucial, as prior imaging-based work has demonstrated that the volume of irreversibly damaged brain tissue expands rapidly until reperfusion occurs.<sup>8</sup> In a recent U.S. registry study, rapid recognition followed by early thrombolytic treatment was associated with improved outcomes. For every 15-minute faster interval to treatment, stroke patients experienced reduced mortality and reduced symptomatic intracranial hemorrhage (sICH), as well as increased rates of independent ambulatory status at time of discharge and discharge directly to home.<sup>9</sup>

However, timely recognition and treatment is complicated by the fact that there are multiple conditions that mimic acute ischemic stroke. A comprehensive review suggested that approximately 74% of patients presenting with apparent acute stroke symptoms were ultimately diagnosed with stroke,<sup>10</sup> thus indicating that 26% of patients had their symptoms produced by "stroke

## EXECUTIVE SUMMARY

- Perform point-of-care glucose testing on patients with acute weakness or altered mental status.
- Many stroke mimics produce positive symptoms, whereas strokes predominately produce negative symptoms.
- Most strokes occur in patients with risk factors for cerebrovascular disease, whereas stroke mimics may or may not have such risk factors.
- Often, the diagnosis of stroke mimic can be made only after an assessment has excluded an ischemic stroke.
- The risk of symptomatic intracranial hemorrhage when tPA is given to a patient with a stroke mimic is about 1 in 200 — a rate at least three times lower than that seen in patients with acute ischemic stroke.

mimics.” This estimation was similar to that from a 10-year study of more than 8000 possible stroke patients evaluated by stroke teams, which found that 30% of patients had a stroke mimic.<sup>11</sup> More recent data reflect similar results, with one study focusing on telestroke cases reporting a stroke mimic rate of almost 25%,<sup>12</sup> while another study of patients transferred from outlying hospitals with concern for acute ischemic stroke had a mimic rate of 17%, nearly all of whom received tPA.<sup>13</sup> Therefore, prompt diagnosis is complicated by a multitude of stroke mimic etiologies, including structural intracranial abnormalities, infection, syncope, vertigo, seizure and migraine patterns, as well as underlying psychiatric causes and demyelinating diseases.

While the incidence of stroke mimics varies among the published studies, there are commonalities. Seizure, migraine, psychogenic disorders, and toxic/metabolic etiologies have been widely reported as among the most common non-vascular conditions that mimic stroke.<sup>14,15,16</sup> Data from a single hospital identified that common stroke mimics included seizures, syncope, sepsis, and functional disorders/conversion disorders.<sup>17</sup> (See *Figure 1*.) Another center reported that toxic/metabolic disorders and seizure etiologies were most common.<sup>14</sup>

The wide array of potential stroke mimic etiologies often renders the clinical picture very difficult to distinguish from an acute ischemic process at the time of ED presentation. A factor that further complicates management is that patients may present within the appropriate thrombolysis administration window and subsequently receive this therapy. While it has been suggested

that up to 14% of mimic patients receive thrombolytics,<sup>14,15,18</sup> it has also been postulated that there is decreased risk of associated adverse effects in this population.<sup>14</sup> There are no current data to suggest that tPA administration leads to adverse events in patients presenting with clinical features concerning for ischemic stroke but who are ultimately diagnosed with a stroke mimic.<sup>19,20</sup>

The key to avoiding unnecessary and potentially harmful therapy is improved understanding of the various mimic entities, as well as attention paid to important historical and physical exam features. In considering stroke symptoms, it can be helpful to think in terms of positive and negative symptoms. It has been explained elsewhere that positive symptoms are associated with excess central nervous system (CNS) activity, and are characterized by visual disturbance such as flashing lights and perception of various geometric shapes. In addition, somatosensory symptoms are associated with pain and/or paresthesias, and motor symptoms with jerking extremity movements. In contrast, negative symptoms are associated with a decrease in CNS function, and commonly manifest as acute vision or hearing loss, as well as sensorimotor deficits in the extremities.<sup>17</sup> As a rule, acute ischemic stroke impairs CNS function manifested by negative symptoms. The three most common are unilateral subjective arm weakness, unilateral subjective leg weakness, and reported speech disturbance; positive symptoms are uncommon. Ischemia in the major cerebral vascular territories produces classic findings. (See *Table 1*.)


Although it may be difficult to rapidly differentiate such a wide array of etiologies from ischemic stroke in the

ED environment, it has been suggested that several key features are associated with a potential stroke mimic profile, including younger patients, milder symptoms, arrival to the ED by means other than emergency medical services (EMS), and lack of cardiovascular risk factors.<sup>11</sup> Furthermore, several studies have suggested that patients with stroke mimics exhibit a lower prevalence of ischemic stroke risk factors.<sup>15,19</sup> One large prospective study reported that specific historical factors associated with increased odds of having a stroke mimic included absence of hypertension, hyperlipidemia, and atrial fibrillation.<sup>11</sup> This is a dilemma that emergency physicians will have to continually address as the treatment for acute stroke continues to evolve. The primary objective in this article is to review common stroke mimic etiologies, as well as discuss the key historical and physical exam characteristics associated with stroke mimics, and to present a diagnostic framework for approaching such cases.

### Common Stroke Mimics: Seizure

Among stroke mimic diagnoses, seizure is one of the most common, being the final diagnosis in up to 20% of stroke mimics.<sup>21,22</sup> For those patients with a prior stroke history, seizures may originate from previous ischemic or hemorrhagic stroke foci.<sup>17</sup> The distinction is less complicated when the patient presents following a witnessed account of seizure activity and has a known history of epilepsy. A constellation of typical symptoms, including generalized convulsions, involuntary movements, postictal confusion, tongue-biting, and incontinence, make an acute ischemic event less likely.<sup>23</sup> The clinical

**Figure 1. The Most Common Stroke Mimics**



Reprinted with permission from: Fernandes PM, Whiteley WN, Hart SR, et al. Strokes: Mimics and chameleons. *Pract Neurol* 2013;13:21–28.

**Table 1. Vascular Territory and Ischemic Stroke Features**

Territory	Features
Anterior cerebral	Weakness and sensory loss, predominantly in the contralateral leg Aphasia Cognitive, behavioral, and emotional disorders may be prominent findings
Middle cerebral (anterior portion involving motor cortex)	Contralateral hemiparesis If involving dominant hemisphere: Broca's (expressive or nonfluent) aphasia
Middle cerebral (posterior portion involving sensory cortex)	Contralateral hemineglect and sensory deficits Ipsilateral gaze preference Hemianopia If involving dominant hemisphere: receptive aphasia
Lacunar (deep penetrating small arteries) — 5 classic subtypes	1. Pure motor — contralateral weakness, facial weakness, ± dysarthria 2. Sensorimotor — contralateral hemiparesis, hemisensory loss 3. Pure sensory — contralateral sensory loss 4. Ataxic hemiparesis — homolateral ataxia and weakness 5. Dysarthria — clumsy hand — dysarthria, one hand clumsiness, ± central facial paralysis, dysphagia, tongue deviation
Posterior cerebral	Ipsilateral cranial nerve palsy with contralateral motor and/or sensory defect Loss of conjugate gaze Homonymous visual field defect

picture is clearly complicated by post-seizure paresis, and in one study this was the most frequent final diagnosis made among stroke mimics that received

tPA.<sup>24</sup> Although actual seizure activity is a relative tPA contraindication, more than 8% of patients in one study exhibited seizure activity after acute stroke

symptom onset.<sup>25</sup> Previous magnetic resonance imaging (MRI)-based studies have suggested that postictal dysfunction may relate to transient cerebral perfusion mismatch associated with epileptic foci but not with an ischemic lesion in a discrete vascular distribution, as would be seen in a stroke.<sup>26</sup>

Postictal paresis, or Todd's paresis (or paralysis), may be difficult to distinguish from stroke. Reports vary with regard to the incidence and duration, with some studies suggesting that it occurs in as many as 10% of patients with generalized seizures, and may last for several hours or even longer in some cases.<sup>23</sup> However, the vast majority of postictal deficits are commonly short, with one video monitoring study reporting a median time of 174 seconds in the case of focal epilepsy.<sup>27</sup> Deficits may persist longer in the case of generalized seizure, with reports of postictal deficits lasting days. Importantly, focal motor deficits are not the only possible deficit following a seizure. Postictal dysphasia has also been reported to follow dominant hemisphere seizures.<sup>23</sup> MRI with diffusion-weighted imaging (DWI) can assist in the differentiation between acute stroke and prior ischemic event, and may provide important information regarding the potential for post-stroke secondary seizure disorder.

## Migraine

Migraine headache is a frequent ED complaint and leads to approximately 1 million patient visits annually.<sup>28</sup> Migraine disproportionately affects females three times as often as males, with a peak incidence around age 40.<sup>28</sup> Patients commonly present with a positive prior history and no evidence of focal neurological deficit on physical exam. However, in the setting of focal neurological deficit, primary headache disorders comprise up to 10% of stroke mimics.<sup>17</sup> For instance, familial hemiplegic migraine is an autosomal dominant disorder with high penetrance. Since this presents with unilateral neurological deficits, it is particularly difficult to differentiate from acute stroke; however, there is a common pattern, as patients tend to be teenagers at the time of onset, and 70% are female.<sup>29</sup> Likewise, migraine in general is thought

to have a heritable component; if both parents have a history of migraine, there is a nearly 75% chance a patient will suffer from the disorder.<sup>30</sup> It is characterized by recurrent severe headaches, classically of a unilateral and throbbing nature. Nausea, vomiting, photophobia, and phonophobia are common associated symptoms. Onset is generally gradual, and headaches may last up to three days.<sup>31</sup> One particular diagnostic challenge is that nearly 30% of acute stroke patients report headache at the time of stroke symptom onset.<sup>32</sup>

Multiple theories have been proposed to explain the phenomenon of migraine headaches. The physiology of migraines is complex, with vasodilatory and inflammatory components. Although some alterations in local blood flow have been noted in patients experiencing migraines, the primary mechanism seems to be a triggered abnormality in the ability of the brain to manage sensory input.<sup>33</sup> There are multiple subtypes that will be described in greater detail below.

Common migraines do not present with aura, and are the majority of cases. However, 20-30% of migraines are associated with aura, and may include a variety of neurological complaints and symptoms.<sup>33</sup> These are known as classic migraines. It is these patients who are likely to raise suspicion for stroke when they present to the ED. By definition, migraine with aura is diagnosed after exclusion of organic disease, specifically TIA, ischemic stroke, and intracranial hemorrhage,<sup>31</sup> in some patients with dense focal neurological deficits, this likely requires neuroimaging. Migraine auras generally exhibit a gradual onset over at least 5 minutes, and last a maximum of 60 minutes.<sup>33</sup> The headache usually starts within one hour of the aura but may not always occur; it will lack the sudden/maximal onset or thunderclap component attributed to acute subarachnoid hemorrhage. Migraine auras can include a variety of neurological symptoms. Visual symptoms are most common, including scotomas, flashes/sparks of light, and zigzag-type visual phenomena.<sup>31</sup>

Although less common, paresthesias, motor disturbances, cognitive changes, and language deficits may

also be seen in conjunction with aura. Ophthalmoplegic migraine includes involvement of cranial nerves III, IV, or VI with associated eye movement palsies and pupil changes. The third cranial nerve is most commonly involved, and the headache is ipsilateral to the deficits. Retinal migraine may result in scotoma or blindness. In the descriptively named above, the aura includes motor symptoms such as hemiparesis or hemiplegia. In contrast to many ischemic stroke syndromes, the onset is typically gradual and associated with a visual, sensory, or speech disturbance. Basilar-type migraines present with an aura suggestive of brainstem vascular territory deficits and may exhibit multiple neurological findings, including visual loss (e.g., blindness), dysarthria, tinnitus, vertigo, bilateral paresthesias, paresis, and altered mentation.

In rare cases, an aura may become permanent. If this occurs, and an area of infarct is confirmed by neuroimaging in the distribution of the aura, it is termed a migrainous infarction or migrainous stroke.<sup>31</sup> The pathophysiology of this phenomenon is unclear and remains, thankfully, rare. In the event that a patient with known migraine with aura experiences new or worsening aura symptoms or experiences aura that persists longer than normally would be expected (e.g., > 60 minutes), consider neurology consultation and advanced imaging.

## Functional Disorders

Underlying psychiatric disease poses one of the most difficult challenges in the clinical diagnosis of acute stroke vs. stroke mimic, as patients may present with acute motor or sensory weakness in the setting of a specific emotional trigger.<sup>29,34</sup> In particular, conversion disorder is a frequent cause of stroke mimic; in one study it was deemed the underlying etiology in nearly 75% of patients ultimately diagnosed with stroke mimic who received thrombolytics.<sup>13</sup> Despite this finding, the prevalence of conversion disorder is rare overall, and it is essentially a diagnosis of exclusion not made in the acute care setting. Conversion disorder typically occurs more commonly in women,

with the majority of cases seen in early childhood and adolescence. Complaints are widely varied, and may include paralysis, aphonia, seizures, ataxia/coordination difficulty, blindness, and anesthesia/paresthesia. Historically speaking, there may be a disproportionate lack of concern relative to the severity of symptoms (*la belle indifférence*). With regard to physical exam, the hallmark is deficit inconsistency observed during serial physical exams, motor-sensory impairments with incongruent vascular territory distribution, and inconsistent task-dependent weakness or “give-away weakness.”<sup>17</sup> However, this has also been observed in patients with concomitant organic disease and cannot be used to definitively diagnose conversion disorder.<sup>35</sup> As such, the diagnosis is generally made only following admission and extensive neurological workup.

Functional disorders as a whole are thought to represent a physical manifestation of emotional distress,<sup>36</sup> with common triggers, including panic attacks and dissociative episodes.<sup>34</sup> Although providers may be tempted to directly confront patients in the acute setting, patients often lack insight into their condition, and the approach may consequently worsen the symptoms. In some cases, providing reassurance to the patient may rapidly improve or resolve symptoms, and if there is an acute stressor identified, then all efforts should be made to remove this from the equation.

Consider an underlying functional disorder if the symptoms are not consistent with a discrete vascular territory, if there are no objective neurological findings, and if there is an inconsistent physical exam. Unfortunately, there is no diagnostic study that can reliably provide this diagnosis. Patients with underlying psychiatric pathology also develop concomitant organic disease, and there are cases of patients being diagnosed with a functional disorder but ultimately found to have organic neurological pathology to explain their symptoms.<sup>35</sup> Providers should be cautious in documenting a non-anatomic exam, atypical presentation, inconsistent physical exam, or a patient's lack of concern about their symptoms as

**Table 2. Peripheral Neuropathies**

Nerve Involved	Mechanism/History	Signs and Symptoms
Radial nerve compression	“Saturday night palsy” compression of the radial nerve in the spiral groove on the humerus, occurs from sleeping on the outstretched arm, prolonged compression over the back of a chair, etc.	Wrist drop, finger extensor weakness, sensory loss on the dorsum of the hand
Ulnar nerve compression	Most commonly occurs at the elbow due to compression from leaning on a table or similar mechanism	May cause weakness in the interosseous muscles, decreased grip, sensory loss of the fourth and fifth digits
Cervical radiculopathy	Nerve root compression at the neck	Radicular pain is a key feature, specific distribution of weakness will depend on the nerve root C5: Shoulder abduction C6: Wrist extension C7: Wrist flexion C8: Finger flexion T1: Finger adduction and abduction
Lumbosacral radiculopathy	Spinal nerve root compression in the vertebral foramina, herniated disc	Radicular pain is a key feature, deficits will depend on the nerve root L2-4: Hip flexion, knee extension L5: Foot dorsiflexion S1: Leg extension, foot plantar flexion
Common peroneal nerve compression	Compression of the peroneal nerve at the neck of the fibula, leg crossing, squatting, cast	Foot drop, sensory loss on dorsum of foot and lateral shin
Sciatic nerve compression	Compression of the sciatic nerve at the sciatic notch, trauma, prolonged bed rest or seated position, piriformis syndrome	Radicular pain to the back of the thigh and leg, weakness in the lower leg muscles, numbness sparing the medial calf

justification for not aggressively evaluating for acute stroke.

## Systemic Infections

Sepsis may present with stroke-like features, and the most common associated infection is a urinary tract infection.<sup>37</sup> Commonly associated symptoms include hyperthermia/hypothermia, tachycardia, and altered mental status, with or without an obvious historical or physical exam indication of an infectious source. However, the fact that fever may not be present and that a source etiology may not be immediately obvious during initial evaluation warrants consideration of an acute neurological insult as a potential cause of altered mental status.

In some cases, sepsis is a stroke risk factor, as severe sepsis has been shown to induce a hypercoagulable state.<sup>38</sup> Furthermore, acute infection may result in recurrence of the focal neurological deficits initially present in patients who have a history of prior stroke, even when there is no residual deficit post-rehabilitation at baseline.<sup>39</sup> Alternatively, there

also may be cases of acute stroke with concomitant secondary infection, such as aspiration pneumonia.<sup>17</sup>

Obtaining a thorough past medical history, either from the patient or a family member, to establish baseline functional status is crucial but not always possible in the acute setting. When the history is unclear, objective focal neurological deficits should be assumed to stem from an acute neurological insult if history, vital signs, and physical exam do not clearly suggest an underlying infectious etiology. However, the timeframe of symptom onset may add some clarity to the evaluation. Sepsis is generally a gradual process and is less likely to present with the abrupt onset of acute, focal neurological deficits. In addition to more common sepsis sources, infections of the CNS, such as brain abscess, meningitis, and encephalitis, should also be considered in the setting of altered mental status.

## Space-Occupying Lesions

Space-occupying brain lesions, including primary CNS tumors,

metastases, and abscesses, have been shown to cause stroke-like focal neurological deficits attributed to alterations in cerebral blood flow associated with mass effect.<sup>40</sup> However, mass effect is thought to progress slowly over a 24–48 hour period, and does not necessarily present with the sudden, acute change typically associated with an ischemic stroke process.<sup>41</sup> Structural brain lesions are an infrequent cause of stroke mimic, and are subsequently less common than seizure, hypoglycemia, syncope, sepsis, functional disorders, and primary headache.<sup>42</sup> Fortunately, noncontrast enhanced brain CT is readily available in almost all EDs and can assist in the rapid diagnosis of most acute space-occupying brain pathology, as well as acute intracranial hemorrhage.

## Hypoglycemia

Transient hypoglycemia, easily assessed by bedside fingerstick glucose, commonly manifests with acute mental status changes and autonomic symptoms such as diaphoresis and tachycardia. However, it also has long been

known to manifest with a constellation of stroke-like symptoms such as hemiplegia and aphasia in certain cases.<sup>43</sup> Hypoglycemia-associated stroke-like presentations have been attributed to a wide variety of etiologies, including insulin or sulfonylurea medication use/overdose and alcohol consumption,<sup>44</sup> as well as less common endocrine sources such as Addisonian crises and insulinomas.<sup>17</sup> When rapidly recognized, hypoglycemia is easily correctable, and patients often quickly improve; however, there are reports of symptoms taking up to several hours to fully resolve.<sup>45</sup> It is very important to address this prior to consideration of tPA administration, and national stroke guidelines mandate that a blood glucose level be obtained prior to tPA administration.<sup>7</sup>

## Peripheral Neuropathy

Peripheral neuropathies are due to cellular level changes in nerves distal to the brain and spinal cord. In addition to potential toxic-metabolic, ischemic, or inflammatory involvement, the anatomic distribution of these structures relative to surrounding bony and soft-tissue structures places them at risk for acute compression and entrapment.<sup>24</sup>

Mononeuropathies may cause focal sensorimotor deficits, and in some instances may easily be mistaken for stroke symptoms when they occur in an acute timeframe. For instance, acute hand weakness may be seen in contralateral cortical strokes that resemble ulnar and radial nerve neuropathy. An acute peripheral nerve injury with subsequent myelin damage may be followed by Wallerian degeneration and result in sensorimotor dysfunction, muscle atrophy, and areflexia.<sup>24</sup>

The history and chronicity of symptoms will aid in distinguishing between acute neuropathy and stroke. However, symptoms associated with acute nerve compression and ischemia may exhibit similar chronicity, and subsequently are more likely to be confused. A high-quality physical exam in addition to detailed history is key to rapidly differentiating between these two entities. Table 2 details common peripheral neuropathy syndromes that can cause significant weakness and thus be confused with stroke.

One common mononeuropathy is radial nerve palsy, which is classically termed “Saturday night palsy.” This may occur with prolonged radial nerve compression (as with an arm draped over a chair) or with other prolonged periods of upper extremity immobilization such as during sleep. Compressive radial neuropathies typically occur distal to the branches that innervate the triceps muscle. Although a patient with radial neuropathy is likely to exhibit weakness with wrist and finger extension, arm extension is expected to remain intact, whereas a stroke patient might have concomitant triceps weakness in addition to the hand and wrist deficits. In addition, radial neuropathy sensory deficit localizes to the hand dorsum, while in acute stroke it is likely to be more volar in nature.<sup>46</sup>

## Unilateral Facial Paralysis

Unilateral facial motor weakness is seen in both central and peripheral nerve pathologies. It involves upper motor neurons/contralateral motor cortex lesions or lower motor neurons, respectively. The two pathological foci may be distinguished by a notable difference in the distribution of motor weakness on physical exam. Peripheral lesions (e.g., Bell’s palsy) typically manifest as impairment in both upper and lower facial muscle function, whereas central lesions exhibit forehead sparing due to bilateral cortical innervation, therefore resulting in intact forehead wrinkling. In addition to differences in the distribution of central compared to peripheral facial paralysis, Bell’s palsy also may exhibit additional findings not typically seen in stroke, including decreased tear production (in some cases may also be excessive), acute gustatory impairment, and increased auditory acuity and facial hyperesthesia.<sup>47</sup>

Bell’s palsy is the most common cause of unilateral idiopathic facial palsy. It is often preceded by prodromal illness and is commonly attributed to herpes simplex virus infection.<sup>48</sup> It has been more recently associated with Lyme disease in endemic regions. Lyme disease is the most common vector-mediated infection in the United States, is caused by the spirochete *Borrelia burgdorferi*, and is vectored by ticks in the genus *Ixodes*.

Bell’s palsy represents the most common neurologic abnormality in the second stage of this important disease, and up to 25% of Bell’s palsy cases are attributed to Lyme disease.<sup>49</sup> Furthermore, it is responsible for half of pediatric facial paralysis cases in endemic regions.<sup>50</sup>

## Vertigo

Acute dizziness is a very common complaint in the ambulatory clinic setting, comprising up to 5% of visits. This represents nearly 10 million such visits annually, with 25% presenting in the ED setting.<sup>51</sup> Acute dizziness is somewhat open to interpretation and is often described as vertigo, presyncope, and other unspecified unsteadiness in space that persists anywhere from minutes to hours, and is often accompanied by nausea, vomiting, nystagmus, and intolerance with head position change.<sup>52</sup> Acute vestibular syndrome (AVS) presents with these symptoms, and is generally presumed to be a viral or post-viral phenomenon. Peripheral AVS generally occurs in the absence of focal neurologic deficits (such as acute unilateral motor and/or sensory deficits and gaze palsy) that would be associated with stroke.<sup>52</sup>

The diagnostic challenge posed in the acute setting is that 35% of posterior stroke cases present with dizziness as the chief complaint,<sup>53</sup> whereas up to 3% of patients with acute dizziness have an ischemic stroke involving the posterior circulation. Ten percent of isolated posterior circulation stroke patients will present with isolated vertigo.<sup>54</sup> However, the vast majority of patients presenting with dizziness are not having an acute stroke, and several groups have reported historical and diagnostic considerations for distinguishing these two entities clinically. Posterior stroke usually manifests with a constellation of neurological deficits and not isolated vertigo. Vertigo red flags suggesting a potential stroke etiology include any neurological exam deficit, total ipsilateral hearing loss, new ataxia with inability to walk without support, and direction-changing nystagmus.<sup>53</sup> In particular, 71% of patients diagnosed with cerebellar stroke reported acute inability to walk without support.<sup>54</sup> The vast majority of patients with serious pathology underlying their dizziness are older,

have multiple cardiovascular risk factors, and have abnormal neurological exams in addition to dizziness, as up to 40% of posterior strokes are attributed to an embolic source.<sup>54</sup>

Thorough neurological examination also helps differentiate between these two disparate causes of dizziness. Ataxia requiring new ambulatory support and direction-changing nystagmus are particularly important in evaluating vertigo, since one of these two signs was seen in 84% of patients with cerebellar infarction and isolated vertigo.<sup>54</sup> Furthermore, performing the Horizontal Head Impulse Test (h-HIT) also may help distinguish between peripheral and central etiologies specifically in the setting of vertigo. The h-HIT is a head acceleration maneuver performed on alert patients with active vertigo and designed to interrogate the vestibulo-ocular reflex.<sup>55</sup> It has been well described in the literature and essentially consists of examiner-directed random movement of the patient's head. Patients visually fixate on the examiner's eyes while the head is moved randomly from side to side. A positive test result indicates a peripheral lesion and manifests as a corrective saccade such that as the patient's head is turned away, the patient's eyes initially do not track the examiner but are followed by a rapid visual correction such that the eyes snap back to the original fixation point. Likewise, the lack of a corrective saccade raises suspicion for a central vertigo etiology.

The Head Impulse-Nystagmus Test of Skew (HiNTS), a variation of the h-HIT, approaches 100% sensitivity and 96% specificity for diagnosing posterior stroke when more than one component is abnormal.<sup>53</sup> However, a considerable amount of training is required for diagnostic efficacy with this approach, and a combination of imaging and neurological consultation may be required to differentiate between peripheral and central etiologies.

## Neurodegenerative Disorders

Acute demyelinating disorders, as well as initial diagnosis or acute exacerbation of multiple sclerosis (MS), may present with a wide variety of focal neurological

deficits, including hemiparesis and cranial nerve or visual deficits.<sup>24</sup> Episodes of acute diplopia, dysarthria, ataxia, and sensory deficits, often exacerbated by warm weather, are common with MS.<sup>56</sup>

Optic neuritis is an acute inflammatory disorder of the optic nerve that may present with sudden unilateral vision loss, and occurs more frequently in young adult women.<sup>57</sup> It is a common initial manifestation of MS, occurring in 15-20% of these patients. In addition, up to half of MS patients may develop the condition at some point during their disease course.<sup>58</sup>

Acute disseminated encephalomyelitis (ADEM) is a post-infectious or post-immunization immune-mediated demyelinating disorder attributed to transient autoimmune destruction of myelin<sup>59</sup> with multifocal brain lesions visible on MRI that generally occurs within three to six weeks of the inciting event.<sup>60</sup> This time course, as well as a wide array of concomitant symptoms, including acute altered mental status, ataxia, optic neuritis, and sensorimotor deficits, aid in distinguishing this from an acute stroke syndrome.

Acute sensorimotor deficits also may be associated with non-compressive transient spinal cord inflammation. Although acute myelopathy is seen in the setting of ADEM,<sup>56</sup> it may also manifest as transverse myelitis.<sup>59</sup> This condition is generally not associated with acute back pain, and patients present with a wide variety of acute neurological deficits, including focal sensorimotor weakness and potential impairment of bowel, bladder, and sexual function below the lesion level.<sup>61,62</sup>

Similarly, Guillain-Barré syndrome is an acute inflammatory demyelinating polyneuropathy (AIDP) comprised of several subtypes. Although the less common of these varieties may be difficult to distinguish from acute stroke, it typically manifests as a progressive, ascending bilateral paralysis characterized by peripheral motor weakness, hyporeflexia, and, in advanced cases, respiratory distress when there is compromise of diaphragm innervation.<sup>24</sup> In contrast, acute strokes typically manifest with unilateral deficits as previously described. However, brainstem infarcts in the basilar artery territory may

present with a myriad of symptoms, including initial hemiparesis that may progress to bilateral limb weakness or locked-in syndrome, altered sensorium, vertigo, ataxia, bulbar motor weakness with gaze palsies or miotic pupils, as well as so-called "crossed-findings" with ipsilateral cranial nerve deficits and motor or sensory findings on the opposite side.<sup>63,64</sup>

## Summary: Approach to Stroke Mimics

Having reviewed common stroke mimics, we can now consider how to approach evaluation of the patient with stroke-like symptoms. It is important to remember that acute stroke likely remains the most life-threatening and time-sensitive diagnosis for patients presenting to the ED with acute, focal neurological deficits. With few exceptions, the diagnosis of stroke mimic will only be made following appropriate resuscitation, laboratory evaluation, and advanced imaging, as well as neurology consultation.

The American Heart Association has developed guidelines for the initial assessment of suspected stroke,<sup>7</sup> and stroke mimics are included in the list of alternative diagnoses to consider during the initial assessment. Recommended ancillary tests are basic blood tests, including serum glucose (rapid fingerstick is initially preferred for expedience), electrolytes, blood urea nitrogen/creatinine (BUN/Cr), complete blood count, troponin, and coagulation studies (PTT, PT/INR, possibly thrombin time [TT] or Ecarin Clotting Test [ECT] if taking direct thrombin inhibitors), as well as 12-lead electrocardiogram (ECG) and oxygen saturation. In most community hospitals, noncontrast enhanced CT scan of the brain is readily available and adequate to rapidly exclude intracranial hemorrhage in a patient eligible for tPA therapy.<sup>7</sup> In specialized stroke centers, emergent MRI may be available to acutely map ischemic brain tissue and provide assistance for reperfusion therapy.

As with all patients, a thorough history and physical exam should be performed in a timely manner, including a baseline NIH stroke scale (NIHSS). Two or more of three key symptoms

**Table 3. Clinical Questions in the Evaluation of Stroke Mimics**

Age and other demographic data: Is there a high a priori probability of a cerebrovascular event?	As noted above, certain demographic and historical factors have been shown to correlate with the likelihood of an event representing a stroke vs. a stroke mimic.
Nature of the symptoms: Positive vs. negative	Stroke is more likely to present with negative symptoms, whereas mimic due to migraine aura, conversion disorder, or other causes is more likely to have positive symptoms.
Onset and progression	Sepsis may be a slower onset, while seizure, stroke, and even tumor may present suddenly.
Duration	Rapidly resolving symptoms raise the question of a TIA or seizure.
Precipitating factors	Deficits preceded by a seizure or illness may indicate an underlying diagnosis.
Associated symptoms, for example, headache, loss of awareness, during or after the attacks	Headache may be associated with stroke but also with migraine. Loss of awareness may increase suspicion of seizure.
Adapted with permission from: Nadarajan V, Perry RJ, Johnson J, et al. Transient ischaemic attacks: Mimics and chameleons. <i>Pract Neurol</i> 2014;14:23–31.	

(acute facial paralysis, arm drift, or speech change) are associated with increased odds of having sustained an acute stroke (odds ratio [OR] 5.5; 95% CI: 3.3–9.1). Conversely, the absence of all three of these symptoms is thought to be helpful in identifying patients unlikely to have had an acute ischemic stroke.<sup>65</sup> Particular historical factors have been associated with acute stroke diagnosis in multiple studies, and include older age, higher NIHSS, and a history of atrial fibrillation, hypertension, or diabetes. Likewise, patients who are young, have a low initial NIHSS, are initially unresponsive, have neurological exams incongruent with an anatomic vascular distribution, and patients with a history of epilepsy were more likely to be diagnosed with stroke mimics.<sup>11,13,21,22,39,66–69</sup> One study reported that a history of psychiatric diagnosis was associated with greater likelihood of ultimate stroke mimic diagnosis.<sup>13</sup>

Some stroke mimics are more easily detectable. For instance, if hypoglycemia is identified, correct blood glucose levels and evaluate for functional response. If a space-occupying lesion is noted per noncontrast brain CT (ICH, tumor, shift, abscess), tPA is clearly contraindicated, and further diagnostics and

treatment should be lesion-specific. If symptoms rapidly resolve and were preceded by a witnessed seizure or there is a history of epilepsy, consider Todd's paralysis as a cause of focal neurological deficits. Although postictal exam findings in the potential unwitnessed seizure that support this diagnosis include tongue biting, bowel or urinary incontinence, and confusion, these objective findings may also be seen with acute stroke in some cases. In the event of a new focal neurological deficit, the initial workup remains unchanged, and if symptoms persist the patient likely warrants admission with neurological consultation. This type of presentation should be assumed to stem from an acute stroke until proven otherwise.

Specific attention should be given to episodes of prior neurological insult or known pathology, including stroke, seizures, migraine, and mass lesions. If available, a well-documented physical exam is very useful for comparison with a particular patient's clinical picture at the time of presentation. For patients having sustained a prior stroke, all potential focal neurological deficits should be considered within the context of their medical history, past presenting symptoms, and subsequent residual

deficits. Although this information may help provide context for the severity of a patient's symptoms, it is important to carefully consider subjective components of the patient's complaint as well as objective exam findings during surveillance for acute stroke pathology.

Table 3 includes a list of questions for providers to consider when a patient presentation is concerning for transient ischemic attack vs. a mimic.<sup>23</sup> These criteria can also be applied to stroke, and may help to develop a differential diagnosis based on symptoms and time course. With any potential stroke patient, frequent re-evaluation during the workup is useful both to help clarify the extent of symptoms as well as to monitor for improvement or deterioration. Temporal changes in symptoms over time may gradually move the level of suspicion closer to or farther from acute stroke.

Nevertheless, ED providers are in a unique position to rapidly identify acute neurological pathology, and it is incumbent on the front-line physician to consider acute stroke first. However, given the wide array of clinical mimics that complicate this process, a detailed understanding of such peripheral entities will enable rapid diagnosis of easily correctable entities (e.g., hypoglycemia) and will expediently resolve exactly which patients are appropriate for acute treatment (e.g., tPA) or require immediate subspecialist support (e.g., neurosurgery) from those for whom a diagnosis of exclusion to explain their symptoms will likely occur during hospitalization.

## References

- Centers for Disease Control and Prevention, National Center for Health Statistics. Compressed Mortality File 1999–2009. CDC Wonder Online Database, Compiled for Compressed Mortality File 1999–2009 Series 20, No. 20, 2012. Underlying Cause-of-Death 1999–2009. [wonder.cdc.gov/Mortsq1.html](http://wonder.cdc.gov/Mortsq1.html). Accessed June 1, 2015.
- Kochanek KD, Xu JQ, Murphy SL, et al. Mortality in the United States, 2013. NCHS Data Brief, No. 178. Hyattsville, MD: National Center for Health Statistics, Centers for Disease Control and Prevention, US Dept. of Health and Human Services; 2014.
- Mozzafarian D, Benjamin EJ, Go AS, et al. Heart disease and stroke statis-

- tics — 2015 update: A report from the American Heart Association. *Circulation* 2015;131:e29–322.
4. Go AS, Mozaffarian D, Roger VL, et al; American Heart Association Statistics Committee and Stroke Statistics Subcommittee. Heart disease and stroke statistics — 2014 update: A report from the American Heart Association. *Circulation* 2014;129:e28–e292.
  5. Centers for Disease Control and Prevention (CDC). Prevalence and most common causes of disability among adults — United States, 2005. *MMWR Morb Mortal Wkly Rep* 2009;58:421–426.
  6. Goldstein LB, Bushnell CD, Adams RJ, et al. Guidelines for the primary prevention of stroke: A guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke* 2010;42:517–584.
  7. Jauch EC, Saver JL, Adams HP Jr, et al. Guidelines for the early management of patients with acute ischemic stroke: A guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke* 2013;44:870–947.
  8. Saver JL. Time is brain — quantified. *Stroke* 2006;37:263–266.
  9. Saver JL, Fonarow GC, Smith EE, et al. Time to treatment with intravenous tissue plasminogen activator and outcome from acute ischemic stroke. *JAMA* 2013;309:2480–2488.
  10. Goldstein LB, Simel DL. Is this patient having a stroke? *JAMA* 2005;293:2391–2402.
  11. Merino JG, Luby M, Benson RT, et al. Predictors of acute stroke mimics in 8187 patients referred to a stroke service. *J Stroke Cerebrovasc Dis* 2013;22:397–403.
  12. Ali SF, Viswanathan A, Singhal AB, et al. The telestroke mimic (TM)-Score: A prediction rule for identifying stroke mimics evaluated in a telestroke network. *J Am Heart Assoc* 2014 Jun 23;3:e000838.
  13. Mehta S, Vora N, Edgell RC, et al. Stroke mimics under the Drip-and-Ship Paradigm. *J Stroke Cerebrovasc Dis* 2014;23:844–849.
  14. Brunser AM, Illanes S, Lavados PM, et al. Exclusion criteria for intravenous thrombolysis in stroke mimics: An observational study. *J Stroke Cerebrovasc Dis* 2013;22:1140–1145.
  15. Chernyshev OY, Martin-Schild S, Albright KC, et al. Safety of tPA in stroke mimics and neuroimaging-negative cerebral ischemia. *Neurology* 2010;74:1340–1345.
  16. Hand PJ, Kwan J, Lindley RI, et al. Distinguishing between stroke and mimics at the bedside: The Brain Attack Study. *Stroke* 2006;37:769–775.
  17. Fernandes PM, Whiteley WN, Hart SR, et al. Strokes: Mimics and chameleons. *Pract Neurol* 2013;13:21–28.
  18. Hess DC, Audebert HJ. The history and future of telestroke. *Nat Rev Neurol* 2013;9:340–350.
  19. Scott PA, Silbergleit R. Misdiagnosis of stroke in tissue plasminogen activator-treated patients: Characteristics and outcomes. *Ann Emerg Med* 2003;42:611–618.
  20. Zinkstok SM, Engelter ST, Gensicke H, et al. Safety of thrombolysis in stroke mimics: Results from a multicenter cohort study. *Stroke* 2013;44:1080–1084.
  21. Forster A, Griebe M, Wolf ME, et al. How to identify stroke mimics in patients eligible for intravenous thrombolysis? *J Neurol* 2012;259:1247–1253.
  22. Tsvigoulis G, Alexandrov AV, Chang J, et al. Safety and outcomes of intravenous thrombolysis in stroke mimics. *Stroke* 2011;42:1771–1774.
  23. Nadarajan V, Perry RJ, Johnson J, et al. Transient ischaemic attacks: Mimics and chameleons. *Pract Neurol* 2014;14:23–31.
  24. Magauran BG, Nitka M. Stroke mimics. *Emerg Med Clin North Am* 2012;30:795–804.
  25. Bladin CF, Alexandrov AV, Bellavance A, et al. Seizures after stroke: A prospective multi-centre study. *Arch Neurol* 2000;57:1617–1622.
  26. Rupperecht S, Schwab M, Fitzek C, et al. Hemispheric hypoperfusion in postictal paresis mimics early brain ischemia. *Epilepsy Res* 2010;89:355–359.
  27. Gallmetzer P, Leutmezer F, Serles W, et al. Postictal paresis in focal epilepsies—Incidence, duration, and causes. *Neurology* 2004;62:2160–2164.
  28. Vinson DR. Treatment patterns of isolated benign headache in US emergency departments. *Ann Emerg Med* 2002;39:215–222.
  29. Thomsen LL, Eriksen MK, Roemer SF, et al. A population-based study of familial hemiplegic migraine suggests revised diagnostic criteria. *Brain* 2002;125:1379–1391.
  30. MacGregor EA. In the clinic: Migraine. *Ann Intern Med* 2013;159:ITC5-1–ITC5-15.
  31. Headache Classification Committee of the International Headache Society. The International Classification of Headache Disorders, 3rd ed. *Cephalalgia* 2013;33:629–808.
  32. Tentschert S, Wimmer R, Greisenegger S, et al. Headache at stroke onset in 2196 patients with ischemic stroke or transient ischemic attack. *Stroke* 2005;36:e1–3.
  33. Kwiatkowski T, Friedman BW. Headache disorders. In: Marx J, ed. *Rosen's Emergency Medicine: Concepts and Clinical Practice*, 8th edition. Elsevier Health Sciences; 2014.
  34. Stone J, Warlow C, Sharpe M. Functional weakness: Clues to mechanism from the nature of onset. *J Neurol Neurosurg Psychiatry* 2012;83:67–69.
  35. Winter AO. Somatoform Disorders. In: Marx J, ed. *Rosen's Emergency Medicine: Concepts and Clinical Practice*, 8th edition. Elsevier Health Sciences; 2014.
  36. Feinstein F. Conversion disorder: Advances in our understanding. *CMAJ* 2011;183:915–920.
  37. Tuntiyatorn L, Saksornchai P, Tunlayadechanont S. Identification of stroke mimics among clinically diagnosed acute strokes. *J Med Assoc Thai* 2013;96:1191–1198.
  38. Valtonen VV. Infection as a risk factor for infarction and atherosclerosis. *Ann Med* 1991;23:539–543.
  39. Hemmen TM, Meyer BC, McClean TL, et al. Identification of nonischemic stroke mimics among 411 code strokes at the University of California, San Diego, Stroke Center. *J Stroke Cerebrovasc Dis* 2008;18:23–25.
  40. Ueno Y, Tanaka A, Nakayama Y. Transient neurological deficits simulating transient ischemic attacks in a patient with 34 meningioma — case report. *Neurol Med Chir (Tokyo)* 1998;38:661–665.
  41. Hacke W, Schwab S, Horn M, et al. 'Malignant' middle cerebral artery territory infarction: Clinical course and prognostic signs. *Arch Neurol* 1996;53:309–315.
  42. Morgenstern LB, Frankowski RF. Brain tumor masquerading as stroke. *J Neurooncol* 1999;44:47–52.
  43. Montgomery BM, Pinner CA, Newberry SC. Transient hypoglycemic hemiplegia. *Arch Int Med* 1964;114:680–684.
  44. Andrade R, Mathew V, Morgenstern MJ, et al. Hypoglycemic hemiplegic syndrome. *Ann Emerg Med* 1984;13:529–531.
  45. Wallis WE, Donaldson I, Scott RS, et al. Hypoglycemia masquerading as cerebrovascular disease (hypoglycemic hemiplegia). *Ann Neurol* 1985;18:510–512.
  46. Tarulli A. *Neurology: A Clinician's Approach*. Cambridge University Press, 2010: 91.
  47. Marendra S, Olsson J. The evaluation of facial paralysis. *Otolaryngol Clin North Am* 1997;30:669.
  48. Murakami S, Mizobuchi M, Nakashiro Y, et al. Bell palsy and herpes simplex virus: identification of viral DNA in endoneurial fluid and muscle. *Ann Intern Med* 1996;124:27–30.

49. Ho K, Melanson M, Desai JA. Bell palsy in Lyme disease-endemic regions of Canada: A cautionary case of occult bilateral peripheral facial nerve palsy due to Lyme disease. *Can J Emerg Med* 2012;14:321-324.
50. Cook SP, Macartney KK, Rose CD, et al. Lyme disease and seventh nerve paralysis in children. *Am J Otolaryngol* 1997;18:320.
51. Newman-Toker DE, Hsieh YH, Camargo CA Jr, et al. Spectrum of dizziness visits to US emergency departments: Cross-sectional analysis from a nationally representative sample. *Mayo Clin Proc* 2008;83:765-775.
52. Tarnutzer AA, Berkowitz AL, Robinson KA, et al. Does my dizzy patient have a stroke? A systematic review of bedside diagnosis in acute vestibular syndrome. *CMAJ* 2011;183:E571-E592.
53. Kattah, JC, Talkad AV, Wang DZ, et al. HINTS to diagnose stroke in the acute vestibular syndrome: Three-step bedside oculomotor examination more sensitive than early MRI diffusion-weighted imaging. *Stroke* 2009;40:3504-3510.
54. Lee H, Sohn SI, Cho YW, et al. Cerebellar infarction presenting isolated vertigo: Frequency and vascular topographical patterns. *Neurology* 2006;67:1178-1183.
55. Halmagyi GM, Curthoys IS. A clinical sign of canal paresis *Arch Neurol* 1988;45:737-739.
56. Li Y, Zeng C, Luo T. Paroxysmal dysarthria and ataxia in multiple sclerosis and corresponding magnetic resonance imaging findings. *J Neurol* 2011;258:273-276.
57. The Optic Neuritis Study Group. Multiple sclerosis risk after optic neuritis: Final optic neuritis treatment trial follow-up. *Arch Neurol* 2008;65:727-732.
58. Marcel C, Anheim M, Flamand-Rouviere C, et al. Symptomatic paroxysmal dysarthria-ataxia in demyelinating diseases. *J Neurol* 2010;257:1369-1372.
59. Tenembaum S, Chitnis T, Ness J, et al. Acute disseminated encephalomyelitis. *Neurology* 2007;68 (16 Suppl 2):S23-36.
60. Alexander M, Murthy JMK. Acute disseminated encephalomyelitis: Treatment guidelines. *Ann Indian Acad Neurol* 2011;14(Suppl 1):S60-S64.
61. Beh SC, Greenberg BM, Frohman T, et al. Transverse myelitis. *Neurol Clin* 2013;31:79-138.
62. Frohman EM, Wingerchuk DM. Clinical practice. Transverse myelitis. *N Engl J Med* 2010;363:564-572.
63. Ferbert A, Brückmann H, Drummen R. Clinical features of proven basilar artery occlusion. *Stroke* 1990;21:1135-1142.
64. Nouh A, Remke J, Ruland S. Ischemic posterior circulation stroke: A review of anatomy, clinical presentations, diagnosis, and current management. *Front Neurol* 2014;5:30.
65. Gibson LM, Whiteley W. The differential diagnosis of suspected stroke: A systematic review. *J R Coll Physicians Edinb* 2013;43:114-118.
66. Arto V, Putaala J, Strbian D, et al. Stroke mimics and intravenous thrombolysis. *Ann Emerg Med* 2012;59:27-32.
67. Vroomen PC, Buddingh MK, Luijckx GJ, et al. The incidence of stroke mimics among stroke admissions in relation to age group. *J Stroke Cerebrovasc Dis* 2008; 17:418-422.
68. Chang J, Teleb M, Yang JP, et al. A model to prevent fibrinolysis in patients with stroke. *J Stroke Cerebrovasc Dis* 2012;21:839-843.
69. Valle J, Lopera E, Guillan M, et al. Stroke mimics: A challenge for the emergency physician. *An Sist Sanit Navar* 2014;37:117-128.

## CME Questions

- The AHA stroke guideline identifies one blood test result that is mandatory prior to the administration of tPA. This information is also useful in ruling out an important and reversible cause of stroke mimic. Which of the following is it?
  - sodium
  - PT/INR
  - glucose
  - hemoglobin
- A 73-year-old white female presents with mild confusion and mild right leg weakness after a witnessed fall at home. Her initial NIH stroke scale is 2, scoring for leg weakness and mild confusion. After quickly reviewing her medical history, she is noted to have a history of hypertension, myocardial infarction, atrial fibrillation, seizures, emphysema, and gout. What combination of historical factors below would collectively identify the patient most likely to be having a true stroke vs. a mimic?
  - female, hypertension, myocardial infarction
  - older age, hypertension, atrial fibrillation
  - atrial fibrillation, emphysema, seizures
  - myocardial infarction, gout, low stroke scale
  - Caucasian, low stroke scale, hypertension
- A 38-year-old female with a history of migraine headache and hypertension presents with unilateral face and tongue paresthesias starting 4 hours ago, followed by a pulsating headache, nausea, and photophobia. She notes a history of similar vision changes prior to her regular headache episodes. Normally this lasts a few minutes and resolves around

E. creatinine

## EMERGENCY MEDICINE REPORTS

### CME Objectives

Upon completion of this educational activity, participants should be able to:

- recognize specific conditions in patients presenting to the emergency department;
- apply state-of-the-art diagnostic and therapeutic techniques to patients with the particular medical problems discussed in the publication;
- discuss the differential diagnosis of the particular medical problems discussed in the publication;
- explain both the likely and rare complications that may be associated with the particular medical problems discussed in the publication.

- the time her headache starts, but this time the visual symptoms have persisted. Which of the following most strongly indicates a need for neuroimaging?
- type of aura symptom
  - duration of aura
  - severity of headache
  - history of hypertension
- Occasionally tPA is given to patients who are subsequently diagnosed with stroke mimic. Which of the following is true of these patients?
    - The risk of intracranial hemorrhage following tPA is higher in stroke mimic patients compared with those patients ultimately diagnosed with stroke.
    - Some of the most common final diagnoses among stroke mimic patients receiving tPA are seizure, migraine, and conversion disorder.
    - It is very rare that patients given tPA are diagnosed with stroke mimic, occurring in < 1% of tPA patients.
    - Stroke mimic patients make up an entirely different demographic set relative to stroke patients.
  - In addition to being a risk factor for stroke, sepsis may also mimic stroke by precipitating focal neurological symptoms. What is the most common type of infection to cause stroke mimic?
    - space-occupying CNS infection (i.e., brain abscess)
    - cellulitis
    - encephalitis/meningitis
    - pneumonia
    - urinary tract infection
  - Postictal focal neurological deficits (Todd's paresis) may mimic stroke; symptoms may last for minutes to hours, and even longer in some cases. What percentage of patients with generalized seizures will have this manifestation?
    - < 1%
    - 10%
    - 25-50%
    - > 50%
  - A 28-year-old male presents with report of having spent the day at a bar, and complains of left hand and wrist weakness with associated paresthesias. Which exam findings suggest a peripheral rather than a central cause of his deficits?
    - diminished palmer sensation
    - inability to extend the left elbow
    - focal weakness with finger and wrist extension
    - dysmetria
  - A 57-year-old male is transported per EMS with aphasia and inability to comply with NIH stroke scale measurement. His wife reports that he consumed several alcoholic drinks per day and that she tries to make sure that he receives all of his medications every day. He is tachycardic and diaphoretic, but afebrile, and there is no recent history of acute illness. What is the most likely etiology of his presentation?
    - intracranial hemorrhage
    - aspiration pneumonia
    - acute ischemic stroke
    - hypoglycemia
  - A 21-year-old female presents per private vehicle with complaint of diplopia and "seeing colors," diffuse paresthesias, and intermittent arm weakness in the setting of a progressively worsened headache that started 6 hours prior to arrival. Patients presenting with a stroke mimic as the underlying cause of their focal neurological deficits are most likely to have the following historical and demographic characteristics:
    - arrival to the ED by means other than EMS.
    - milder symptoms.
    - lack of cardiovascular risk factors.
    - all of the above.
  - A 27-year-old, otherwise healthy female presents with history of recurrent prior episodes of diplopia, and one day of unilateral vision loss. She reports that the diplopia has often been worse during warm summer days. Her symptoms warrant evaluation for:
    - occipital brain neoplasm.
    - multiple sclerosis.
    - ischemic cerebellar stroke.
    - pituitary adenoma.

Is there an article or issue you'd like posted to your website? Interested in a custom reprint? There are numerous opportunities to leverage editorial recognition to benefit your brand. Call us at 877-652-5295 or email [ahc@wrightsmedia.com](mailto:ahc@wrightsmedia.com) to learn more.

To obtain information and pricing on group discounts, multiple copies, site-licenses, or electronic distribution please contact:

**TRIA KREUTZER**

Phone: (800) 688-2421, ext. 5482  
Email: [tria.kreutzer@ahcmedia.com](mailto:tria.kreutzer@ahcmedia.com)

To reproduce any part of AHC newsletters for educational purposes, please contact The Copyright Clearance Center for permission:

Email: [info@copyright.com](mailto:info@copyright.com)  
Website: [www.copyright.com](http://www.copyright.com)  
Phone: (978) 750-8400

## CME INSTRUCTIONS

To earn credit for this activity, please follow these instructions:

- Read and study the activity, using the references for further research.
- Scan the QR code at right or log onto [AHCMedia.com](http://AHCMedia.com) and click on My Account. *First-time users must register on the site.*
- Pass the online tests with a score of 100%; you will be allowed to answer the questions as many times as needed to achieve a score of 100%.
- After completing the test, your browser will be directed to the activity evaluation form.
- Once the completed evaluation is received, a credit letter will be e-mailed to you instantly.



## EDITORS

**Sandra M. Schneider, MD**  
Professor, Emergency Medicine  
Hofstra North Shore—LIJ  
School of Medicine  
North Shore University Hospital  
Manhasset, New York

**J. Stephan Stapczynski, MD**  
Clinical Professor of Emergency Medicine  
Scholarly Projects Advisor  
University of Arizona College of Medicine  
- Phoenix  
Emergency Department, Maricopa  
Integrated Health System

## EDITORIAL BOARD

**Paul S. Auerbach, MD, MS, FACEP**  
Professor of Surgery  
Division of Emergency Medicine  
Department of Surgery  
Stanford University School of Medicine  
Stanford, California

**William J. Brady, MD, FACEP, FAAEM**  
Professor of Emergency Medicine and  
Medicine, Medical Director, Emergency  
Preparedness and Response, University  
of Virginia Operational Medical  
Director, Albemarle County Fire Rescue,  
Charlottesville, Virginia; Chief Medical  
Officer and Medical Director, Allianz  
Global Assistance

**Michael L. Coates, MD, MS**  
Professor  
Department of Family and Community  
Medicine  
Wake Forest University School  
of Medicine  
Winston-Salem, North Carolina

**Alasdair K.T. Conn, MD**  
Chief of Emergency Services  
Massachusetts General Hospital  
Boston, Massachusetts

**Charles L. Emerman, MD**  
Chairman  
Department of Emergency Medicine  
MetroHealth Medical Center  
Cleveland Clinic Foundation  
Cleveland, Ohio

**Chad Kessler, MD, MHPE**  
Deputy Chief of Staff, Durham VAMC  
Chairman, VHA Emergency Medicine  
Field Advisory Committee  
Clinical Associate Professor, Departments  
of Emergency Medicine and Internal  
Medicine  
Duke University School of Medicine  
Durham, North Carolina

**Kurt Kleinschmidt, MD, FACEP, FACMT**  
Professor of Surgery/Emergency  
Medicine  
Director, Section of Toxicology  
The University of Texas Southwestern  
Medical Center and Parkland Hospital  
Dallas, Texas

**Frank LoVecchio, DO, FACEP**  
Vice-Chair for Research  
Medical Director, Samaritan Regional  
Poison Control Center  
Emergency Medicine Department  
Maricopa Medical Center  
Phoenix, Arizona

**Larry B. Mellick, MD, MS, FAAP, FACEP**  
Professor, Department of Emergency  
Medicine and Pediatrics  
Georgia Regents University  
Augusta, Georgia

**Paul E. Pepe, MD, MPH, FACEP, FCCM, MACP**  
Professor of Medicine, Surgery,  
Pediatrics, Public Health and Chair,  
Emergency Medicine  
The University of Texas Southwestern  
Medical Center and Parkland Hospital  
Dallas, Texas

**Charles V. Pollack, MA, MD, FACEP**  
Chairman, Department of Emergency  
Medicine, Pennsylvania Hospital  
Associate Professor of Emergency  
Medicine  
University of Pennsylvania School of  
Medicine  
Philadelphia, Pennsylvania

**Robert Powers, MD, MPH**  
Professor of Medicine and Emergency  
Medicine  
University of Virginia  
School of Medicine  
Charlottesville, Virginia

**David J. Robinson, MD, MS, MMM, FACEP**  
Professor and Vice-Chairman of  
Emergency Medicine  
University of Texas Medical School at  
Houston  
Chief of Emergency Services, LBJ General  
Hospital, Harris Health System  
Houston, Texas

**Barry H. Rumack, MD**  
Professor Emeritus of Pediatrics and  
Emergency Medicine  
University of Colorado School of  
Medicine  
Director Emeritus  
Rocky Mountain Poison and Drug Center  
Denver, Colorado

**John A. Schriver, MD**  
Chief, Department of Emergency Services  
Rochester General Hospital  
Rochester, New York

**David Sklar, MD, FACEP**  
Professor of Emergency Medicine  
Associate Dean, Graduate Medical  
Education  
University of New Mexico School of  
Medicine  
Albuquerque, New Mexico

**Charles E. Stewart, MD, EMDM, MPH**  
Claremore Indian Hospital  
Claremore, Oklahoma

**Gregory A. Volturo, MD, FACEP**  
Chairman, Department of Emergency  
Medicine  
Professor of Emergency Medicine and  
Medicine  
University of Massachusetts Medical  
School  
Worcester, Massachusetts

**Steven M. Winograd, MD, FACEP**  
St. Barnabas Hospital  
Clinical Assistant Professor, Emergency  
Medicine  
New York College of Osteopathic  
Medicine  
Old Westbury, New York

**Allan B. Wolfson, MD, FACEP, FACP**  
Program Director,  
Affiliated Residency in Emergency  
Medicine  
Professor of Emergency Medicine  
University of Pittsburgh  
Pittsburgh, Pennsylvania

### CME Question Reviewer

**Roger Farel, MD**  
Retired  
Newport Beach, CA

© 2015 AHC Media LLC. All rights reserved.

## EMERGENCY MEDICINE REPORTS™

(ISSN 0746-2506) is published every other week by  
AHC Media LLC, One Atlanta Plaza, 950 East Paces  
Ferry Road NE, Suite 2850, Atlanta, GA 30326.  
Telephone: (800) 688-2421 or (404) 262-7436.

**Editorial & Continuing Education  
Director:** Lee Landenberger

**Executive Editor:** Shelly Morrow Mark

**GST Registration No.:** R128870672

Periodicals Postage Paid at Atlanta, GA 30304 and at  
additional mailing offices.

**POSTMASTER:** Send address changes  
to **Emergency Medicine Reports**,  
P.O. Box 550669, Atlanta, GA 30355.

Copyright © 2015 by AHC Media LLC, Atlanta, GA.  
All rights reserved. Reproduction, distribution, or  
translation without express written permission is strictly  
prohibited.

**Back issues: \$31.** Missing issues will be fulfilled  
by customer service free of charge when contacted  
within one month of the missing issue's date.

## SUBSCRIBER INFORMATION

### CUSTOMER SERVICE: 1-800-688-2421

Customer Service E-Mail Address:  
customerservice@ahcmedia.com

Editorial E-Mail Address:  
shelly.mark@ahcmedia.com

Website:  
www.AHCMedia.com

### SUBSCRIPTION PRICES

1 year with 65 ACEP/65 AMA/39 AAFP  
Category 1/Prescribed credits: \$564

1 year *without credit*: \$419  
Add \$19.99 for shipping & handling

### MULTIPLE COPIES:

Discounts are available for group subscriptions,  
multiple copies, site-licenses, or electronic  
distribution. For pricing information, call  
Tria Kreutzer at 404-262-5482.

One to nine additional copies:  
**\$359 each;**  
10 or more additional copies:  
**\$319 each.**

All prices U.S. only. U.S. possessions and  
Canada, add \$30 plus applicable GST. Other  
international orders, add \$30.

## ACCREDITATION

AHC Media is accredited by the Accreditation Council for Continuing  
Medical Education to provide continuing medical education for physicians.

AHC Media designates this enduring material for a maximum of 65 *AMA  
PRA Category 1 Credits™*. Each issue has been designated for a maximum  
of 2.50 *AMA PRA Category 1 Credits™*. Physicians should claim only credit  
commensurate with the extent of their participation in the activity.

Approved by the American College of Emergency Physicians for a  
maximum of 65.00 hour(s) of ACEP Category I credit.

This Enduring Material activity, *Emergency Medicine Reports*, has  
been reviewed and is acceptable for up to 39.00 Prescribed credits by  
the American Academy of Family Physicians. Term of approval begins  
01/01/2015. Term of approval is for one year from this date. Each  
monograph is approved for 1.50 Prescribed credits. Credit may be claimed  
for one year from the date of each monograph. Physicians should claim  
only the credit commensurate with the extent of their participation in the  
activity.

The American Osteopathic Association has approved this continuing  
education activity for up to 65 AOA Category 2-B credits.


This is an educational publication designed to present scientific  
information and opinion to health professionals, to stimulate thought,  
and further investigation. It does not provide advice regarding medical  
diagnosis or treatment for any individual case. It is not intended for  
use by the layman. Opinions expressed are not necessarily those of  
this publication. Mention of products or services does not constitute  
endorsement. Clinical, legal, tax, and other comments are offered for  
general guidance only; professional counsel should be sought for specific  
situations.

This CME activity is intended for emergency and family physicians. It is in  
effect for 36 months from the date of the publication.

# EMERGENCY MEDICINE REPORTS

## Stroke Mimics: A Clinical Dilemma for the Emergency Physician

### The Most Common Stroke Mimics



Reprinted with permission from: Fernandes PM, Whiteley WN, Hart SR, et al. Strokes: Mimics and chameleons. *Pract Neurol* 2013;13:21–28.

### Vascular Territory and Ischemic Stroke Features

Territory	Features
Anterior cerebral	Weakness and sensory loss, predominantly in the contralateral leg Aphasia Cognitive, behavioral, and emotional disorders may be prominent findings
Middle cerebral (anterior portion involving motor cortex)	Contralateral hemiparesis If involving dominant hemisphere: Broca’s (expressive or nonfluent) aphasia
Middle cerebral (posterior portion involving sensory cortex)	Contralateral hemineglect and sensory deficits Ipsilateral gaze preference Hemianopia If involving dominant hemisphere: receptive aphasia
Lacunar (deep penetrating small arteries) — 5 classic subtypes	1. Pure motor — contralateral weakness, facial weakness, ± dysarthria 2. Sensorimotor — contralateral hemiparesis, hemisensory loss 3. Pure sensory — contralateral sensory loss 4. Ataxic hemiparesis — homolateral ataxia and weakness 5. Dysarthria — clumsy hand — dysarthria, one hand clumsiness, ± central facial paralysis, dysphagia, tongue deviation
Posterior cerebral	Ipsilateral cranial nerve palsy with contralateral motor and/or sensory defect Loss of conjugate gaze Homonymous visual field defect

### Peripheral Neuropathies

Nerve Involved	Mechanism/History	Signs and Symptoms
Radial nerve compression	“Saturday night palsy” compression of the radial nerve in the spiral groove on the humerus, occurs from sleeping on the outstretched arm, prolonged compression over the back of a chair, etc.	Wrist drop, finger extensor weakness, sensory loss on the dorsum of the hand
Ulnar nerve compression	Most commonly occurs at the elbow due to compression from leaning on a table or similar mechanism	May cause weakness in the interosseous muscles, decreased grip, sensory loss of the fourth and fifth digits
Cervical radiculopathy	Nerve root compression at the neck	Radicular pain is a key feature, specific distribution of weakness will depend on the nerve root C5: Shoulder abduction C6: Wrist extension C7: Wrist flexion C8: Finger flexion T1: Finger adduction and abduction
Lumbosacral radiculopathy	Spinal nerve root compression in the vertebral foramina, herniated disc	Radicular pain is a key feature, deficits will depend on the nerve root L2-4: Hip flexion, knee extension L5: Foot dorsiflexion S1: Leg extension, foot plantar flexion
Common peroneal nerve compression	Compression of the peroneal nerve at the neck of the fibula, leg crossing, squatting, cast	Foot drop, sensory loss on dorsum of foot and lateral shin
Sciatic nerve compression	Compression of the sciatic nerve at the sciatic notch, trauma, prolonged bed rest or seated position, piriformis syndrome	Radicular pain to the back of the thigh and leg, weakness in the lower leg muscles, numbness sparing the medial calf

## Clinical Questions in the Evaluation of Stroke Mimics

Age and other demographic data: Is there a high a priori probability of a cerebrovascular event?	As noted above, certain demographic and historical factors have been shown to correlate with the likelihood of an event representing a stroke vs. a stroke mimic.
Nature of the symptoms: Positive vs. negative	Stroke is more likely to present with negative symptoms, whereas mimic due to migraine aura, conversion disorder, or other causes is more likely to have positive symptoms.
Onset and progression	Sepsis may be a slower onset, while seizure, stroke, and even tumor may present suddenly.
Duration	Rapidly resolving symptoms raise the question of a TIA or seizure.
Precipitating factors	Deficits preceded by a seizure or illness may indicate an underlying diagnosis.
Associated symptoms, for example, headache, loss of awareness, during or after the attacks	Headache may be associated with stroke but also with migraine. Loss of awareness may increase suspicion of seizure.
Adapted with permission from: Nadarajan V, Perry RJ, Johnson J, et al. Transient ischaemic attacks: Mimics and chameleons. <i>Pract Neurol</i> 2014;14:23–31.	