

# PRIMARY CARE REPORTS

The Practical CME Journal for Primary Care and Family Physicians

April 2018

VOL. 24, NO. 4

## AUTHORS

**Elizabeth Fernandez, MD,**  
Attending Physician, St. John's  
Riverside Hospital, Yonkers, NY;  
Adjunct Clinical Assistant Professor,  
Lake Erie College of Osteopathic  
Medicine, Erie, PA

**Bjorn Bernhardsen, MD,  
MBA, MPH,** Resident Physician,  
St. John's Riverside Hospital,  
Yonkers, NY

## PEER REVIEWER

**Steven M. Winograd, MD,  
FACEP,** Mt. Sinai Queens  
Hospital Center, Jamaica Queens,  
NY; Assistant Clinical Professor of  
Emergency Medicine, Mt. Sinai  
Medical School, New York City;  
Assistant Clinical Professor of  
Emergency Medicine, NYITCOM,  
Old Westbury, NY

## STATEMENT OF FINANCIAL DISCLOSURE

To reveal any potential bias in this publication, and in accordance with Accreditation Council for Continuing Medical Education guidelines, Dr. Wise (editor) reports he is involved with sales for CNS Vital Signs. Dr. Hocum (pharmacist reviewer) reports he is an employee of United Therapeutics. Dr. Fernandez (author), Dr. Bernhardsen (author), Dr. Winograd (peer reviewer), Ms. Coplin (executive editor), Ms. Mark (executive editor), and Ms. Hatcher (editorial group manager) report no financial relationships with companies related to the field of study covered by this CME activity.

**RELIAS**  
Formerly AHC Media

## Wound Care Management

Patients with lacerations commonly present to primary care physicians. Although managing most lacerations is routine, some cases are complex, requiring physician judgment for effective treatment. This article will discuss the principles of laceration management, incorporating recent research in wound care. The authors performed a PubMed search using terms such as laceration, sutures, closure, and wound to identify relevant articles to review. This article reviews evidence-based wound care practices, including prehospital presentation, anesthesia, irrigation, free closure methods, and post-closure care. As will be noted, many common practices in wound care have not been subjected to randomized, controlled trials, suggesting there might be room for improvement.

### Wound Healing Etiology/Pathophysiology

Natural wound healing is a complex process involving multiple steps. After the initial insult, the first process is coagulation, which then progresses through the stages of inflammation, collagen deposition, contraction, and epithelialization.<sup>1</sup>

Platelets activated during coagulation stimulate an inflammatory cell response, attracting phagocytic cells to remove debris and bacteria. The inflammatory phase lasts three to four days, followed by the proliferative phase, which is characterized by collagen deposition within the wound. Typically, through day 21, collagen is cross-linked in a disorganized manner, producing firm tissue that is characteristic of a healed wound. This is followed by epithelialization and wound contraction the year following injury.<sup>2</sup>

While it is not clinically relevant to readily recall all aspects of the wound healing cycle, it is important that the provider understand the factors that may interfere with wound healing. Patient factors, such as inherited collagen-vascular disorders, poor tissue oxygenation (typically found in smokers and patients with poorly controlled diabetes), and even poor diet, all can contribute to poor wound healing. The provider's role is to ensure a clean, well-approximated wound that will best aid the body's natural wound repair ability.

### Differential Diagnosis

Lacerations are caused most often by straightforward processes, such as impact from falls or sharp edges. Even in these common circumstances, consider complicating factors that may alter care needs. The history can be invaluable in guiding care considerations, but sometimes may be incomplete or missing entirely. All wounds should be assessed for cause and

## EXECUTIVE SUMMARY

Primary care physicians usually can care for minor wounds in their offices. This article highlights key elements in first-line management.

- There is no “golden hour” for wound repair. Clinically uninfected wounds in highly vascular areas can be closed many hours after injury with an acceptable risk for infection.
- Topical anesthetic or cryotherapy can reduce the pain of local anesthetic infiltration.

- Adhesive strips, skin staples, and tissue adhesives are time-efficient methods to close wounds not under tension.
- The cosmetic outcome for absorbable and nonabsorbable sutures is generally equivalent.
- Prophylactic antibiotics are indicated for all cat, most human, and some dog bites.

potential for morbidity due to possible foreign bodies and bacterial contamination. Also, patient cooperation can complicate wound care. Tetanus vaccination status and the need for prophylactic antibiotics should be considered. With some mechanisms of injury, such as a fall, other and less obvious injuries should not be overlooked. Finally, since providers may not reliably estimate wound size visually, proper measuring should be used.<sup>3</sup>

### Prehospital Care

Prehospital care should follow Advanced Trauma Life Support (ATLS) guidelines, including hemorrhage control, dressing placement, and immobilization as needed. If direct pressure fails to control bleeding, elevation and/or tourniquet placement may be necessary. Any foreign objects within the wound, either from the initial injury or subsequent injury, ideally should be left in place until definitive care is undertaken at a hospital. Other ATLS principles, such as airway maintenance or protection and oxygenation, should be followed. Patients with complex lacerations, especially with severe bleeding or deep tissue involvement, may require direct transport to a trauma center for evaluation.<sup>4</sup>

### Evaluating the Wound

#### History

Traditional practice has dictated the importance not only of the mechanism of injury, but also the duration of time since the injury in influencing

the risk for wound infection. This is based on the concept of bacterial contamination of the wound at the time of injury or colonization of the wound after injury due to the exposure of subepithelial tissue. In 1975, Robson et al reported that between three and five hours after injury, bacteria count within traumatic wounds was at a level associated with infection, suggesting the potential for infection if the wound was closed after that time period.<sup>5</sup> This principle was observed in a study by Waseem et al published in 2012.<sup>6</sup> Of 297 patients analyzed, the 10 patients who developed a wound infection had a median wound closure time of 867 minutes compared to 330 minutes in the 287 patients who did not develop a wound infection ( $P = 0.03$ ). As the authors noted, this result did not control for confounding factors known to influence the rate of wound infection after closure, nor were they able to identify a threshold value that differentiated the two groups.

The idea that there is a threshold value for the risk of wound infection after closure of traumatic lacerations has been challenged in clinical observational studies over the past 30 years, although the early studies were relatively small, had a low rate of follow-up, did not use current approaches to wound closure, and had less rigorous methods for assessing outcome. There are two recent studies of sufficient size and rigor to address the issue of traumatic wound closure timing.

The first was published by Van Den Baar et al in 2010.<sup>7</sup> It was a study of 408 patients treated at a Dutch Level

I trauma center that found there was no difference in the rate of infection of wounds treated before and after six hours of injury. The second study, published by Quinn et al in 2014, evaluated 2,663 patients treated for traumatic lacerations in three U.S. EDs.<sup>8</sup> The researchers found no statistically significant difference in the rate of infection for wounds closed before or after 12 hours from injury. In a multivariate model for predictors of infection, they found that diabetes (odds ratio [OR], 3.1; 95% confidence interval [CI], 1.2-8.0), location not in the head or neck (OR, 2.5; 95% CI, 1.4-4.5), length > 5 cm (OR, 2.4; 95% CI, 1.4-4.0), and moderate to heavy contamination (OR, 1.9; 95% CI, 1.04-3.3) were independent risk factors for infection, not length of time from injury to closure.

Based on published studies, there is no “golden hour” for wound repair.<sup>9</sup> Studies indicate scalp and face wounds have no time limit, provided there are no signs of infection. And, although the study numbers were small, the risk of infection with closure up to 19 hours after injury is essentially no different from early repair if the wound is not clinically infected.<sup>8</sup>

#### Exam

As with any trauma patient, the airway, breathing, and circulation must be assessed and stabilized before moving on to other concerns. Once the patient is stable, a careful neurovascular exam is necessary prior to anesthesia administration. A wound may be much more severe than it initially appears. Many injuries

**Table 1. Maximal Local Anesthetic Infiltrative Dosing**

Anesthetic	Without Epinephrine	With Epinephrine	Maximum Dose (80 kg Adult)
Bupivacaine	2 mg/kg	3 mg/kg	160 / 240 mg
Lidocaine	4.5 mg/kg	7 mg/kg	360 / 560 mg
Procaine	7 mg/kg	9 mg/kg	560 / 720 mg

can happen quickly, and the patient may not realize the extent of injury. Injuries also often occur in the setting of a heightened emotional state, and this can mask pain. As such, wound size and depth should be evaluated with this in mind. Simple and gentle exploration often will reveal the extent of injury as well as any foreign material. Likewise, any deficit in sensation or function can be used to pinpoint the extent of neurological injury.<sup>10</sup> At this time, wounds should be evaluated for distal neurovascular and/or tendon injury. Anesthesia usually is required for full wound exploration, as well as to facilitate wound cleansing, as these procedures can be extremely painful.

### Diagnostics

Laboratory testing is rarely necessary, but may be considered if there is concern for infection, hemorrhage leading to anemia, or if the patient is on anticoagulants. Plain films and other imaging may be useful to rule out radiopaque foreign body and/or fracture that otherwise may remain undetectable.<sup>11</sup> Bedside ultrasound may be a useful adjunct to evaluate for non-radiopaque foreign body. CT also can be useful in determining the extent of injury in traumatic wounds if there is concern for underlying organ damage.

## Treatment

### Topical Anesthesia

Topical anesthesia has been used mainly in pediatric wounds; however, it has gained momentum for adults in recent years either alone or as pretreatment before local infiltration. Lidocaine, epinephrine, tetracaine (LET) and eutectic mixture of local

anesthetics (EMLA) are efficacious in reducing the pain of local injection. EMLA cream as pretreatment before infiltrative local anesthesia increases patient satisfaction with treatment.<sup>13,14,15</sup> A Cochrane review published in 2011 identified that most data-generating studies regarding topical anesthesia efficacy were at medium to high risk of bias, and concluded that low-risk bias studies still are needed.<sup>16</sup>

If considering topical anesthesia, keep in mind that there will be an increased length of stay, as the onset typically is slower than local infiltration. Topical anesthetics usually require 10 to 30 minutes for maximum effect. There has been some consideration of the possibility of placing these medications on wounds upon arrival to ED triage for efficiency.<sup>17,18</sup> A randomized, double-blind trial exploring the ability of triage nurses to pretreat lacerations with topical anesthetics showed that the topical anesthetics significantly anesthetized the tissue to decrease pain of local injection by the time of physician evaluation.<sup>19</sup>

A recent study investigating cryotherapy for relief of local injection pain found that application of a cold compress to the wound site for just five minutes was enough to decrease pain scores significantly.<sup>20</sup> Other novel methods for local anesthesia may be available in the near future. A recent noninferiority trial compared the effectiveness of local infiltration with a topical anesthetic putty application, comparing pain scores with outcomes.<sup>21</sup> The lidocaine putty was found to be noninferior to an infiltrative technique. Lidocaine putty had

the added benefit of not disrupting or distorting the tissue when applied, unlike with local anesthetic infiltration.

### Infiltrative Anesthesia

With or without pretreatment, intradermal anesthesia is a quick and effective method to control pain before irrigation, exploration, and closure of the wound. Lidocaine 1% is the most commonly used local anesthetic with broad applications for approaching many types of pain, not only dermal but also visceral and central pain, renal colic, and more.<sup>22</sup> Other local anesthetics, such as procaine 1% and the longer-acting bupivacaine 0.25%, also may be used. (See Table 1.)

The addition of epinephrine 1% has the benefit of vasoconstriction, which allows the provider to have a clearer field with less blood loss. The addition of epinephrine traditionally has been discouraged if vascular compromise was suspected, although there is no evidence of harmful effect, even if used on the fingers, toes, or nose.<sup>23</sup> Epinephrine also allows higher total doses of anesthetic to be used, increasing the maximum dose of lidocaine from 4.5 mg/kg of body weight to 7.0 mg/kg of body weight. This raises the maximal total dose of lidocaine 1% from 36 mL to 56 mL in an 80-kg adult, which is particularly useful when dealing with larger wounds.

The injection of local anesthetics can be very painful. This can be mitigated by injecting slowly, warming the medication to body temperature, using small (27-30 gauge) needles, avoiding large injections in one location, and using distracting vibratory stimulation in the area. Another option is to add 1 mL sodium bicarbonate 8.4% for every 9 mL of 1% lidocaine, which not only decreases pain but also shortens time to anesthesia onset.

### Nerve Blocks

Regional nerve blocks can be extremely useful in the ED. They have the dual benefits of anesthetizing a large area with minimal time,

effort, and pain to the patient while also requiring smaller volumes of local anesthetic medication. As the anesthetic is not injected directly into the wound area, it does not distort the tissue, allowing for alignment during repair and perhaps a better cosmetic outcome. Nerve blocks also may be considered when anesthetizing areas that are large or difficult to access, such as the face or fingers. This practice can lower patient pain scores by diverting the injection site from the inflamed local area of the wound and requiring fewer injections in total.<sup>24</sup>

While a full analysis of nerve blocks is outside the scope of this review, it is worth noting that a 2014 review of hand anesthesia revealed new best practice for performing digital anesthesia.<sup>25</sup> Evidence from multiple studies supports the use of single volar (palmar) injection in the middle of the proximal phalanx over traditional dual dorsal web space injection. The pain of injection was found to be no different than in the dorsal webbing of the finger and requires only one injection instead of two.

### Personal Protective Equipment

Sterile gloves may be used if preferred for closer fit and more flexibility; however, studies have shown no decrease in infection rate when compared with non-sterile exam gloves.<sup>24</sup>

### Irrigation and Cleaning

Normal saline has long been used for irrigation of wounds, but recent studies from Stanford University have shown that tap water is safe and effective in wound irrigation.<sup>26</sup> In fact, fewer wound infections actually were observed in the tap water group compared to the sterile saline group, although this finding was just shy of statistical significance. Tap water has the added benefit of easily allowing for copious irrigation without the expense of sterile fluids. However, this does come under the assumption that potable water is readily available. When this is not the case, sterile normal saline should be used.<sup>27,12</sup>

**Table 2. Superficial Laceration Suture Choice**

Size	Location	Duration
3-0 to 4-0	Sole of foot	7-10 days
4-0	Joints	10-14 days
4-0 to 5-0	Trunk/extremities/hands	7-10 days
5-0	Fingers/toes	12-14 days
5-0 to 6-0	Face/eyebrow	3-5 days

Perhaps counterintuitively, antibiotic solution use for washout actually has been shown to increase wound healing complications. One prospective, randomized study found a statistically significant increase in wound healing failures and dehiscence in the antibiotic group when compared to the group using non-sterile soap for irrigation.<sup>27</sup>

Low-pressure irrigation, defined as approximately 0.5 psi or lower, is achieved with a slow, gentle wash. Such pressure is sufficient for cleaning simple, non-bite, uncontaminated wounds in a highly vascular area, such as the scalp or face. High-pressure irrigation, defined as approximately 7 psi, can be obtained with forceful depression on the piston of a 30 mL or larger syringe through a 19-gauge catheter or needle, or with a commercially available splashguard. High-pressure irrigation is regarded as more effective for removal of debris and reduction in post-repair wound infection.

Many lacerations have little visible contamination, and there is little evidence regarding the amount of irrigation required to minimize post-repair infection. Recommendations by wound care experts range from 25 to 100 mL of irrigation fluid per centimeter of laceration.<sup>17</sup>

### Wound Closure

**Wound Adhesive/Adhesive Strips.** Tissue adhesives and adhesive strips have been shown to be noninferior to sutures when used appropriately.<sup>28,29,30</sup> Both options have been well-evaluated over the last 20 years and found to be excellent, less painful or painless


**Table 3. Common Absorbable Suture Options**

Type	Tensile Strength Time
Chromic gut	3-4 weeks
Absorbable synthetic	3 weeks
Fast-absorbing gut	5-7 days
Nonabsorbable options of nylon and polypropylene do not lose tensile strength.	

alternatives to sutures for the closure of appropriate lacerations. Either option can expedite care and improve efficiency, while improving patient experience. Suitable lacerations are those that are under low tension (so as not to risk wound dehiscence), do not require deep-layer sutures, and are not complicated by hair growth. Infection rates and cosmetic results have not been found to be significantly different in wounds closed in this manner when compared with wounds closed with sutures. Whereas tissue adhesive is used with a painting motion directly atop the wound edges, proper use of adhesive strips consists of placing closure devices perpendicular to the wound edge in the same locations where a suture would be placed.<sup>29,31,32</sup>

**Staples.** Staples are a quick, cost-effective, and noninferior technique for wound closure in the ED. One pediatric prospective, randomized trial showed staples resulted in shorter procedures at a lower cost with statistically equal outcomes as sutures.<sup>33</sup>

**Figure 1. Running Sutures**



Top left: placement of running sutures; top right: securing sutures in place using final open loop and suture tail; bottom left: completed sutures; bottom right: one damaged suture can lead to a reopened wound

Staples should not be used when CT or MRI may be needed while they are in place, as they may interfere with image quality.

**Hair Apposition.** For scalp laceration closure, the alternative method of hair apposition may be used.<sup>34,35</sup> Hair at least 1 cm long may be twisted together and secured with tissue adhesive to achieve wound closure. This method actually has been shown by retrospective observational study to be superior to scalp sutures for wounds that were less than 10 cm with hair greater than 3 cm in length.


**Secondary or Delayed Primary Repair.** Healing by secondary intention is considered when there is concern for infection (e.g., bite wounds or highly contaminated wounds) in non-cosmetically important locations. Alternatively, wounds at high risk also can be considered for delayed primary closure after four days of open wound management: cleaning, dressing, and with or without prophylactic oral antibiotics. The risk of infection rises as time passes after the injury, and

closure after four days of such care can be done with an acceptable risk for wound infection.

**Sutures.** The major choice for sutures is between absorbable and nonabsorbable ones. Absorbable sutures are convenient for patients in that they do not require a follow-up visit for removal (*see Tables 2 and 3*); however, their use has been discouraged in visible and cosmetically important areas because of the belief that the enzymatic process responsible for suture dissolution will leave a visible mark. Newer research has shown no increase in adverse appearance between the two types of sutures when used properly, and that cosmetic outcomes for absorbable and nonabsorbable options generally are equivalent.<sup>36,37</sup>

It is important to consider the amount of time that the sutures will last, compared with how much time is needed for the wound to heal.<sup>36</sup> For example, an absorbable synthetic suture (such as Vicryl™) will last approximately 21 days, so it would

**Figure 2. Vertical Mattress Sutures**




Top: vertical mattress suture placement; middle: demonstrating the deep and superficial elements of vertical mattress sutures; bottom: secured suture

be unnecessary for a wound that is expected to heal in less than a week. Likewise, an absorbable suture that will dissolve in five days would not be appropriate for use over joints or on the extremities, which typically take longer to heal to tolerate the stress of movement. When choosing a suture, consider monofilament vs. braided material. Although monofilament sutures have less of a local inflammatory effect than braided sutures, braided sutures, by their nature, will maintain knot integrity more readily.

**Techniques.** Simple interrupted sutures are the most versatile type but may be the most time-consuming option. A running or “baseball” stitch is a much faster technique; however, should a single stitch break, the entire repair may fail and lead to wound dehiscence. (*See Figure 1.*) For wounds under high tension, consider either undermining the wound with a deep suture or using mattress sutures.

Vertical mattress sutures have the benefit of placing a deep and

**Figure 3. Placement of Horizontal Mattress Sutures**



Top: horizontal mattress suture placement; bottom: secured suture

superficial suture simultaneously and have an excellent cosmetic effect. (See Figure 2.) In addition, vertical mattress sutures place the knots outside the wound, so there is less concern for the knots fusing into the wounds. Horizontal mattress sutures are particularly useful for edematous, large, and/or high-tension wounds. (See Figure 3.) Another alternative for higher-tension wounds is the locked running suture. This differs from the running suture in that each “throw” is locked by passing the suture through the loop that has been created. (See Figure 4.) Compared to the standard running suture, each loop of the locked running suture will hold more tension.

If the wound includes a pointed edge, consider a corner stitch when there is sufficient tissue on the pointed edge to run the suture parallel to the skin surface. This minimizes tip retraction and scar formation. Finally,

a running subcuticular stitch may be used, but it often is less favored because it is the most technically difficult suture. This technique eliminates any possibility of visible scarring from the thread running into and out of the surface of the skin, as horizontal stitches are placed just below the epidermis and run back and forth. However, if not placed correctly, this technique may distort the appearance of the wound.

## Additional Treatment

### Prophylactic Antibiotics

There is no clear evidence that antibiotic prophylaxis reduces the rate of post-repair infections for the majority of traumatic wounds repaired in the ED. Moreover, many wounds considered at high risk, such as hand lacerations, do not have a lower rate of infection if antibiotics are prescribed. A prospective observational

study found that age, gender, diabetes, wound closure method, and prophylactic antibiotics did not affect wound infection outcomes for simple hand lacerations.<sup>38</sup>


Most non-bite wound infections are due to staphylococci or streptococci. Despite the increase in community-associated methicillin-resistant *Staphylococcus aureus* (MRSA) skin infections, MRSA has not been a common isolate in traumatic wound lacerations. For prophylactic coverage of these non-bite lacerations, a beta-lactam is adequate. For plantar puncture wounds through athletic shoes, there is the theoretical risk for *Pseudomonas* infection, and prophylaxis with ciprofloxacin can be considered; however, local antibiotic resistance patterns must be taken into account.

### Tetanus

According to Centers for Disease Control and Prevention (CDC) guidelines, tetanus antibiotic prophylaxis is not required for clean, minor wounds in a vaccinated individual (three or more doses to build initial immunity, as well as less than 10 years since the last tetanus toxoid). For deep wounds, crush wounds, or any wound contaminated by dirt, soil, or saliva, Td or Tdap should be given if it has been more than five years since the last tetanus toxoid booster. Anyone with a laceration that is “neither clean nor minor,” and who has had fewer than three prior doses of tetanus toxoid or who has an unknown history should receive tetanus immunoglobulin (TG) as well as Td or Tdap to provide immediate immunity and to promote immune response.<sup>40</sup>

Of note, the Academy of Emergency Medicine and Care and the World Society of Emergency Surgery recently published a position paper concerning management of traumatic wounds that considered tetanus prophylaxis.<sup>41</sup> The position paper was based on 10 questions submitted to an expert panel, and two of its eight recommendations on traumatic wounds concerned tetanus and proper administration of the

**Figure 4. Placement of Locked Running Sutures**



toxoid. Expert consensus agreed with the CDC recommendations that every patient who presents with a traumatic wound should be evaluated for immunization status and that tetanus infection is a potential risk in all traumatic wounds.<sup>41</sup> Vaccination status should be documented in all lacerations, and Tdap may be given in the thigh muscle or deltoid as a 0.5 mL dose when needed.<sup>41,42</sup>

### Wound Care

When it comes to outpatient wound care recommendations for patients, traditional teaching has been to place non-adherent dressings over the wound to provide a moist, clean environment for wound healing to take place. While such dressings have not been proven to prevent bacterial contamination, they help contain drainage and minimize dried crust formation.<sup>43</sup>

Patients typically are advised not to get sutured wounds wet. Studies

previously have shown that uncovered wounds that were exposed to water (e.g., washing or showering) after 12 hours had no increase in infection rate.<sup>44</sup> There are no data on immersion restrictions, although it makes sense to avoid immersing the wound in water until the wound has closed and the sutures are removed. Of course, tissue adhesives and adhesive strips cannot get wet, or they will lose their strength.

### Sub-specialty Follow-up

With complicated lacerations or highly visible areas where cosmetic appearance is important, there is often a question of whether a plastic surgery consultation would yield better outcomes. Specialty consultation is not always available, such as when presentation occurs in a remote environment or late at night. Fortunately, a study of facial laceration repair found that patient satisfaction scores were similar between wounds repaired by plastic

surgeons and those repaired by ED physicians.<sup>45</sup> A subset of females and parents of small children tended to be more satisfied with plastic surgeon repair, but this bias is thought to be because of patient awareness of provider specialty. Dermatology referral also may be considered for scar management, as advances with laser surgery and pharmacologic management can assist with scar mitigation once the wound has healed.<sup>46</sup>

### Nail Bed Laceration

Nail bed injuries add another layer of complexity to wound closure. (See *Figure 5*.) Preserving the nail serves more than just a cosmetic outcome. The fingernail allows for increased tactile control, promotes proper circulation, and serves as protection for the sensitive fingertip. At this time, there is some controversy regarding proper repair techniques, but it has long been agreed that an improperly repaired nail bed will lead to permanent deformity.<sup>47</sup>

Classic teaching is that leaving a nail bed laceration open can lead to abnormal regrowth of the fingernail, and, as such, all nail bed lacerations should be repaired with removal of the nail to facilitate primary closure. This was challenged in 1999 by Rose et al in a prospective study comparing outcomes in 52 children with crush injuries involving the nail bed.<sup>48</sup> Patients were divided into two equal groups: one group had nail removal with primary repair and one had simple nail trephination. No significant difference was found between the two groups regardless of subungual hematoma size, mechanism, child age, or underlying fracture.<sup>48</sup>

Often, the nail has been removed because of the actual injury prior to the patient's arrival, and, as such, the decision to remove or keep the nail is already made. However, when there is a choice, the options are to replace the natural nail (when present), stent the matrix with an artificial substance, or leave the nail bed open following primary laceration closure. A 2014 retrospective study of 401 patients revealed

## Figure 5. Nail Bed Laceration



Left: initial injury noting partial avulsion of nail; middle: nail removed intact; right: after nail bed repaired, nail placed back under basal cuticle and on nail bed, then held in place with two lateral distal sutures

Images courtesy of Salim R. Rezaie, MD

far superior outcomes with native nail replacement when compared to the use of silicone nail splints.<sup>49</sup> Native nail replacement yielded significantly fewer nail deformities, whereas silicone nail splints led to twice the infection rate. It is also worth noting, however, that a 2015 randomized feasibility trial from the United Kingdom of 60 children who underwent nail bed repair found increased complication and infection rates among patients with nail replacement when compared to patients whose nails were discarded with the nail bed left open.<sup>50</sup> Because of the small study size, the results were not statistically significant, and the authors noted plans for a definite randomized, controlled trial. More evidence is needed regarding nail splinting vs. discarding after nail bed repair.

If the nail is to be replaced, common practice is to secure the nail with absorbable sutures, as there is less risk of granuloma formation when compared with nonabsorbable sutures. A 2008 prospective trial compared the use of standard suture fixation with the use of tissue adhesive to reattach the nail and found no difference in both physician- and patient-perceived outcomes, pain scores, and subsequent functional ability between the two groups.<sup>51</sup> This

trial also evaluated the use of tissue adhesive against absorbable sutures for primary laceration repair with similar results. The researchers found that the average time required for nail bed repair with tissue adhesive was 9.5 minutes, compared with an average time of 27.8 minutes for suture repair. Thus, tissue adhesive may be considered for both primary nail bed laceration repair and nail replacement.

When there is a laceration through the nail with minimal adjacent damage, the laceration and nail may be re-approximated with the same application of tissue adhesive, without removing the nail. When the nail must be reattached, some providers anecdotally recommend pre-drilling the nail with an 18-gauge needle and exiting sutures through this hole to prevent foreign body granuloma. This also may cause less trauma to the nail bed, as it avoids having to forcibly push the suture needle through the nail. However, no comparison studies are known to have evaluated this method.

## Bite Wounds

### Canine

In 2015, 80-90% of the estimated 3 to 6 million animal bites per year in the United States were from

canines.<sup>52</sup> They occur most often in young children (especially males) between the ages of 5 to 9 years. Knowledge of the proper care of such wounds is extremely important. Often there is the question of whether these wounds should be closed or if this will lead to a significant risk for infection. A randomized, controlled trial evaluated 200 patients over three years for this very reason, looking at full-thickness wounds presenting to the ED within 48 hours of injury.<sup>53</sup> All wounds were cleansed using high-pressure irrigation with iodine antiseptics, and all patients received the same antibiotic treatment of amoxicillin/clavulanic acid, 500/125 mg BID for five days. Outcomes were measured for presence of infection and for cosmetic appearance. No significant difference in infection rates was found between the two groups. However, cosmetic appearance of the wounds was found to be significantly better in the primary closure group. It is important to note that wounds treated later than eight hours from injury demonstrated a 22.2% infection rate vs. 4.5% for those with earlier treatment times.

Regardless of closure, the first-line antibiotic choice is amoxicillin-clavulanate in patients who are not allergic to penicillin. In severe

**Figure 6. Hand Infection Due to Cat Bite**



*Pasteurella multocida* isolated from wound

Image courtesy of J. Stephan Stapczynski, MD

wounds, consider an initial dose of IV ampicillin-sulbactam, ticarcillin-clavulanate, piperacillin-tazobactam, or a carbapenem. If wounds are infected at presentation, a 10-day minimum course is recommended.<sup>52,54</sup> For penicillin-allergic patients, metronidazole plus doxycycline or trimethoprim/sulfamethoxazole or ciprofloxacin may be used.<sup>52</sup>

### Feline

Cat bites are more likely puncture wounds, while dog bites are more likely to tear or crush tissue. Approximately 50% of untreated cat bites will become infected compared to about 8% of dog bites.<sup>55</sup> (See Figure 6.) In a retrospective analysis from 2004, Mitnovetski et al recommended prophylactic antibiotics for small cat bite wounds, avoidance of primary closure (with evaluation at 24 and 48 hours for delayed closure), and drainage and debridement with irrigation for deep space infection.<sup>56</sup> Of note, most patients already had signs of infection — erythema and swelling — when seen in the ED. Puncture wounds from felines should

not be closed, but should be allowed to heal by secondary intention.<sup>57</sup>

Antibiotics must include coverage for *Pasteurella multocida*. First-line antibiotic coverage consists of amoxicillin-clavulanate (500/125 mg TID or 875/125 BID) in combination with dicloxacillin (500 mg QID) for *S. aureus* coverage.<sup>52</sup> A retrospective review from 2013 also suggested hospitalization be considered for bites affecting tendon sheaths and joints, or when erythema, pain, and swelling are present.<sup>59</sup> These infections can spread along fascial planes quite rapidly and ultimately may require surgical wash-out and/or debridement.

### Human

As with cat bites, human bites should be irrigated and debrided as needed. Although human oral flora can have up to 190 bacteria species, most human bite infections are caused by *Eikenella corrodens* or *Streptococcus* sp.<sup>24</sup> Antibiotic prophylaxis after human bites includes amoxicillin/clavulanate. Thereafter, human bites may be closed by primary closures as needed.

### Snake/Scorpion

There is no increase in wound infection or specific pathogen associated with snake bites or scorpion stings, so prophylactic antibiotics for these bites/stings have no benefit. However, these wounds may require antivenom.<sup>52</sup>

### Rabies

When caring for bite wounds, give rabies prophylaxis when there is any suspicion of rabies exposure. Although there are now five cases of survival from human rabies documented using the Milwaukee protocol, treatment with vaccine before the onset of symptoms is a more efficacious and cost-effective rabies solution.<sup>59,60</sup>

### Wound Follow-up

#### Hypertrophic Scars and Keloids

Hypertrophic scars and keloids can be misidentified. Hypertrophic scars are hard, red or pink, raised scars, elevated but remaining within the limits of the original wound, usually emerging within the first month after injury, and may regress over time. Keloids are raised, reddish-purple, nodular scars, harder than hypertrophic scars, invading adjacent tissue extending beyond the margins of the original wound, and rarely regressing over time. Keloid formation is a fairly common occurrence, particularly in patients with African-American, Asian, and Hispanic heritage. The actual pathogenesis is as yet unclear. While there is evidence that hypertrophic scar formation can be minimized by careful wound closure, there is no evidence that keloid formation can be prevented.<sup>61</sup>

#### Suture Removal

A 2012 prospective single-blind trial found that most patients are willing to remove and are capable of removing their own sutures from simple lacerations, provided they receive sufficient removal instructions.<sup>62</sup> Complications were not significantly higher than among patients

who presented to the ED for suture removal. However, high-risk wounds that should be re-evaluated within 48 hours by a physician do not fall into this category.

## References

- Childs DR, Murthy AS. Overview of wound healing and management. *Surg Clin North Am* 2017;97:189-207.
- Simon BC, Hern HG. Wound Management Principles. In: *Rosen's Emergency Medicine — Concepts and Clinical Practice*. 8th ed., Philadelphia: Elsevier; 2014:751-766.
- Bourne CL, Jenkins MA, Brewer KL. Estimation of laceration length by emergency department personnel. *West J Emerg Med* 2014;15:889-891.
- Initial Assessment and Management. In: *Advanced Trauma Life Support Student Course Manual*. 9th ed., Chicago: American College of Surgeons; 2012:6-12.
- Hamer ML, Robson MC, Krizek TJ, Southwick WO. Quantitative bacterial analysis of comparative wound irrigations. *Ann Surg* 1975;181:819-822.
- Waseem M, Lakdawala V, Patel R, et al. Is there a relationship between wound infections and laceration closure times? *Int J Emerg Med* 2012;5:32.
- van den Baar MT, van der Palen J, Vroon MI, et al. Is time to closure a factor in the occurrence of infection in traumatic wounds? A prospective cohort study in a Dutch level 1 trauma centre. *Emerg Med J* 2010;27:540-543.
- Quinn JV, Polevoi SK, Kohn MA. Traumatic lacerations: What are the risks for infection and has the 'golden period' of laceration care disappeared? *Emerg Med J* 2014;31:96-100.
- Peterson AS. The "golden period" for wound repair. *The Journal of Lancaster General Hospital* 2010;5:134-135.
- Casal D, Cunha T, Pais D, et al. A stab wound to the axilla illustrating the importance of brachial plexus anatomy in an emergency context: A case report. *J Med Case Rep* 2017;11:6.
- Fowler TR, Crellin SJ, Greenberg MR. Detecting foreign bodies in a head laceration. *Case Rep Emerg Med* 2015;2015:801676.
- Worster B, Zawora MQ, Hsieh C. Common questions about wound care. *Am Fam Physician* 2015;91:86-92.
- Singer AJ, Stark MJ. LET versus EMLA for pretreating lacerations: A randomized trial. *Acad Emerg Med* 2001;8:223-230.
- Park SW, Oh TS, Choi JW, et al. Topical EMLA cream as a pretreatment for facial lacerations. *Arch Plast Surg* 2015;42:28-33.
- Ernst AA, Marvez-Valls E, Nick TG, et al. Topical lidocaine adrenaline tetracaine (LAT gel) versus injectable buffered lidocaine for local anesthesia in laceration repair. *West J Med* 1997;167:79-81.
- Eidelman A, Weiss JM, Baldwin CL, et al. Topical anaesthetics for repair of dermal laceration. *Cochrane Database Syst Rev* 2011;(6):CD005364.
- Nicks BA, Ayello EA, Woo K, et al. Acute wound management: Revisiting the approach to assessment, irrigation, and closure considerations. *Int J Emerg Med* 2010;3:399-407.
- Ali S, McGrath T, Drendel AL. An evidence-based approach to minimizing acute procedural pain in the emergency department and beyond. *Pediatr Emerg Care* 2016;32:36-42; quiz 43-44.
- Singer AJ, Stark MJ. Pretreatment of lacerations with lidocaine, epinephrine, and tetracaine at triage: A randomized double-blind trial. *Acad Emerg Med* 2000;7:751-756.
- Mahshidfar B, Cheraghi Shevi S, Abbasi M, et al. Ice reduces needle-stick pain associated with local anesthetic injection. *Anesth Pain Med* 2016;6:e38293.
- Jenkins MG, Murphy DJ, Little C, et al. A non-inferiority randomized controlled trial comparing the clinical effectiveness of anesthesia obtained by application of a novel topical anesthetic putty with the infiltration of lidocaine for the treatment of lacerations in the emergency department. *Ann Emerg Med* 2014;63:704-710.
- Golzari SE, Soleimanpour H, Mahmoodpoor A, et al. Lidocaine and pain management in the emergency department: A review article. *Anesth Pain Med* 2014;4:e15444.
- Ilicki J. Safety of epinephrine in digital nerve blocks: A literature review. *J Emerg Med* 2015;49:799-809.
- Sanders JE. Pediatric wound care and management in the emergency department. *Pediatr Emerg Med Pract* 2017;14:1-24.
- Al Youha S, Lalonde DH. Update/review: Changing of use of local anesthesia in the hand. *Plast Reconstr Surg Glob Open* 2014;2:e150.
- Weiss EA, Oldham G, Lin M, et al. Water is a safe and effective alternative to sterile normal saline for wound irrigation prior to suturing: A prospective, double-blind, randomised, controlled clinical trial. *BMJ Open* 2013;3(1): pii: e001504. doi: 10.1136/bmjopen-2012-001504.
- Anglen JO. Comparison of soap and antibiotic solutions for irrigation of lower-limb open fracture wounds. A prospective, randomized study. *J Bone Joint Surg Am* 2005;87:1415-1422.
- Mattick A, Clegg G, Beattie T, Ahmad T. A randomised, controlled trial comparing a tissue adhesive (2-octylcyanoacrylate) with adhesive strips (SteriStrips) for paediatric laceration repair. *Emerg Med J* 2002;19:405-407.
- Wilbur L, Seupaul R. Evidence-based emergency medicine. Are tissue adhesives an acceptable alternative for simple lacerations? *Ann Emerg Med* 2011;58:373-374.
- Singer AJ, Hollander JE, Valentine SM, et al. Prospective, randomized, controlled trial of tissue adhesive (2-octylcyanoacrylate) vs standard wound closure techniques for laceration repair. Stony Brook Octylcyanoacrylate Study Group. *Acad Emerg Med* 1998;5:94-99.
- Bruns TB, Worthington JM. Using tissue adhesive for wound repair: A practical guide to dermabond. *Am Fam Physician* 2000;61:1383-1388.
- Forsch RT, Little SH, Williams C. Laceration repair: A practical approach. *Am Fam Physician* 2017;95:628-636.
- Khan AN, Dayan PS, Miller S, et al. Cosmetic outcome of scalp wound closure with staples in the pediatric emergency department: A prospective, randomized trial. *Pediatr Emerg Care* 2002;18:171-173.
- Ozturk D, Sonmez BM, Altinbilek E, et al. A retrospective observational study comparing hair apposition technique, suturing and stapling for scalp lacerations. *World J Emerg Surg* 2013;8:27.
- Forsch RT. Essentials of skin laceration repair. *Am Fam Physician* 2008;78:945-951.
- Tejani C, Sivitz AB, Rosen MD, et al. A comparison of cosmetic outcomes of lacerations on the extremities and trunk using absorbable versus nonabsorbable sutures. *Acad Emerg Med* 2014;21:637-643.
- Al-Abdullah T, Plint AC, Fergusson D. Absorbable versus nonabsorbable sutures in the management of traumatic lacerations and surgical wounds: A meta-analysis. *Pediatr Emerg Care* 2007;23:339-344.
- Roodsari GS, Zahedi F, Zehetabchi S. The risk of wound infection after simple hand laceration. *World J Emerg Med* 2015;6:44-47.
- Lee JH, Jeon MS, Lee DL, et al. Analysis of patients with facial lacerations repaired in the emergency room of a provincial hospital. *Arch Plast Surg* 2015;42:34-39.
- Mootrey G, Tiwari T, Weinbaum C, eds. Tetanus. In: Centers for Disease Control and Prevention. *Epidemiology and Prevention of Vaccine-Preventable Diseases*. Hamborsky J, Kroger A, Wolfe S, eds. 13th ed. Washington D.C. Public Health Foundation, 2015. Available at: <https://www.cdc.gov/vaccines/pubs/pink-book/tetanus.html>. 2015:341-352.
- Prevaldi C, Paolillo C, Locatelli C, et al. Management of traumatic wounds in the emergency department: Position paper from the Academy of Emergency Medicine and Care (AcEMC) and the World Society of Emergency Surgery (WSES). *World J Emerg Surg* 2016;11:30.
- Wolicki JE, Weinbaum C, Weaver D. Vaccine administration. In: Centers for Disease Control and Prevention. *Epidemiology and Prevention of Vaccine-Preventable Diseases*. Hamborsky J, Kroger A, Wolfe S, eds. 13th ed. Available at: <https://www.cdc.gov/vaccines/pubs/pinkbook/vac-admin.html>. 2015:79-106.
- Chrintz H, Vibits H, Cordtz TO, et al. Need for surgical wound dressing. *Br J Surg* 1989;76:204-205.
- Heal C, Buettner P, Raasch B, et al. Can sutures get wet? Prospective randomised controlled trial of wound management in general practice. *BMJ* 2006;332:1053-1056.
- Lee SJ, Cho YD, Park SJ, et al. Satisfaction with facial laceration repair by provider specialty in the emergency department. *Clin Exp Emerg Med* 2015;2:179-183.
- Miletta NR, Donelan MB, Hivnor CM. Management of trauma and burn scars: The dermatologist's role in expanding patient access to care. *Cutis* 2017;100:18-20.
- Ashbell TS, Kleinert HE, Putcha SM, Kutz JE. The deformed finger nail, a frequent result of failure to repair nail bed injuries. *J Trauma* 1967;7:177-190.

## Help Us Help You

Share your expert opinion and help us tailor future articles to meet your professional needs. Please take our reader survey at <https://www.surveymonkey.com/r/3RLMRQP> and tell us which topics intrigue you most.

48. Roser SE, Gellman H. Comparison of nail bed repair versus nail trephination for subungual hematomas in children. *J Hand Surg Am* 1999;24:1166-1170.
49. Weinand C, Demir E, Lefering R, et al. A comparison of complications in 400 patients after native nail versus silicone nail splints for fingernail splinting after injuries. *World J Surg* 2014;38:2574-2579.
50. Greig A, Gardiner MD, Sierakowski A, et al; NINJA Pilot Collaborative. Randomized feasibility trial of replacing or discarding the nail plate after nail-bed repair in children. *Br J Surg* 2017;104:1634-1639.
51. Strauss EJ, Weil WM, Jordan C, Paksima N. A prospective, randomized, controlled trial of 2-octylcyanoacrylate versus suture repair for nail bed injuries. *J Hand Surg* 2008;33:250-253.
52. Aziz H, Rhee P, Pandit V, et al. The current concepts in management of animal (dog, cat, snake, scorpion) and human bite wounds. *J Trauma Acute Care Surg* 2015;78:641-648.
53. Paschos NK, Makris EA, Gantsos A, Georgoulis AD. Primary closure versus non-closure of dog bite wounds. A randomised controlled trial. *Injury* 2014;45:237-240.
54. Cavalcanti AL, Porto E, Dos Santos BF, et al. Facial dog bite injuries in children: A case report. *Int J Surg Case Rep* 2017;41:57-60.
55. Jaindl M, Oberleitner G, Endler G, et al. Management of bite wounds in children and adults — an analysis of over 5000 cases at a level I trauma centre. *Wien Klin Wochenschr* 2016;128:367-375.
56. Mitnovetski S, Kimble F. Cat bites of the hand. *ANZ J Surg* 2004;74:859-862.
57. Singer AJ, Dagum AB. Current management of acute cutaneous wounds. *N Engl J Med* 2008;359:1037-1046.
58. Babovic N, Cayci C, Carlsen BT. Cat bite infections of the hand: Assessment of morbidity and predictors of severe infection. *J Hand Surg Am* 2014;39:286-290.
59. Willoughby RE Jr, Tieves KS, Hoffman GM, et al. Survival after treatment of rabies with induction of coma. *N Engl J Med* 2005;352:2508-2514.
60. Lampejo T, Bruce M, Teall A, et al. Caring for a patient with rabies: Implications of the Milwaukee protocol for infection control and public health measures. *J Hosp Infect* 2017;96:385-391.
61. Han T, Lin DF, Jiang H. Wound natural healing in treatment of tumor-like hypertrophic scar. *An Bras Dermatol* 2017;92:474-477.
62. Macdonald P, Primiani N, Lund A. Are patients willing to remove, and capable of removing, their own nonabsorbable sutures? *CJEM* 2012;14:218-223.

## CME Questions

1. Wound irrigation should be equal to:
  - a. 0-50 mL/cm laceration.
  - b. 25-100 mL/cm laceration.
  - c. 100-150 mL/cm laceration.
  - d. 150-200 mL/cm laceration.
2. What size suture is recommended over joints?
  - a. 3-0
  - b. 4-0
  - c. 5-0
  - d. 6-0
3. Which closure technique is best for a high-tension laceration?
  - a. Single interrupted
  - b. Running closure
  - c. Staples
  - d. Horizontal mattress
4. Which injury does *not* require a tetanus booster shot?
  - a. A fully immunized child with last dose two years ago who stepped on a metal nail
  - b. A child with two tetanus vaccine doses who got cut on the playground
  - c. A woman immunized six years ago with a laceration caused by heavy steel girder that landed on her leg
  - d. A middle-aged man last immunized in high school who cut himself chopping carrots for his son's lunch
5. Which of these injuries is *improperly* paired to the bacteria commonly associated with it?
  - a. Human "fight bite," *Eikenella corrodens*
  - b. Plantar puncture wound, *Pseudomonas*
  - c. Cat bite, *Pasteurella multocida*
  - d. Snake bite, *Streptococcus pyogenes*

## PRIMARY CARE REPORTS

### CME Objectives

Upon completion of this educational activity, participants should be able to:

- Summarize recent, significant studies related to the practice of primary care medicine;
- Evaluate the credibility of published data and recommendations related to primary care medicine;
- Discuss the advantages and disadvantages of new diagnostic and therapeutic procedures in the primary care setting.

Is there an article or issue you'd like posted to your website? Interested in a custom reprint?

There are numerous opportunities to leverage editorial recognition to benefit your brand.

Call us at (800) 688-2421 or email [Reprints@AHCMedia.com](mailto:Reprints@AHCMedia.com) to learn more.

Discounts are available for group subscriptions, multiple copies, site-licenses, or electronic distribution. For pricing information, please contact our Group Account Managers at:

Phone: (866) 213-0844  
Email: [Groups@AHCMedia.com](mailto:Groups@AHCMedia.com)

To reproduce any part of AHC newsletters for educational purposes, please contact The Copyright Clearance Center for permission:

Email: [info@copyright.com](mailto:info@copyright.com)  
Website: [www.copyright.com](http://www.copyright.com)  
Phone: (978) 750-8400

Access Your Issues  
Online!

Visit [AHCMedia.com](http://AHCMedia.com)  
and go to  
My Account to log in.

## EDITOR IN CHIEF

**Gregory R. Wise, MD, FACP**  
Associate Professor of Medicine  
Oscar Boonshoft School of Medicine  
Wright State University  
Sole Shareholder  
Kettering Physicians Network  
Dayton, OH

## EDITORIAL BOARD

**Nancy J.V. Bohannon, MD, FACP**  
Private Practice  
San Francisco, CA

**Clara L. Carls, DO**  
Program Director  
Hinsdale Family Medicine Residency  
Hinsdale, IL

**Norton J. Greenberger, MD**  
Clinical Professor of Medicine  
Harvard Medical School  
Senior Physician  
Brigham & Women's Hospital  
Boston, MA

**Brian Hocum, PharmD**  
Adjunct Faculty, Washington State University College of Pharmacy,  
Spokane

**Udaya Kabadi, MD**  
Professor  
University of Iowa  
School of Medicine  
Iowa City, IA

**Norman Kaplan, MD**  
Professor of Internal Medicine  
Department of Internal Medicine  
University of Texas Southwestern  
Medical School  
Dallas, TX

**Dan L. Longo, MD, FACP**  
Professor of Medicine  
Harvard Medical School  
Deputy Editor,  
*The New England Journal of Medicine*  
Boston, MA

**David B. Nash, MD, MBA**  
Dean  
Jefferson School of Population Health  
Thomas Jefferson University  
Philadelphia, PA

**Karen J. Nichols, DO, FACOI**  
Dean  
Professor, Internal Medicine  
Midwestern University  
Chicago College of Osteopathic  
Medicine  
Downers Grove, IL

**Allen R. Nissenson, MD**  
Professor of Medicine  
Director of Dialysis Program  
University of California Los Angeles  
School of Medicine

**Kenneth L. Noller, MD**  
Professor and Chairman  
Department of OB/GYN  
Tufts University School of Medicine  
Boston, MA

**Robert W. Piepho, PhD, FCP**  
Professor Emeritus of Pharmacology  
and Toxicology  
Dean Emeritus  
University of Missouri Kansas City  
School of Pharmacy  
Kansas City, MO

**Robert E. Rakel, MD**  
Department of Family and  
Community Medicine  
Baylor College of Medicine  
Houston, TX

**Glen D. Solomon, MD, FACP**  
Professor and Chair  
Department of Internal Medicine  
Wright State University  
Boonshoft School of Medicine  
Dayton, OH

**Leon Speroff, MD**  
Professor of Obstetrics and  
Gynecology  
Oregon Health Sciences University  
School of Medicine  
Portland, OR

**Robert B. Taylor, MD**  
Professor and Chairman  
Department of Family Medicine  
Oregon Health Sciences University  
School of Medicine  
Portland, OR

**Roger D. Woodruff, MD**  
Associate Professor and Chair  
Department of Family Medicine  
Loma Linda University  
Loma Linda, CA

© 2018 by AHC Media, a Relias Learning company. All rights reserved.

**PRIMARY CARE REPORTS™** (ISSN 1040-2497) is published 12 times annually by AHC Media, a Relias Learning company, 111 Corning Road, Suite 250, Cary, NC 27518-9238. Telephone: (800) 688-2421.

**Executive Editor:** Leslie Coplin  
**Executive Editor:** Shelly Mark  
**Editorial Group Manager:** Terrey L. Hatcher  
**Senior Accreditations Officer:** Lee Landenberger

**GST Registration No.:** R128870672

Periodicals Postage Paid at Cary, NC, and additional mailing offices.

**POSTMASTER:** Send address changes to Primary Care Reports, Relias Learning, 111 Corning Road, Suite 250, Cary, NC 27518-9238.

Copyright © 2018 by AHC Media, a Relias Learning company. All rights reserved. Reproduction, distribution, or translation without express written permission is strictly prohibited.

**Back issues: \$26.** Missing issues will be fulfilled by customer service free of charge when contacted within one month of the missing issue's date.

## SUBSCRIBER INFORMATION

### CUSTOMER SERVICE: (800) 688-2421

Customer Service Email Address:  
Customer.Service@AHCMedia.com

Editorial Email Address:  
lcoplin@relias.com

Website:  
AHCMedia.com

### SUBSCRIPTION PRICES

1 year with free AMA  
Category 1/Prescribed credits: \$379

Add \$19.99 for shipping & handling

Online-only, single user price: \$329

### MULTIPLE COPIES:

Discounts are available for group subscriptions, multiple copies, site-licenses, or electronic distribution. For pricing information, please contact our Group Account Managers at Groups@AHCMedia.com or (866) 213-0844.

All prices U.S. only. U.S. possessions and Canada, add \$30 plus applicable GST. Other international orders, add \$30.

## ACCREDITATION

Relias Learning is accredited by the Accreditation Council for Continuing Medical Education (ACCME) to provide continuing medical education for physicians.

Relias Learning designates this enduring material for a maximum of 3 AMA PRA Category 1 Credits™. Physicians should claim only credit commensurate with the extent of their participation in the activity.

This Enduring Material activity, *Primary Care Reports*, has been reviewed and is acceptable for credit by the American Academy of Family Physicians. Term of approval begins 01/01/2018. Term of approval is for one year from this date. Physicians should claim only the credit commensurate with the extent of their participation in the activity. Approved for 3 AAFP Prescribed credits.

The American Osteopathic Association has approved this continuing education activity for up to 2.5 AOA Category 2-B credits.

Successful completion of this CME activity, which includes participation in the evaluation component, enables the participant to earn up to 3 MOC Medical Knowledge points in the American Board of Internal Medicine's (ABIM) Maintenance of Certification (MOC) program. Participants will earn MOC points equivalent to the amount of CME credits claimed for the activity. It is the CME activity provider's responsibility to submit participant completion information to ACCME for the purpose of granting ABIM MOC credit.

This is an educational publication designed to present scientific information and opinion to health professionals, to stimulate thought, and further investigation. It does not provide advice regarding medical diagnosis or treatment for any individual case. It is not intended for use by the layman. Opinions expressed are not necessarily those of this publication. Mention of products or services does not constitute endorsement. Clinical, legal, tax, and other comments are offered for general guidance only; professional counsel should be sought for specific situations.

This CME activity is intended for primary care and family practice physicians. It is in effect for 36 months from the date of the publication.

