

# Trauma Reports

Vol. 6, No. 2

Supplement to *Emergency Medicine Reports, Pediatric Emergency Medicine Reports, ED Management, and Emergency Medicine Alert*

Mar./Apr. 2005

*Burn injuries frequently present to the emergency department (ED). In the majority of cases, the burns are minor, yet, they require a careful assessment, cleaning, dressing, and careful follow-up. In the pediatric and geriatric populations, careful attention, to the history and physical examination, and an awareness of burn patterns associated with abuse, may protect the patient from further inflicted injury.*

*Patients with more severe burn injuries, especially those associated with house fires or explosions, should be assessed carefully for multiple trauma, and care should be taken to protect the spine until injury can be excluded clinically or radiographically. The airway of a burn patient may be particularly challenging; early aggressive intervention, when indicated, may make a potentially disastrous situation manageable. The authors review the diagnosis, classification, and management of patients who have sustained burns.*

— The Editor

## Epidemiology

More than one million burn injuries occur in the United States each year; 700,000 of these injured individuals will seek care in

an ED, and 45,000 will be hospitalized for their injuries.<sup>1</sup> In the United States, \$2 billion are spent on burn care annually.<sup>2</sup>

Seventy percent of burned patients are male, and the average age of patients sustaining burn injury is 30 years. Infants account for 13% of cases, and adults older than 60 years for 11%. The

extremes of age have been associated with an increased risk of death from burns and burn-related injuries.<sup>3</sup>

Burns can occur by several mechanisms. Scald burns arise from exposure to hot liquids or steam. Thermal burns are the result of contact with flames. Contact burns are caused by contact of the skin with hot or cold surfaces. Burns also occur from exposure to radiation, chemicals, and electricity.

Thermal, scald, and contact are the most common categories of burn injuries. In particular age groups, certain types of burns occur more commonly: scald and contact burns are prevalent from birth to 2 years of age, whereas thermal burns are common in the 5- to 20-year range.<sup>3</sup>

Most children dying as a result of burns sustained their injuries in house fires, and as many as 20% of pediatric burns are the result of abuse or neglect.<sup>4</sup> Low income, minority children are

## The Burned Patient: Assessment, Diagnosis, and Management in the ED

**Authors:** **Heidi Teague, MD**, Assistant Professor, Department of Surgery, Division of Emergency Medicine, University of Maryland, Baltimore; **Sharon A. Swencki, MD**, Resident, Department of Surgery, Division of Emergency Medicine, University of Maryland, Baltimore; **Alice Tang, DO, MPH**, Resident, Department of Surgery, Division of Emergency Medicine, University of Maryland, Baltimore.

**Peer Reviewer:** **Carl Menckhoff, MD, FACEP, FAEM**, Assistant Professor and Residency Director, Department of Emergency Medicine, Medical College of Georgia, Augusta

Now available online at [www.ahcpub.com/online.html](http://www.ahcpub.com/online.html) or call (800) 688-2421 for more information.

### EDITOR IN CHIEF

**Ann Dietrich, MD, FAAP, FACEP**  
Associate Clinical Professor  
Ohio State University  
Attending Physician  
Columbus Children's Hospital  
Associate Pediatric Medical Director  
MedFlight  
Columbus, Ohio

### EDITORIAL BOARD

**Mary Jo Bowman, MD**  
Associate Professor of Clinical Pediatrics  
Ohio State University College of Medicine  
Attending Physician, Children's Hospital of Columbus  
Columbus, Ohio

**Larry N. Diebel, MD**  
Associate Professor of Surgery  
Detroit Medical Center  
Wayne State University  
Detroit, Michigan

### Robert Falcone, MD

President  
Grant Medical Center  
Columbus, Ohio

**Theresa Rodier Finerty, RN, BSN, MS, CNA, BC**  
Director, Emergency and Trauma Services,  
OSF Saint Francis Medical Center  
Peoria, Illinois

### Dennis Hanlon, MD

Director  
Emergency Medicine Residency Program  
Assistant Professor of Emergency Medicine  
Allegheny General Hospital  
Pittsburgh, Pennsylvania

### S.V. Mahadevan, MD, FACEP

Assistant Professor of Surgery  
Associate Chief, Division of Emergency Medicine  
Stanford University School of Medicine  
Stanford, California

### Janet A. Neff, RN, MN, CEN

Trauma Program Manager  
Stanford University Medical Center  
Stanford, California

### Ronald M. Perkin, MD, MA, FAAP, FCCM

Professor and Chairman  
Department of Pediatrics  
Brody School of Medicine at East Carolina University  
Medical Director, Children's Hospital University  
Health Systems of Eastern Carolina  
Greenville, North Carolina

### Steven A. Santanello, DO

Medical Director, Trauma Services  
Grant Medical Center  
Columbus, Ohio

### Eric Savitsky, MD

Assistant Professor of Medicine  
Emergency Medicine/Pediatric Emergency Medicine  
UCLA Emergency Medicine Residency Program  
Los Angeles, California

### Perry W. Stafford, MD, FACS, FAAP, FCCM

Head, Pediatric Surgery  
Jersey City Medical Center  
Jersey City, New Jersey.

© 2005 Thomson American Health Consultants  
All rights reserved

three times more likely to die in a house fire than children in high-income categories.<sup>5,6</sup>

Likely causes of burn injury also vary among different ages. Mishaps with flammable liquids are common causes of burn injury in teenagers and young adults,<sup>7</sup> but for the elderly population, kitchen accidents and the resultant exposure to hot liquid or to an open flame are most prevalent.<sup>8</sup>

Burn injury also may occur in connection with industrial incidents as well as other major trauma. One-fifth to one-fourth of severe burns are work related, and 5% to 10% of burned patients sustain multisystem trauma concurrent with their burn injury.<sup>9-11</sup>

A study performed at Massachusetts General Hospital and Shriners Burns Institute in Boston found three distinct risk factors associated with higher mortality: age more than 60 years, burn injury greater than 40% of the total body surface area (TBSA), and the presence of concomitant inhalation injury. If none of these risk factors were present, the death rate associated with burn injury was 0.3%. The presence of one risk factor raised the mortality to 3%. Two risk factors correlated with a 33% mor-

tality, and when all three risk factors were present, the risk of death jumped to 90%. These numbers apply only to adults younger than 90 years. If a burned patient survives the first week after the incident, the chance of death drops to 2.5%. Only 1.1% of burn injured patients surviving after 2 weeks will die from their injuries.<sup>2</sup>

A recent study from the Shriners Burn Institute at the University of Texas sheds some light on pediatric burn mortality numbers. On average, the investigators found that burns of 85% of the TBSA were 30% lethal. Concomitant inhalation injury increases the risk of death. The mortality rate for patients with burns affecting 50% of the TBSA and concomitant inhalation injury was 10%. In comparison, patients with burns covering 73% of the TBSA without inhalation injury had the same 10% mortality. This study also found that males younger than 3 years had a higher risk of death from their injuries than males of other ages. Lastly, Hispanic teenagers had a higher mortality from burn injury than African-American or Caucasian teenagers of the same age.<sup>5</sup>

**Trauma Reports™** (ISSN 1531-1082) is published bimonthly by Thomson American Health Consultants, 3525 Piedmont Road, N.E., Six Piedmont Center, Suite 400, Atlanta, GA 30305. Telephone: (800) 688-2421 or (404) 262-7436.

**Vice President/Group Publisher:** Brenda Mooney

**Editorial Group Head:** Glen Harris

**Managing Editor:** Martha Jo Dendinger

**POSTMASTER:** Send address changes to **Trauma Reports**, P.O. Box 740059, Atlanta, GA 30374. Copyright © 2005 by Thomson American Health Consultants, Atlanta, GA. All rights reserved. Reproduction, distribution, or translation without express written permission is strictly prohibited.

### Accreditation

**Trauma Reports™** continuing education materials are sponsored and supervised by Thomson American Health Consultants. Thomson American Health Consultants designates this continuing education activity for up to 2.5 hours in Category 1 credit toward the AMA Physician's Recognition Award. Each physician should claim only those hours of credit that he/she actually spent in the educational activity. This CME activity was planned and produced in accordance with the ACCME Essentials. Approved by the American College of Emergency Physicians for 2.5 hours of CEP Category 1 credit.

Thomson American Health Consultants (AHC) is accredited by the Accreditation Council for Continuing Medical Education (ACCME) to provide continuing medical education for physicians.

**Trauma Reports®** is approved for approximately 2.5 nursing contact hours. This offering is sponsored by Thomson American Health Consultants, which is accredited as a provider of continuing nursing education by the American Nurses' Credentialing Center's Commission on Accreditation. Provider

**THOMSON**  
★  
**AMERICAN HEALTH  
CONSULTANTS**

### Conflict of Interest Disclosure

In order to reveal any potential bias in this publication, and in accordance with Accreditation Council for Continuing Medical Education guidelines, Dr. Dietrich (editor in chief) reports that she is the medical director for the Ohio Chapter of the American College of Emergency Physicians. Editorial board members Bowman, Diebel, Falcone, Finerty, Hanlon, Mahadevan, Perkin, Santanello, Savitsky, and Stafford report no relationships with companies related to the field of study covered by this CE/CME program. Ms. Neff reports that she is a stockholder in Biopure. Drs. Teague, Swencki, and Tang (authors), and Dr. Menckhoff (reviewer) report no relationships with companies related to the field of study covered by this CE/CME program.

### Subscriber Information

**Customer Service: 1-800-688-2421**

**Customer Service E-Mail:** [customerservice@ahcpub.com](mailto:customerservice@ahcpub.com)

**Editorial E-Mail:** [martha.dendinger@thomson.com](mailto:martha.dendinger@thomson.com)

**World Wide Web page:** <http://www.ahcpub.com>

**FREE** to subscribers of *Emergency Medicine Reports*, *Pediatric Emergency Medicine Reports*, *Emergency Medicine Alert*, and *ED Management*.

approved by the California Board of Registered Nursing, Provider Number CEP 10864, for approximately 2.5 contact hours. This program is approved by the American Association of Critical-Care Nurses (AACN) for 2.5 nursing contact hours. Provider # 10852.

This is an educational publication designed to present scientific information and opinion to health professionals, to stimulate thought, and further investigation. It does not provide advice regarding medical diagnosis or treatment for any individual case. It is not intended for use by the layman. Opinions expressed are not necessarily those of this publication. Mention of products or services does not constitute endorsement. Clinical, legal, tax, and other comments are offered for general guidance only; professional counsel should be sought for specific situations.

This CME/CE activity is intended for emergency, family, osteopathic, trauma, surgical, and general practice physicians and nurses who have contact with trauma patients. It is in effect for 36 months from the date of publication.

### For Customer Service,

Please call our customer service department at **(800) 688-2421**. For editorial questions or comments, please contact **Martha Jo Dendinger**, Managing Editor, at [martha.dendinger@thomson.com](mailto:martha.dendinger@thomson.com).

### Pathophysiology

Burn wounds have three distinct zones of tissue damage: the coagulation zone, the stasis zone, and the hyperemic zone. The coagulation zone consists of the tissue that has been irreversibly destroyed from the primary injury and cannot recover. Surrounding this area is the stasis zone, where damaged tissue with decreased perfusion is located, but the potential for recovery still exists. Adjacent to the stasis zone is the hyperemic zone, the tissue that has sustained minimal damage and will recover spontaneously.<sup>7,12,13</sup> The tissue in the zones of stasis and hyperemia, while still viable, is at risk of destruction from poor perfusion, edema, and inflammation.<sup>13</sup>

Blisters form when damaged capillaries, with increased vascular permeability, leak plasma into the interstitial space. The damaged epidermis separates from the underlying dermis. The fluid of the blister contains inflammatory mediators, such as arachidonic acid metabolites, thromboxane, and calmodulin, as well as plasma proteins and cellular debris. The high osmolarity of this fluid can cause additional water absorption from underlying tissue into the blister, which increases local wound tissue pressure and may cause ischemia in the already compromised tissue.<sup>12</sup>

As the largest organ in the human body, the skin provides protection, immunologic defense, and acts as a barrier to fluid loss. When the skin is burned, these functions are lost, enhancing the victim's risk of systemic illness, sepsis, and multiple organ failure.

Loss of the barrier function of the skin leads to massive fluid losses from evaporation. This fluid loss impedes tissue perfusion and oxygenation. The resultant hypovolemia can result in relative hypoperfusion to distant organs.<sup>7</sup> Injured tissues release inflammatory mediators and vasoactive substances, causing interstitial edema and organ dysfunction. Multi-organ failure usually develops between the second and eighth week after injury and accounts for one-third of burn-related deaths.<sup>2,14</sup> Increased age, increased TBSA burned, male sex, and the presence of inhalation

## Figure 1. Inflicted Immersion Burn



**Figure 1.** Photograph shows stocking distribution of an inflicted immersion burn. Note the clear demarcation between burned and spared skin. The child told medical staff of being forced to stand in hot water for several minutes.

*Reprinted with permission from Bechtel, K. Identifying the subtle signs of pediatric physical abuse. Ped Emerg Med Rep 2001;6:61.*

injury all are associated with an increased likelihood of the development of multi-organ failure after burn injury.<sup>14,15</sup>

Loss of the skin's immunologic defenses leads to an increased susceptibility to infection. Systemic infection results from invasion of bacteria into the body through the burn eschar. Immediately following injury, the burned area is sterile, but then bacteria quickly colonize the wound. Rapidly reproducing in this avascular environment, the bacteria are able to gain access to the rest of the body.<sup>8</sup> Sepsis is common in burned patients and has been noted to precede the development of multi-organ failure. Advanced age and the presence of full-thickness burns are risk factors for the development of severe sepsis.<sup>15</sup>

In the first 1 to 3 hours after burn injury, edema develops and may increase up to 24 hours after injury.<sup>16-18</sup> The development of edema in the setting of burn injury is multifactorial. Vasodilation and increased transcapillary pressure in conjunction with increased extravascular osmotic activity of the burned tissue, increased microvascular permeability, and impaired cell membrane function with swelling of the cells all contribute to the development of edema.<sup>19,20</sup>

**Cardiac Effects.** Immediately after injury, myocardial function may be depressed; however, this typically improves within three days. This depression in myocardial function may be caused by circulating myocardial depressants or persisting hypovolemia, despite aggressive fluid therapy and the lack of classic signs of hypoperfusion.<sup>21</sup> A hyperdynamic cardiovascular response then occurs, with an up-to-twofold increase in cardiac output.<sup>8,22</sup>

**Metabolic Effects.** Metabolic derangements (e.g., metabolic acidosis, respiratory alkalosis, and electrolyte disturbances) are common in patients with burn injury. Intracellular concentrations

of sodium and calcium rise, while intravascular levels of potassium increase as the result of cell membrane alteration.<sup>23</sup>

In comparison with other critically ill patients, patients suffering from burn injuries have the highest metabolic rate.<sup>24</sup> The burned patient has increased energy expenditure, accelerated glycogen and protein breakdown, and lipolysis. This hypermetabolic state is the result of increased circulating catabolic hormones, catecholamines, cortisol, and glucagon.<sup>22,24,25</sup>

Catabolism begins by five days after injury.<sup>24,26</sup> Although it was once thought that catabolism resolved with wound closure, catabolism actually continues up to nine months after the initial injury.<sup>27</sup> The level of catabolism is increased with age, weight, and delay in surgical treatment. Catabolism intensifies with increases in TBSA burned, up to a TBSA of 40%. Sepsis, hyperglycemia, and decreased ambient air temperature also increase catabolism in these critically ill patients.<sup>24,26,28</sup>

Thermoregulation is altered after burn injury. In addition to the loss of the skin's protective function, which results in loss of body heat and hypothermia, the hypothalamic temperature regulation set point increases by 2° C above normal body temperature.<sup>24,29</sup>

The metabolic effects of burn injury can have serious sequelae for the patient. Patients experience loss of lean body mass and body weight, delayed wound healing, and immune depression as a result of their hypermetabolism. Fatty infiltration of the liver develops as a result of the increase in lipolysis.<sup>30</sup> Pediatric patients with severe burns have been found to have a delay in linear growth for two years after injury.<sup>31</sup>

## Clinical Features


**History.** A thorough history is required from every burned patient and may require information from paramedics, family, or witnesses to the injury. Standard elements, such as medical history, surgical history, medications, allergies, and last tetanus immunization, are essential. Beyond those, the mechanism of injury is perhaps the most important information. Knowing how the burn was incurred will help direct the workup and physical examination and help delineate whether the burn injury occurred intentionally. The mechanism of burn injury also may indicate whether the inhalation of toxic gases, such as carbon monoxide or cyanide, may have occurred. Current use of alcohol or illicit drugs also is important to ascertain; it may have contributed to an alteration in mental status, the mechanism of injury, or comorbidities. In pediatric patients, it is essential to ascertain the circumstances surrounding the injury, maintaining a high index of suspicion for intentional acts.

**Physical Examination.** The physical examination of the burned patient begins with assessment of the ABCs (airway, breathing, circulation). After management of these crucial elements, a secondary survey focusing on recognizing concurrent traumatic injuries, if there is associated trauma, should be completed. Evaluate the patient's face and oropharynx for carbonaceous sputum, circumoral burns, and singed nasal hair, which could indicate the presence of an inhalation injury. During this survey, patterns of injury in pediatric patients that may indicate abuse, such as a stocking or glove-like appearance with sharp

**Figure 2. Rule of Nines (Adult)**

**ADULT BODY PART % OF TOTAL BSA**

Arm	9%
Head	9%
Perineum	1%
Leg	18%
Anterior trunk	18%
Posterior trunk	18%



**Figure 2.** The Rule of Nines is designed to rapidly access the percentage of TBSA for a burn injury.

*Illustration courtesy of the authors.*

margins, should be recognized. (See Figure 1.) The severity of the burn injury also should be appraised.

**Severity of Burn Injury.** To determine the severity of a burn injury, assess both the TBSA burned and the depth of the burn injury. The TBSA measurement is used to estimate fluid resuscitation requirements and to assess the risk of death. Burn depth is used to assess the burned patient's need for hospitalization, the need for surgical intervention, as well as the probability of scar development after the wound heals.<sup>32</sup>


There are three methods of estimating the total body surface area burned (First-degree or superficial burns are excluded from BSA calculations). The Rule of Nines is the most commonly used system to estimate the extent of burn injury. It is much more accurate for adults than for children. The body is divided into areas that represent 9% of body surface area or multiples of 9%. The exception is the perineum, which is assigned the value of 1%. (See Figure 2.) For the pediatric patient, the Rule of Nines is altered by taking 4% from each leg and 1% from the perineum to add an additional 9% to the surface area of the head. (See Figure 3.) The values associated with affected areas are summed to estimate the TBSA burned.<sup>3,33</sup>

The Lund and Browder chart is considered a more accurate estimation tool to determine TBSA, especially for pediatric patients. This chart also divides the body into areas and assigns a

**Figure 3. Rule of Nines (Child)**

**CHILD BODY PART % OF TOTAL BSA**

Arm	9%
Head and neck	18%
Leg	14%
Anterior trunk	18%
Posterior trunk	18%



**Figure 3.** Modification to the adult Rule of Nines to reflect the different proportions in the pediatric population.

*Illustration courtesy of the authors.*


percentage body surface area based on the patient's age. This chart accounts for the differences in proportionality between newborns, children, and adults.<sup>3,34</sup> (See Figure 4.)

Another frequently used technique for estimation of injury uses the surface area of the patient's palm, considered to represent 1% of the TBSA. This method is best used for patients with scattered small burns and is believed to be the least accurate of the three methods. Investigators at the University College London Medical School called this method into question. Their research found that the palmar surface of the hand represented only 0.4% body surface area of adults and 0.45% body surface area in children.<sup>35</sup>

Given the increasing incidence of obesity in our society, it is important to consider how estimation of TBSA in the obese patient differs from that process in nonobese patients. Underestimation of burned area on the trunk and legs becomes more common with increasing obesity. The trunk may constitute up to 50% of TBSA in the obese patient, while each leg may account for 20%. The head and arms of the obese patient account for a smaller body surface area than that assigned to them by the Rule of Nines.<sup>36</sup>

The depth of burn injury commonly has been referred to in terms of first-, second-, and third-degree burns. First-degree burns affect only the epidermis. They are warm and red, have no blisters, and generally are painful. In second-degree burns, both the dermis and epidermis are involved. Most burns of this type will be painful, red, and blistered, with moist bases. In third-degree burns, the entire dermis and epidermis are destroyed.

**Figure 4. Lund and Browder Chart**



**Figure 4.** The Lund and Browder Chart is the most accurate estimation tool to determine TBSA and accounts for differences in proportions between children and adults.

*Illustration courtesy of the authors.*

burns are extremely painful. The burned area is erythematous, and the surface blanches readily, with brisk capillary refill. Blisters develop rapidly. Owing to the extensive vasculature of the epidermis, these are moist wounds and produce moderate edema.

Extending to the deeper layers of the dermis, deep partial-thickness burns may have a red and white waxy or mottled appearance. Although these wounds continue to blanch, capillary refill may be slow or entirely absent. Blisters usually are not present. The surface of the wound is moist, notable edema is present, and sensation is altered.<sup>39</sup> Most partial-thickness burns (both superficial and deep) heal spontaneously within 14 days. The extent of scarring resulting from these burns depends upon the depth of the burn. If located over a joint, these burns may require skin grafting.<sup>38</sup>

Full-thickness burns destroy epidermis and dermis and extend into the subcutaneous tissue. They have a white or charred appearance without any blistering. These burns are insensate secondary to destruction of sensory nerves; however, the area of a full-thickness burn will be bordered by an area of less severely burned tissue, which is painful. Subdermal burns extend into muscle, fascia, and bone. Burns of this magnitude require surgical intervention and have an associated risk of systemic disease. Fluid and protein shifts cause intense edema.<sup>38,39</sup> Skin grafting leads to extensive scarring, development of contractures, and impaired mobility.

Customarily, the determination of burn depth is made clinically; however, in serious burn injury, this assessment can be complicated. Wound biopsy has been used to histologically identify burn depth more accurately by examining the tissue for evidence of blocked and patent vasculature.<sup>40</sup> The less invasive technology of laser Doppler imaging (LDI) also is being used to assess microvascular blood flow in the dermis to determine burn depth.

LDI combines laser Doppler technology with scanning techniques to produce an image of tissue perfusion by tracking red blood cell movement.<sup>41,42</sup> This technology has been found to assess burn depth accurately in 97% of injuries compared with 60% to 80% by clinical assessment alone for adult patients.<sup>41</sup> In pediatric burned patients, this technology has been found to have a sensitivity of 90% and specificity of 96% for the detection of deep partial or full-thickness burns when used 36 to 72 hours after injury.<sup>42</sup> This technology has yet to become widespread in

These burns appear leathery and dry and are typically tan in color. Third-degree burns usually are painless, secondary to destruction of pain receptors in the dermis.<sup>37</sup> In some instances, the term “fourth-degree burn” also is used. These are burns that involve underlying tissue such as muscle, bone, or fascia.

However, these categories have been replaced recently by the more accurate and less confusing terms of superficial, partial-thickness burns (which include both superficial and deep partial thickness), and full-thickness burns. This system of categories accounts for the anatomic structures affected by the burn (Table 1).<sup>38</sup>

Superficial burns affect only epidermis. Erythema is present, but no blisters form. The surface is usually dry and painful to touch. In certain areas, such as around the eyes, there also may be edema. These burns usually heal in 3 to 7 days and generally do not lead to scarring.

In superficial partial-thickness burns, injury extends through the epidermis into the superficial layers of the dermis. These

**Table 1. Burn Depth**

CLASSIFICATION	FORMER CLASSIFICATION	STRUCTURES INVOLVED	COLOR	BLISTERS	PAIN
Superficial	First degree	Epidermis	Red, dry	No	Yes
Partial-thickness, superficial	Second degree	Epidermis into superficial dermis	Red, moist	Yes	Yes
Partial-thickness, deep	Second degree	Epidermis into deep dermis	Red/white, waxy	No	Altered
Full-thickness	Third degree	Epidermis, dermis, into subcutaneous tissue	White or charred	No	None
Subdermal	Fourth degree	Epidermis, dermis, into fascia, muscle, bone	White, charred, variable depending on involved structures	No	None

use, but it has the potential to become a useful adjunct in the assessment of burn injury depth in the future.

### Diagnostic Studies

Burned patients should be placed on a cardiac monitor, and pulse oximetry should be assessed and monitored, if indicated. Basic laboratory studies should be obtained in patients with severe burns or concomitant trauma, including a complete blood count, type and crossmatch, chemistries, coagulation profiles, arterial blood gas measurement, and a pregnancy test, when appropriate. All patients with thermal burns should have arterial or venous blood sent for measurement of the carboxyhemoglobin level to evaluate for carbon monoxide poisoning.

An initial chest radiograph is warranted in all burned patients when an inhalation injury is possible. A normal study does not rule out pulmonary injury however, and serial chest radiographs may show delayed development of pulmonary edema or findings of pulmonary contusions. Computed tomography scans should be obtained as indicated in the patient with accompanying traumatic injuries or decreased mental status. In addition, the history and physical examination should guide radiologic examination of the extremities and cervical spine.

### Initial Management

Stabilization of the ABCs is essential in managing any medical emergency. The initial approach to managing severe burn victims is no exception. Because severe burns often are associated with nonthermal injuries, seriously burned patients must be viewed as trauma patients and should be stabilized according to Advanced Trauma Life Support and the American College of Surgeons Committee on Trauma protocol.<sup>8,43</sup>

The first priority in stabilizing these patients is ensuring a patent airway, which can be challenging, secondary to oropharyngeal and laryngeal edema.

Airway edema may progress rapidly in a burned patient who has inhaled heated gases or toxic products of combustion. Signs that indicate the patient may have had a significant inhalational injury include singed nasal hairs, facial burns, oral burns, sooty sputum, and stridor or wheezes. Fiberoptic laryngoscopy or bronchoscopy may be helpful in assessing the degree of airway trauma.

Once the airway is established, it is paramount to secure it; laryngeal edema, even more than oropharyngeal edema, makes intubation difficult in burn patients. If circumferential burns of the chest or an extremity are present, emergent escharotomy may be necessary. If the chest eschar compromises ventilatory motion incisions along the the costal margin, anterior axillary lines and across the top of the chest may be necessary to allow adequate chest movement. Circumferential eschar of an extremity may act as a tourniquet restricting adequate blood flow to an extremity. In this case, an escharotomy should be performed along the lateral aspect of the extremity, through the entire depth of the eschar, to allow the return of adequate blood flow. Obtaining intravenous access also is extremely important for adequate fluid resuscitation. If peripheral access is unobtainable, central access, or intraosseous access in children, generally is required.

### Fluid Resuscitation

Fluid resuscitation in severe burn victims is controversial in many aspects. Despite many years of research in the area of burns, there does not seem to be standardization regarding the type of infusion fluid to use or definitive resuscitation endpoints.

The American Burn Association (ABA) suggests that patients with burns greater than 15% TBSA should undergo fluid replacement according to the Parkland Formula.<sup>3</sup>

$$4 \text{ mL (of IV fluid) } \times \text{ weight (kg) } \times \% \text{ TBSA burn} = \text{Total amount of fluid for the first 24 hours}$$

**Table 2. Consensus Formulas for Fluid Resuscitation of Burned Patients after 24 Hours**

TBSA BURNED (%)	FLUID ADMINISTRATION
0-30	None
30-50	0.3 mL/kg/%burn/24 hr
50-70	0.4 mL/kg/%burn/24 hr
70-100	0.5 mL/kg/% burn/24 hr

**Key:** TBSA = total body surface area.

**Table 3. Resuscitation Endpoints**

• Sensorium	Arousable and comfortable
• Digital temperature	Warm peripherally
• Systolic blood pressure	
- For infants –	60 mmHg systolic
- For older children –	70–90 mmHg + 2 x age in years
- For adults –	mean arterial pressure >60 mmHg
• Pulse	80–180/min (age dependent)
• Urine output	0.5–1 mL/kg/hr (glucose negative)
• Base deficit	<2 mEq/L

*Adapted from Sheridan RL. Comprehensive treatment of burns. Curr Probl Surg 2001;38:641-756.*

This formula currently is the gold standard and applies only to adults. However, the Parkland formula is for replacement fluid administration and does not include approximation of maintenance fluids. Pediatric considerations will be discussed later in this article. Half of the calculated amount is given intravenously during the first 8 hours, and the rest is given during the remaining 16 hours. If initial fluid administration is delayed, half of the calculated volume is to be completed by the end of the eighth hour after injury. Interestingly, even though current teaching is to give fluids aggressively to burn victims, some studies have suggested that this type of fluid management increases oxygen delivery to ischemic tissue, triggering free radicals that can further damage tissue.<sup>44</sup>

The fluid of choice for initial resuscitation is an isotonic crystalloid fluid, such as lactated Ringer's solution. The lactate replaces the chloride in the solution, decreasing the likelihood of hyperchloremic acidosis.<sup>8</sup> Many studies have explored the use of hypertonic solution (i.e., 3% saline solution) as an alternative for burn victims.<sup>22</sup> Some studies have shown that use of a hypertonic solution may decrease the extent of edema.<sup>45,46</sup> However, the outcomes of studies that compared hypertonic solution versus crystalloid solution are inconclusive and suggest the use of such hypertonic solution does not improve, and may worsen, outcomes.

During the second 24 hours of resuscitation, fluid administration should decrease. This premise is based on the belief that after 18 to 24 hours, if resuscitation is successful, capillary

integrity improves and therefore fluid requirements decrease.<sup>8</sup> The consensus formulas for fluid administration after the first 24 hours are listed in Table 2.

At this point in fluid management, the fluid of choice is a colloid formula, such as 5% albumin lactated Ringer's solution. In addition, electrolytes should be monitored closely.

The aforementioned formulas should be used in burn victims to achieve suggested endpoints of resuscitation. Infusion rates of fluids can be adjusted based on the resuscitation endpoints. (See Table 3.)<sup>8</sup> Resuscitation endpoints include stable vitals signs, normal mentation and sensorium. One important endpoint is maintaining adequate urine output, specifically 30-50 mL per hour in adults and 1mL/kg per hour in children.<sup>47,48</sup>

### Wound Care

Immediate wound care for burns is important for many reasons. Topical agents for less serious burns provide a means of pain control and decrease the rate of bacterial growth. Deeper burns may require surgical management or subsequent transfer to a burn center. How a burn wound is treated depends upon the depth of the wound. Overall, outpatient management of burns can be divided into the six Cs: clothing, cooling, cleaning, chemoprophylaxis, covering, and comforting.<sup>32</sup>

**Clothing.** Clear the patient's body of all materials that are hot or burned. In addition, clothing that appears to have come into contact with any chemicals also should be removed.

**Cooling.** Cold water has many purposes for burn wounds.

Applying gauze soaked with cold water stops the burning process, relieves pain, and removes chemicals from direct contact with skin. Use caution with cooling methods for patients with burns greater than 10% TBSA; they may cause hypothermia, especially in children. Cold water should be applied to the burn area for at least 10 minutes and a maximum of 20 minutes.<sup>49</sup>

**Cleaning.** Cleaning a wound is essential to prevent infection; however, the procedures can cause a great deal of pain. Local, regional, or systemic anesthesia should be induced before cleaning a wound; topical anesthesia and injection of anesthetic agents directly into the wound should be avoided. There is increasing support for using mild soap and tap water to wash burns.<sup>50-53</sup> Disinfectants, such as povidone-iodine solution or chlorhexidine gluconate solution, should be avoided because the agents will hamper the healing process. If there are any residues adhering to the wound, such as tar or asphalt, they should be removed with the aid of large amounts of bacitracin ointment applied during a period of many days.

Blister formation commonly is associated with burns. Intact blisters can allow re-epithelialization 40% faster than blisters that are aspirated or deroofed.<sup>13</sup> However, studies have shown that blister fluid contains proteins that increase the likelihood of sepsis by decreasing normal lymphocytic and neutrophilic function. Deroofing minor blisters is controversial and needs further research. Some study results have suggested that small-to-moderate-sized blisters be covered with occlusive dressing for the first 72 hours after injury. Large blisters or blisters over a joint may be aspirated with a needle and a syringe leaving the roof of the blis-

ter intact. After 72 hours, the blistered skin may be excised using aseptic technique.<sup>12</sup>

**Chemoprophylaxis.** Burns are considered to be tetanus-prone wounds. Active immunization against tetanus should be given to burn patients when tetanus status is in doubt. For patients who are not immunized or who have incomplete tetanus status, passive tetanus immunization is recommended.<sup>22</sup>

Burn wounds are more vulnerable to infections and, ultimately, sepsis. The common pathogens that cause infection in burn victims are *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Streptococcus pyogenes*, and other coliform bacilli.<sup>54</sup>

Topical antibiotics are an essential element in burn wound management. Classically, the medication of choice is silver sulfadiazine. It is a good selection for most burns, especially for deep partial-thickness burns, because it may permit wound healing without the need for a skin graft. Silver sulfadiazine cream should not be used on the face or in patients who are pregnant, newborns, or nursing mothers with children younger than 2 months because of the risk of sulfonamide kernicterus. In addition, silver sulfadiazine is contraindicated in patients with sulfa allergies. A cerium nitrate silver sulfadiazine cream also is commercially available. Cerium is a lanthanide metal that interacts with calcium, which is an important element of epidermal cell growth. A few studies suggest that cerium, in conjunction with silver sulfadiazine, can decrease local inflammation and sepsis.<sup>55</sup>

Bacitracin also can be used as a topical antibiotic for wound management. The advantage of using bacitracin over silver sulfadiazine is its lower cost. However, studies have not compared the efficacies of one topical antibiotic with another.

Biologic dressings, such as xenograft and allograft, also may be used to prevent wound contamination and fluid loss. These dressings are associated with lower infections rates and faster healing compared with silver sulfadiazine.<sup>39</sup> Biologic dressings need to be applied within 6 hours after burn injury. These dressings allow skin epithelialization and eventually will peel off as the skin heals.

Nonbiologic dressings provide a moist wound environment and fast healing. These dressings require fewer changes and induce less pain compared with topical antibiotics.<sup>39</sup> However, nonbiologic dressings need a bulky dressing that must be changed daily.

**Covering.** Superficial burns generally do not need wound dressings. Patients with this type of burn should be instructed to see their physician if blisters form. Also, a skin lubricant, such as aloe vera, can be applied to the burn wound.

Partial and full-thickness burns should be covered with sterile dressings after the wound is cleansed and a topical antibiotic is applied. Patients should be instructed to change dressings with recommended frequencies of twice a day to once a week.<sup>56</sup> At each dressing change, the wound should be cleaned gently, a topical antibiotic should be applied, and the wound re-dressed.

**Comforting.** A burn injury can be extremely painful. Patients with small burns may be instructed to take nonsteroidal anti-inflammatory drugs and acetaminophen. Nonsteroidal anti-inflammatory medications also decrease inflammation and

## Table 4. Burn Center Referral Criteria

### PATIENTS WHO MEET THE CRITERIA LISTED BELOW SHOULD BE REFERRED TO A BURN CENTER

- Partial-thickness and full-thickness burns greater than 10% of the total body surface area (TBSA) in patients younger than 10 years or older than 50 years
- Partial-thickness and full-thickness burns greater than 20% TBSA in other age groups
- Partial-thickness and full-thickness burns involving the face, eyes, ears, hands, feet, genitalia, or perineum, or those that involve skin overlying major joints
- Full-thickness burns greater than 5% BSA in any age group
- Electrical burns, including lightning injury. (Significant volumes of tissue beneath the surface may be injured and result in acute renal failure and other complications.)
- Significant chemical burns
- Inhalation injury
- Burn injury in patients with pre-existing illness that could complicate management, prolong recovery, or affect mortality
- Any burn patient in whom concomitant trauma poses an increased risk of morbidity or mortality may be treated initially in a trauma center until stable before transfer to a burn center.
- Children with burns seen in hospitals without qualified personnel or equipment for their care should be transferred to a burn center with these capabilities.
- Burn injury in patients who will require special social and emotional or long-term rehabilitative support, including cases involving suspected child abuse or neglect

*Adapted from Committee on Trauma, American College of Surgeons. Guidelines for the operation of burn units. Resources for Optimal Care of the Injured Patient 1999. 1998:55.*

edema and increase blood flow. Opioids taken orally may be added for pain control. For more painful burns, adults should be given morphine intravenously or intramuscularly. Aggressive pain control should be pursued with some adults requiring large doses of intravenous morphine.

### Special Considerations

**Pediatric Burns.** In general, the management of pediatric burns is similar to that of adult burns. Children have smaller and shorter airways, which can make intubation difficult, especially if the child has fast-developing edema. Children require larger amounts of fluid during resuscitation because they have larger insensible fluid losses. Pediatric patients also have a larger surface-to-mass ratio, which makes temperature control difficult. As always, in managing any emergency, the ABCs in the pediatric burned patient must be secured during initial resuscitation. To make up for fluid loss, including insensible fluid loss, the following formula can be used:

$$(5,000 \text{ mL/M}^2 \text{ BSA burned/24 hr}) + \\ (2,000 \text{ mL/M}^2 \text{ BSA nonburned/24 hr})^9$$

Half the calculated fluid is given during the first eight hours, and the rest is given during the next 16 hours. As in adults, pain

## Figure 5. Oral Commissure Burn



**Figure 5.** An oral commissure burn may occur when a child chews on a live electrical cord.

*Reprinted with permission from Stewart C. Electrical injuries. Ped Emerg Med Rep 2001;1:4.*

medication is fundamental in managing burns in children. A variety of pain medications can be used, such as nonsteroidal anti-inflammatory agents, opiates, benzodiazepines, neuroleptics, and dissociative drugs.<sup>57</sup> For example, a suggested starting dose of opiates (e.g., morphine) can be given in a dose of 0.1 to 0.2 mg/kg intravenously every 30 to 60 minutes as the patient's blood pressure and respiratory status tolerate. Some patients may require significantly higher doses.

**Geriatric Patients.** As the population ages, geriatric patients become increasingly common in EDs throughout the United States and other countries. Elderly females are more likely than elderly males to sustain a burn injury, and a lack of supervision is common.<sup>58</sup> The elderly are more likely to sustain a flame injury or a scald injury, and the majority of burn injuries have been reported to occur secondary to impaired judgment, mobility or both.<sup>58</sup> With increasing age, the survival decreases with reported survival rates in one series 86% in the 59-69 year old age group, 69% in the 70-79 year old age group, and 47% in patients older than 80 years.<sup>58</sup> In addition, the degree of concomitant chronic illness has a detrimental effect on survival.

**Inhalation Injury.** Inhalation injury is common in burn patients. Many people who sustain a burn injury were confined in a smoke-filled area and subsequently were exposed to large amounts of carbon monoxide. Usually 100% oxygen via face mask or intubation will reduce the half-life of carbon monoxide and is sufficient to treat an inhalational injury. However, hyperbaric oxygen also is a treatment option and usually is reserved for patients with carbon monoxide levels greater than 25%.<sup>59</sup> Other potential criteria for use of hyperbaric oxygen include patients with coma, transient loss of consciousness, ischemic electrocardiogram changes, focal neurologic deficits, and who are pregnant.<sup>60</sup>

**Electrical Injuries.** Electrical injuries also are considered a type of burn injury and often are managed at burn centers. Exposures to voltages of 200 to 1000 volts are considered moderate and are associated with local injury. Exposure to more than 1000

volts induces high-voltage injuries. They are associated with compartment syndromes, loss of consciousness, and myoglobinuria. Patients with these injuries must be monitored for cardiac arrhythmias for at least 24 to 72 hours.<sup>8</sup> In addition, delayed ophthalmologic and neurologic injuries can occur. Urine also must be monitored for myoglobinuria, and appropriate fluid must be administered.

**Oral Electrical Burns in Children.** Oral electrical burns in children usually are incurred by the sucking or chewing on the female end of a live extension cord or by biting through an electrical cord.<sup>61</sup> This type of injury is most common in children younger than 2 years, with a male predominance.<sup>61</sup> Electrolyte-rich saliva completes the circuit between the two electric poles. This results in an arc burn generating intense heat between 2500° and 3000° C that causes tissue necrosis.<sup>62,63</sup> The severity of the burn depends upon the length of contact, type of current, voltage, resistance of the tissue, and path of the electrical current through the body.<sup>62,63</sup> The low-voltage mechanism of this injury, usually resulting from household appliances consuming between 110 and 220 volts, leads to muscle contraction actually prolonging the length of exposure to the current.<sup>63</sup>

Oral electrical burn wounds have a central necrotic zone that appears gray or white. Surrounding this area, the tissue is edematous and raised. The most common area of involvement is the oral commissure. (See Figure 5.) The lower lip and cheek more likely are involved than the upper lip.<sup>64</sup> The heat of the electrical arc causes coagulation of tissues and thrombosis of blood vessels. As a result, bleeding is not very common in the acute phase of the injury. Edema of the surrounding tissues peaks within the first 24 hours.<sup>65</sup> Due to involvement of neural structures, the burns are usually painless and may result in deficits in sensory and motor function.<sup>61,62</sup>

In the acute care setting, these burn patients initially should be evaluated for systemic effects including arrhythmias and other injuries, which are unusual.<sup>65,66</sup> The wound should be irrigated, and topical antibiotics should be applied. Distinguishing between the viable and non-viable tissue is difficult in the acute phase, and debridement should not be attempted.<sup>67</sup> After an observation period in the ED, patients with isolated oral burn injury may be managed as outpatients with expeditious follow-up by a burn specialist and by a pediatric dentist.<sup>64</sup>

One of the most serious complications of oral electrical burns is delayed bleeding from the labial artery. This is most common 7 to 10 days post burn when swelling subsides and the eschar separates from the underlying tissue.<sup>63,66</sup> The patient's guardians should be warned of this possibility and should receive instruction on the proper application of pressure during transport back to the hospital should that occur. In the ED, the bleeding should be managed by applying direct pressure, if necessary, followed by applying epinephrine-soaked packing, and ultimately by suturing.<sup>62</sup>

Oral electric burns can be treated by different modalities. The use of oral splints to reduce wound contracture and to maintain position of the affected tissues during the healing process has been advocated.<sup>62</sup> The appliance is created by a pediatric dentist

and applied in the first few weeks post injury and commonly is worn for 6 to 12 months.<sup>62</sup> Surgical repair of tissue defects usually is delayed for at least 6 months after injury. At this time, the amount of scarring and the functional deficit can be better assessed.<sup>2</sup> The goals of reconstructive surgery are restoration of functionality and aesthetics.<sup>67</sup> These patients should be followed by a plastic surgeon well into their adolescence until fully grown.<sup>7</sup> Ultimately, most of these patients will have almost complete functional recovery of the mouth.<sup>64</sup>

## Disposition

Most small superficial burns and some partial-thickness burns can be managed on an outpatient basis. Adequate patient education on dressing changes, topical medications, and appropriate follow-up is necessary. A patient who does not have follow-up resources should be encouraged to return to the ED for wound checks.

Some patients with partial-thickness burns need to be transferred to a burn center. The American Burn Association has published criteria for patients needing specialized care that are listed in Table 4.<sup>68</sup> Refer questions regarding specific patients to a burn center physician for consultation.

## References

- Burn Incidence and Treatment in the U.S. 2000 Fact Sheet. American Burn Association. www.ameriburn.org. Accessed on 5/24/2004.
- Ryan CM, Schoenfeld DA, Thorpe WP, et al. Objective estimates of the probability of death from burn injuries. *N Engl J Med* 1998; 338:362-366.
- Danks RR. Burn management: A comprehensive review of the epidemiology and treatment of burn victims. *JEMS* 2003;28:118-141.
- Passaretti D, Billmire DA. Management of pediatric burns. *J Craniofac Surg* 2003;14:713-718.
- Barrow RE, Spies M, Barrow LN, et al. Influence of demographics and inhalation injury on burn mortality in children. *Burns* 2004; 30:72-77.
- U.S. Department of Health injury mortality: National summary on injury mortality date, 1984-1990. Washington, DC: U.S. Department of Health and Human Services;1993.
- Kao CC, Garner WL. Acute burns. *Plast Reconstr Surg* 2000;105; 2482-2493.
- Sheridan RL. Comprehensive treatment of burns. *Curr Probl Surg* 2001;38:641-756.
- Herndon DN, Spies M. Modern burn care. *Sem Pediatr Surg* 2001;10:28-31.
- Varghese TK, Kim AW, Kowal-Vern A, et al. Frequency of burn-trauma patients in an urban setting. *Arch Surg* 2003;138:1292-1296.
- Brandt CP, Yowler CJ, Fratianna RB. Burns with multiple trauma. *Am Surg* 2002;68:240-244.
- Flanagan M, Graham J. Should burn blisters be left intact or debrided? *J Wound Care* 2001;10:41-45.
- duKamp A. Deroofing minor burn blisters – What is the evidence? *Accid Emerg Nurs* 2001;9:217-221.
- Cumming J, Purdue GF, Hunt JL, et al. Objective estimates of the incidence and consequences of multiple organ dysfunction and sepsis after burn trauma. *J Trauma* 2001;50:510-515.
- Palmieri TL, Greenhalgh DG. Topical treatment of pediatric patients with burns. *Am J Clin Dermatol* 2002;3:529-534.
- Monafo WW. Current concepts: Initial management of burns. *N Engl J Med* 1996;335:1581-1586.
- Demling RH, Mazess RB, Witt RM, et al. The study of burn wound edema using dichromatic absorptiometry. *J Trauma* 1978;18: 124-128.
- Arturson G, Jonsson CE. Transcapillary transport after thermal injury. *Scand J Plast Reconstr Surg* 1979;13:29-38.
- Arturson G. Forty years in burns research – The post burn inflammatory response. *Burns* 2000;26:599-604.
- Arturson G. Pathophysiology of the burn wound and pharmacological treatment. The Rudi Hermans Lecture, 1995. *Burns* 1996;22: 255-274.
- Papp A, Uusaro A, Parviainen I, et al. Myocardial function and haemodynamics in extensive burn trauma: Evaluation by clinical signs, invasive monitoring, echocardiography and cytokine concentrations: A prospective clinical study. *Acta Anaesthesiol Scand* 2003;47:1257-1263.
- Sheridan RL. Burns. *Crit Care Med* 2002;30:500-514.
- Horton JW. Free radicals and lipid peroxidation mediated injury in burn trauma: The role of antioxidant therapy. *Toxicology* 2003;189: 75-88.
- Lee JO, Herndon DN. Modulation of the post-burn hypermetabolic state. Nestle Nutrition Workshop Series Clinical & Performance Program 2003;8:39-56.
- Demling RH, Seigne P. Metabolic management of patients with severe burns. *World J Surg* 2000;24:673-680.
- Hart DW, Wolf SE, Chinkes DL, et al. Determinants of skeletal muscle catabolism after severe burn. *Ann Surg* 2000;232:455-465.
- Hart DW, Wolf SE, Micak R, et al. Persistence of muscle catabolism after severe burn. *Surgery* 2000;128:312-319.
- Gore DC, Chinkes DL, Hart DW, et al. Hyperglycemia exacerbates muscle protein catabolism in burn-injured patients. *Crit Care Med* 2002;30:2438-2442.
- Herndon DN, Tompkins RG. Support of the metabolic response to burn injury. *Lancet* 2004;363:1895-1902.
- Barret JP, Jeschke MG, Herndon DN. Fatty infiltration of the liver in severely burned pediatric patients: Autopsy findings and clinical implication. *J Trauma* 2001;51:736-739.
- Rutan RL, Herndon DN. Growth delay in postburn pediatric patients. *Arch Surg* 1990;125:392-395.
- Morgan ED, Bledsoe SC, Barker J. Ambulatory management of burns. *Am Fam Phys* 2000;62:2015-26, 2029-30, 2032.
- Knaysi GA, Crikelair GF, Cosman B. The role of nines: Its history and accuracy. *Plast Reconstr Surg* 1968;41:560-563.
- Lund CC, Browder NC. Estimation of areas of burns. *Surg Gynecol Obstet* 1944;79:352-358.
- Perry RJ, Moore CA, Morgan BD, et al. Determining the approximate area of a burn: an inconsistency investigated and re-evaluated. *BMJ* 1996;312:1338.
- Livingston EH, Lee S. Percentage of burned body surface area determination in obese and nonobese patients. *J Surg Res* 2000;

- 91:106-110.
37. Palmieri TL, Greenhalgh DG. Topical treatment of pediatric patients with burns. *Am J Clin Dermatol* 2002;3:529-534.
  38. Harulow S. Burn wounds: Assessment and first aid treatment. *Aust Nurs J* 2000;7:1-4.
  39. Johnson RM, Richard R. Partial-thickness burns: identification and management. *Adv Skin Wound Care* 2003;16:178-189.
  40. Watts AM, Tyler MP, Perry ME, et al. Burn depth and its histological measurement. *Burns* 2001;27:154-160.
  41. Pape SA, Skouras CA, Byrne PO. An audit of the use of laser Doppler imaging (LDI) in the assessment of burns of intermediate depth. *Burns* 2001;27:233-239.
  42. Holland AJA, Martin HCO, Cass DT. Laser Doppler imaging prediction of burn wound outcome in children. *Burns* 2002;28:11-17.
  43. American College of Surgeons Committee on Trauma. Resources for Optimal Care of the Injured Patient. Chicago: American College of Surgeons, 1993.
  44. Horton JW. Free radicals and lipid peroxidation mediated injury in burn trauma: The role of antioxidant therapy. *Toxicology* 2003;189:75-88.
  45. Brown MD. Hypertonic versus isotonic crystalloid for fluid resuscitation in critically ill patients. *Ann Emerg Med* 2002;20:113-114.
  46. Bunn F, Roberts I, Tasker R, et al. Hypertonic versus isotonic crystalloid for fluid resuscitation in critically ill patients (Cochrane Review). In: The Cochrane Library. Issue 4. Oxford, United Kingdom, Update Software Ltd; 2002.
  47. Purdue GF, Hunt JL, Burris AM. Pediatric surgical emergencies: Pediatric burn care. *Clin Pediatr Emerg Med* 2003;3:76-82.
  48. Warden GD. Burn shock resuscitation. *World J Surg* 1992;16:16-23.
  49. Harulow S. Burn wounds: Assessment and first aid treatment. *Aust Nurs J* 2000;7:1-4.
  50. Mertens DM, Jenkins ME, Warden GD. Outpatient burn management. *Nurs Clin North Am* 1997;32:343-64.
  51. Baxter CR. Management of burn wounds. *Dermatol Clin* 1993;11:709-714.
  52. Waitzman AA, Negligan PC. How to manage burns in primary care. *Can Fam Physician* 1993;39:2394-2400.
  53. Greenhalgh DG. The healing of burn wounds. *Dermatol Nurs* 1996;8:13-23.
  54. Edward-Jones V, Greenwood J. What's new in burn microbiology. *Burns* 2003;29:15-24.
  55. Lansdown A, Myers S, Ciarice J, Sullivan P. A reappraisal of the

- role of cerium in burn wound management. *J Wound Care* 2003;12:113-120.
56. Hartford CE. Care of outpatient burns. In: Herndon DN, ed. *Total Burn Care*. Philadelphia:Saunders, 1996:71-80.
57. Stoddard FJ, Sheridan RL, Saxe GN, et al. Treatment of pain in acutely burned children. *J Burn Care Rehabil* 2002;23:135-156.
58. McGill V, Kowal-Ven A, Gamelli RL. *Arch Surg* 2000;135:320-325.
59. Heimbach D. What's new in general surgery: Burns and metabolism. *J Am Coll Surg* 2002;194:156-164.
60. Van Hoesen K. Hyperbaric oxygen therapy. In: Rosen P, et al, eds. *Emergency Medicine: Concepts and Clinical Practice*. 2nd ed. St. Louis, Mo: Mosby-Year Book; 1998:1032-1042.
61. Milano M. Oral electrical and thermal burns in children: Review and report of case. *J Dent Child* 1999;66:116-119.
62. Donly KJ, Nowak AJ. Oral electrical burns: Etiology, manifestations, and treatment. *Gen Dent* 1988;36:103-107.
63. Canady JW, Thompson SA, Bardach J. Oral commissure burns in children. *Plast Reconstr Surg* 1996;97:738-744.
64. Rai J, Jeschke MG, Barrow RE, et al. Electrical injuries: A 30-year review. *J Trauma* 1999;46:933-936.
65. Banks K, Merlino PG. Minor oral injuries in children. *Mount Sinai J Med* 1998; 65:333-342.
66. Zubair M, Besner GE. Pediatric electrical burns:management strategies. *Burns* 1997;23:413-420.
67. Thomas SS. Electrical burns of the mouth: still searching for an answer. *Burns* 1996;22:137-140.
68. Committee on Trauma, American College of Surgeons. Guidelines for the operation of burn units. Resources for Optimal Care of the Injured Patient 1999. 1998:55.

## CME/CE Questions

1. Burns can occur by several different mechanisms. What are the three most common types of burn injuries?
  - A. chemical, scald, and electrical
  - B. electrical, thermal, and scald
  - C. scald, contact, and thermal
  - D. radiation, chemical, and contact

## CE/CME Objectives

Upon completing this program, the participants will be able to:

- a) recognize or increase suspicion for traumatic injuries that present to the emergency department;
- b) describe the various modalities used to identify different traumatic conditions covered in the newsletter;
- c) describe how to correctly and quickly stabilize, and then to manage patients with the particular condition covered in the newsletter; and
- d) identify both likely and rare complications that may occur with traumatic injuries.

## CE/CME Instructions

Physicians and nurses participate in this continuing medical education/continuing education program by reading the article, using the provided references for further research, and studying the questions at the end of the article. Participants should select what they believe to be the correct answers, then refer to the list of correct answers to test their knowledge. To clarify confusion surrounding any questions answered incorrectly, please consult the source material. **After completing this activity, you must complete the evaluation form provided and return it in the reply envelope provided in order to receive a certificate of completion.** When your evaluation is received, a certificate will be mailed to you.

2. Burn wounds consist of three distinct zones of tissue damage: the coagulation zone, the stasis zone, and the hyperemic zone. Which of the following is an accurate description of the stasis zone?
  - A. Tissue irreversibly destroyed by heat, which cannot recover
  - B. Damaged tissue with decreased perfusion and the potential for recovery
  - C. Intermediate damage to tissue, with no chance of recovery
  - D. Minimal damage to tissue, with the chance of spontaneous recovery
  
3. The depth of burn injury commonly has been classified in terms of first, second, and third degree. However, these categories have been replaced recently by the terms superficial, partial thickness, and full thickness. Which statement below best describes partial-thickness burns?
  - A. They only affect the epidermis, have no blister formation, heal in 3 to 7 days, and do not lead to scarring.
  - B. They involve the underlying tissue, such as muscle, bone, or fascia; are extremely painful; and lead to systemic disease.
  - C. They destroy the epidermis and dermis, extend into the subcutaneous tissue, and are insensate secondary to the destruction of nerves.
  - D. They include the subcategories of superficial and deep and extend through the epidermis into the superficial and deeper layers of the dermis, respectively.
  
4. Which of the following statements accurately describes why it is important to assess the total body surface area (TBSA) of the burned patient?
  - A. It is used to estimate fluid resuscitation requirements and to assess the risk of death.
  - B. It is used to assess the patient's need for surgical intervention and stabilization.
  - C. It is used to determine the probability of scar development after the wound heals.
  - D. It uses the Rule of Nines, equally accurately in adults and children, to estimate the extent of burn injury.
  
5. To evaluate the burn patient for carbon monoxide poisoning, one must complete which of the following tests?
  - A. serial chest radiographs and pulse oximetry
  - B. measurement of arterial or venous blood carboxyhemoglobin level
  - C. complete blood count, chemistries, and coagulation studies
  - D. arterial blood gas measurements and liberal CT scans
  
6. The American Burn Association (ABA) recommends using which of the following for burns greater than 15% TBSA during the first 24 hours?
  - A. The Resuscitation Consensus Formula
  - B. Hypertonic solution (i.e., 3% saline solution)
  - C. The Parkland Formula as the gold standard for adults
  - D. A combination of colloid formula and isotonic crystalloid fluid
  
7. What is the essential reason to clean burn wounds?
  - A. To ensure decreased blister formation
  - B. To immediately remove adherents such as tar and asphalt
  - C. To prevent the separation of the epidermis from the dermis
  - D. To prevent future infection of the burn wound
  
8. Which are the most common pathogens that cause burn wound infections?
  - A. *Staphylococcus aureus*, *Streptococcus pneumoniae*, and *Clostridium perfringens*
  - B. *Pseudomonas aeruginosa*, *Streptococcus pyogenes*, and *Staphylococcus aureus*
  - C. *Streptococcus pyogenes*, *Clostridium perfringens*, and *Staphylococcus epidermidis*
  - D. *Staphylococcus epidermidis*, *Pseudomonas aeruginosa*, and *Streptococcus pneumoniae*
  
9. Topical antibiotics are an essential element in burn wound management. Classically, silver sulfadiazine is the medication of choice and can be used in which of the following patients?
  - A. A pregnant woman with  $\geq 15\%$  TBSA burns
  - B. A patient who has burn wounds to the face
  - C. A patient with deep partial-thickness burns and no sulfa allergies
  - D. A newborn with superficial extremity burn wounds only
  
10. Management of burns in children is similar to that of burns in adults with a few exceptions. Which of the following statements is true?
  - A. Children require larger amounts of fluid during resuscitation because they have larger insensible losses.
  - B. Pediatric patients have smaller surface-to-mass ratio, which makes their temperature easier to control.
  - C. Children have a smaller and shorter airway, which makes intubation easier and right main stem intubation less likely.
  - D. Pain medication is not as important in managing pediatric burns, and nonsteroidal anti-inflammatory agents usually are adequate.

**Answer Key:**

1. C
2. B
3. D
4. A
5. B
6. C
7. D
8. B
9. C
10. A